

Creepers: Real Quadratic Orders with Large Class Number

Roger Patterson,
B.Sc(Hons) Monash

Department of Mathematics
Division of Information and Communication Sciences
Macquarie University
Sydney, Australia

November, 2003

Acknowledgements

I am undoubtedly indebted to my two supervisors, Alf van der Poorten and Hugh Williams. Most students would feel lucky to have one good supervisor whereas I had the privilege of having two excellent mathematicians to guide me. Their help, wisdom and infinite patience has been invaluable, and I am very grateful for it.

The Mathematics Department at Macquarie University has provided a pleasant and enjoyable environment for the past few years. I would also like to thank both the University of Calgary and the University of Manitoba for their warm and generous hospitality during my visits to Canada. Those visits were made possible by an ARC IREX grant, and these visits were vital in the development of the thesis. Also, I owe a great debt to my Canadian hosts, Lynne Romuld in Winnipeg and Kris and Gardy Vasudevan in Calgary, who took me in and made me feel welcome.

Last, but not least I would also like to thank all my family and friends, in particular my lovely wife Cath for all her patience and inspiration.

Certificate

This thesis is presented for the degree of Doctor of Philosophy at Macquarie University. It has not been submitted for a higher degree at any other university or institution.

Roger Patterson

TABLE OF CONTENTS

Overview	1
Chapter I. Introduction	5
1. Quadratic forms	5
2. Continued fractions	6
3. Units and orders	13
4. Ideals, infrastructure and composition	17
5. Function fields	21
Chapter II. History of sleepers, beepers and creepers	27
6. Creepers and beepers	28
Chapter III. Sleepers	33
7. Sleepers from curves of genus 1	34
8. Finding exceptional function fields	39
9. More sleepers	45
Chapter IV. Elementary aspects of creepers	47
10. Easy Creepers	47
11. When l is negative	52
12. When both l and m are not units	55
Chapter V. Determining creepers	59
13. Preliminary Results	59
14. Basic observations on creepers	62
15. Independent ideals in creepers	66
16. Part of the continued fraction expansion of ω_n	71
17. The length of the continued fraction expansion of A_{2i}/B_{2i}	84
Chapter VI. Additional remarks about creepers	87
18. Don't be $\square$	87
19. The proper way to handle square factors	90
20. A composition approach	92
21. The fundamental unit of creepers	95
22. Function field creepers	98
Chapter VII. The general expansion of a creeper	101
23. Determination of the specific elements in the expansion	102
24. Determining the number of partial quotients	110
25. Finding an element of norm 1	113
26. Returning to the regular continued fraction expansion	116
Chapter VIII. Higher order creepers	119
27. More creepers	119
28. From torsion to creepers	125
29. Unusual representations of creepers	130
30. Large torsion creepers	131
31. Jeepers?	133
Conclusion	135

Tables	137
Sleepers	137
Easy Kreepers	139
When l is negative	141
When both l and m are not units	142
Function Field Kreepers	149
Higher Creepers	150
Jeepers	156
Bibliography	163

Overview

MOTIVATION

One of the original problems of modern number theory is the determination of class numbers of quadratic fields. This was first investigated by Gauss at the conclusion of the eighteenth century following earlier work of Fermat, Lagrange and others. It was at this time that Gauss proposed that there are infinitely many real quadratic fields whose class number is 1. This still unsolved problem motivated Shanks in the 1960s to look at tables of class numbers in order to establish any irregularities. He discovered a family of fields whose class numbers grew abnormally large in a nontrivial manner. This sequence has come to be known as *Shanks' sequence* and is given by

$$S_n = (2^n + 1) + 4 \cdot 2^n. \quad (0.1)$$

Shanks, as well as Gauss, knew of the correspondence between binary quadratic forms and continued fractions of quadratic irrationals, and so he was able to express this phenomenon in the language of continued fractions. Namely, that the period length of the continued fraction of $\sqrt{S_n}$ is abnormally short.

Generally, one has no idea a priori how the continued fraction expansion of $\sqrt{D}$ will behave. The so-called “Cohen-Lenstra” heuristics [11] suggest that 75% of all prime discriminants have a class number of 1. In terms of continued fractions, very roughly, this says that the continued fraction for $\sqrt{D}$ would have a period length of about $\sqrt{D}$ in most cases. However, it is notoriously difficult to predict the continued fraction expansion of $\sqrt{D}$. Nevertheless, there are instances where one can explicitly write the expansion. For example:

$$\sqrt{D} = \begin{cases} [n, \overline{2n}] & \text{if } D = n^2 + 1 \\ [n, \frac{n-1}{2}, 1, 1, \frac{n-1}{2}, 2n] & \text{if } D = n^2 + 4 \text{ and } n \equiv 1 \pmod{2} \end{cases}$$

These examples were generalized by Schinzel, who proved that $D_n = an^2 + bn + c$ has a bounded period length if and only if $a = \square$ and $\Delta \mid 4(2a, b)^2$. (Here $\Delta = b^2 - 4ac$).

Following Shanks' discovery (which had actually been found earlier by Nyberg, yet was neglected), others began to uncover further examples of sequences of real quadratic fields with large class numbers and short period lengths. This problem intrigued Kaplansky and led him to conjecture that all such families had been given in a paper of Hugh Williams. It was this conjecture of Kaplansky's which led to this work; namely that of establishing all *kreepers*, the name which we have given to such unusual sequences of real quadratic fields.

BREAKDOWN OF PARTS

The first chapter of this thesis is an introduction which establishes the notation used throughout; this is relevant to the notation for continued fractions, for which there is no strict standard. Since continued fractions will feature throughout, a review of some facts regarding them is given. We also show how continued fractions can be related to certain matrix products, and while this correspondence will only explicitly occur occasionally, it underlies the thinking in this thesis. We also examine the well known correspondence between indefinite binary quadratic forms, continued fractions of quadratic irrationals and ideals of real quadratic fields. Lastly, there is a brief overview of function fields.

Chapter II gives a history of the search for real quadratic fields with large class numbers. This is, in effect, the same thing as determining continued fraction expansions of quadratic irrationals whose period length is quite small. The size of

this “smallness” partitions the known examples into various sets: that of sleepers, beepers and creepers (with thanks to Kaplansky and Van der Poorten for the fine nomenclature). Sleepers have bounded period length, beepers have unbounded period length but a regulator of size only $O(\log(D_n))$, and creepers, which are polynomially parametrized, have a period length of $an + b$ and a regulator of order n^2 . A copy of Kaplansky’s letter is included as an appendix, like him we are primarily interested in creepers.

Chapter III gives a slight detour into the realm of sleepers. The quadratic instances of sleepers were fully determined by Schinzel back in the 1960s. The coupling of Mazur’s result, which gives all possible instances of torsion on elliptic curves, as well as another theorem of Schinzel’s allows us to detail the cases of sleepers for quartics. However, unlike the quadratic case this survey is not comprehensive; that is we don’t give every instance of a quartic sleeper, which would be feasible given the results in this part, but not of overly great importance. It is determined which quartics $f(X)$, where $\sqrt{f(X)}$ is periodic, yield numerical sleepers $\sqrt{f(x)}$ for x in some congruence class. Also included in this section is a review of the methods known for finding function fields with a nontrivial unit. In particular, Flynn determined an infinite sequence of hyperelliptic curves,

$$Y^2 = \left(\frac{1}{2} (X^{g+1} - t(X-1)^g - X^g(X-1)) \right)^2 - tX^g(X-1)^{g+1} \quad (0.2)$$

which possess a torsion divisor of order $2g^2 + 2g + 1$. These results become relevant in Chapter VIII during the investigation of higher order creepers.

Chapter IV investigates some of the simpler aspects of creepers, mainly through some explicit examples. Our interest will focus on new examples, but not, at this point, on the full generality. This way we obtain a gentle extension of the earlier known creepers. It is here that our procedure for handling creepers is introduced.

Chapter V is then where the hard work is done. This chapter establishes the necessary form a creeper must take as well as certain divisibility requirements. The sequence of discriminants given by a creeper are $\{D_n\}_{n \in I}$, for some infinite set I , where

$$D_n = (c/d)^2 \left[(qrx^n + (mz^2x^k - ly^2)/q)^2 + 4ly^2rx^n \right] \quad (0.3)$$

and

$$\begin{aligned} m, l, r \text{ are squarefree, } \quad c^2 r l y^2 m z^2 \mid d^2 D_n, \quad (qr, x) = 1, \\ (qrx, m z l y) = 1, \quad (yl, m z) = 1. \end{aligned}$$

The first few sections establish some easy properties as well as extending a theorem of Yamamoto which allows us to determine a form for creepers. It is in showing that this particular form can then be reduced to the instance of (0.3) that we run into difficulties. The verification of all the required results is elementary, but unfortunately tedious.

Chapter VI covers material which was not dealt with in Chapter IV. The effect of the square factors y^2 and z^2 is the first issue considered. We detail both the expansions of D_n and $D_n/(zy)^2$. From this we discover that the expansion of $D_n/(zy)^2$ is much simpler. However, in Chapter VII we will have to expand the more complicated D_n since there is no efficient way of determining the expansion of $\sqrt{D_n}$ from that of $\sqrt{D_n}/zy$. The continued fraction algorithm is often referred to as taking “Baby-steps”, in Shanks’ language. Composition, on the other hand, is called “Giant-steps” since it involves taking many Baby-steps in one calculation. Up to this point, we have only used Baby-steps to describe creepers. But here, we investigate creepers from the Giant-step viewpoint. That is, by appropriately composing certain ideals we determine a unit. Although we don’t do it, with more work it would be possible to use this approach to also show that the number of

Baby-steps would have to be linear in n . This Giant-step approach is similar to how some of the other authors tackled the earlier examples of creepers.

Since creepers have a small unit, we explicitly give this unit, using methods distinct from those in Chapter V. As a concluding note for this part, we examine function field creepers. These are sequences of polynomials, $\{f(X)\}$ of degree $2g+2$, such that the period length of $\sqrt{f(X)}$ is linear in g and the regulator is of order g^2 . Our interest is primarily numerical, consequently the coverage of these functional matters will be brief.

Chapter VII confirms that the form found in Chapter V, namely (0.3) does in fact yield a creeper. Much of the required work has already been completed in Chapter V. All that is required at this point is to find an element of norm 1 and to show that this element must occur after $an + b$ steps of the continued fraction expansion. This procedure also explicitly constructs the fundamental unit, in a manner distinct from that used in Chapter VI.

Chapter VIII further highlights the connection between function fields and number fields. The motivation for this part is to find higher order creepers. In other words, we seek sequences of real quadratic fields with large class numbers whose discriminants are given by polynomial families in x^n with a degree greater than 2. We find that the earlier results regarding torsion of elliptic and hyperelliptic curves become useful. More precisely, by appropriately manipulating these curves we are able to determine instances of creepers which appear to have little resemblance to creepers. This connection is further demonstrated by showing that Shanks' sequence of quadratic fields (0.1) is in fact just a specialisation of Flynn's sequence of hyperelliptic curves (0.2), and moreover that each of them is constructed in an identical manner. The results in this chapter are less comprehensive, but they raise interesting questions about the nature of higher order creepers.

We conclude by summarising what has been shown throughout the thesis and what the future directions for this work are. The final section is just a collection of tables which are cited throughout the thesis. There are some detailed expansions of creepers, which assist in the understanding of Chapters IV and VI. The tables of higher order creepers are interesting, if only to indicate the complex nature of some creepers.

Chapter I. Introduction

Modern number theory essentially began with the work of Fermat in the 17th Century. Results like

$$\text{for an odd prime } p, \quad p = a^2 + b^2 \iff p \equiv 1 \pmod{4} \quad a, b \in \mathbb{Z}$$

led following generations of mathematicians to develop the theory of quadratic forms. This development led to the realisation that the property of unique factorization, which is so important and natural in the rational integers, may fail to hold in an algebraic number field. The class number is, in a sense, a measure of how bad this failure is (i.e. class number 1 indicates that unique factorization holds).

Much of the early headway was made by Gauss. However, his theory of binary quadratic forms was deemed by some to be superseded with the introduction of ideals by Dedekind. It had already been known that the continued fraction expansion of $\sqrt{D}$ gives much information about the forms of discriminant D . So one now had a connection between binary quadratic forms, ideals and continued fractions.

1. QUADRATIC FORMS

A binary quadratic form

$$f(x, y) = ax^2 + bxy + cy^2 \tag{1.1}$$

is said to have *discriminant* $b^2 - 4ac = D$ and if $\gcd(a, b, c) = 1$ the form is called *primitive*. If $D < 0$ and $a > 0$ ($a < 0$) the form is called positive (negative) definite since these are precisely the values f can then take¹. On the other hand, if $D > 0$ the form is called indefinite. Henceforth, our interest will purely be on positive discriminants. A discriminant D is called *fundamental* if $D \equiv 1 \pmod{4}$ and D is squarefree or if $d \equiv 2, 3 \pmod{4}$ is squarefree and $D = 4d$. A shorthand notation frequently used to represent the quadratic form (1.1) is (a, b, c) .

Detailed explanations of quadratic forms can be found in [12], [13], [17], as well as the ancient originals [20] and [14].

Two forms $f_1(x, y) = a_1x^2 + b_1xy + c_1y^2$ and $f_2(X, Y) = a_2X^2 + b_2XY + c_2Y^2$ are called *equivalent* if they have the same discriminant and there exists a unimodular transformation between the two. That is there exists $\alpha, \beta, \gamma, \delta \in \mathbb{Z}$ such that

$$\alpha\delta - \beta\gamma = \pm 1 \quad \text{and} \quad \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \begin{pmatrix} X \\ Y \end{pmatrix}$$

which transforms f_1 into f_2 . If the transformation satisfies $\alpha\delta - \beta\gamma = 1$ then the forms are *properly equivalent*, otherwise they are *improperly equivalent*. We will only be interested in proper equivalence, so henceforth the phrase equivalence will mean proper equivalence.

This is an equivalence relation and the point is that two equivalent forms are going to be assuming the same values so there really is no point in distinguishing between the two. As an example, $33x^2 + 28xy + 6y^2$, in shorthand $(33, 28, 6)$, is equivalent to $(1, 0, 2)$ via the transformation $\begin{pmatrix} -1 & 2 \\ 2 & -5 \end{pmatrix}$.

Another example is the so-called Pell equation,

$$f(x, y) = x^2 - Dy^2$$

which is equivalent to

$$g(X, Y) = X^2 + 2XY - (D - 1)Y^2$$

¹The relation $4af(x, y) = (2ax + by)^2 - Dy^2$ may help.

under the transformation $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$. But if $D \equiv 1 \pmod{4}$, then $g(X, Y)$ is not equivalent to

$$h(x, y) = x^2 + xy + \frac{1-D}{4}y^2,$$

since the transformation $X \mapsto x, Y \mapsto y/2$ is not unimodular and even worse g and h don't have the same discriminant. Nonetheless, for a fundamental discriminant D we can define the *principal form*:

$$f(x, y) = \begin{cases} x^2 - \frac{D}{4}y^2 & \text{if } D \equiv 0 \pmod{4} \\ x^2 + xy + \frac{1-D}{4}y^2 & \text{if } D \equiv 1 \pmod{4}. \end{cases}$$

Fermat was able to observe that

$$(x^2 + y^2)(u^2 + v^2) = X^2 + Y^2 \quad \text{via } X = xu - yv \text{ and } Y = xv + uy \quad (1.2)$$

which says that the product of two sums of two squares is again a sum of two squares. Formulas like these were handy in proving things like:

$$\begin{aligned} p &= x^2 + 5y^2 & \text{if and only if } p &\equiv 1, 9 \pmod{20}, \\ p &= 2x^2 + 2xy + 3y^2 & \text{if and only if } p &\equiv 3, 7 \pmod{20}, \end{aligned}$$

which is assisted by knowing

$$\begin{aligned} (2x^2 + 2xy + 3y^2)(2u^2 + 2uv + 3v^2) \\ = (2xu + xv + uy - 2vy)^2 + 5(xv + yu + yv)^2. \end{aligned}$$

Consequently, one is led to consider what is the product of two quadratic forms? It turns out that the product is well-defined and the operation is called *composition*. There is a formula for determining a composite, which we shall give later. Moreover, for $f(x, y)$ the *conjugate* form $f(x, -y)$ acts an inverse. For a given discriminant, the set of primitive forms modulo equivalent forms is a group, which is called the *form class group*; the order of which is called the *form class number*.

Naturally, there are infinitely many forms of any given discriminant, but we are really interested in the number of *inequivalent* forms. To assist in this calculation, a form $ax^2 + bxy + cy^2$ of discriminant $D > 0$ is called *reduced* if it satisfies

$$\frac{b + \sqrt{D}}{2|a|} > 1 \quad \text{and} \quad -1 < \frac{b - \sqrt{D}}{2|a|} < 0.$$

This is an appropriate definition because every form is equivalent to a reduced form. A reduced form (a, b, c) must satisfy

$$0 < b < \sqrt{D}, \quad \sqrt{D} - b < 2|a| < \sqrt{D} + b, \quad \sqrt{D} - b < 2|c| < \sqrt{D} + b.$$

Thus the set of reduced forms for any given discriminant is finite. However, a reduced form is not necessarily unique; a form may be equivalent to many reduced forms. To determine the class number one needs to know the reduced forms and which ones are equivalent to each other. A method of establishing this is through the use of continued fractions.

2. CONTINUED FRACTIONS

There are many references for this section, but frequently there is a clash in notation. Our notation predominately follows [54], [71], [79]. The bible of all things continued fractionally is [58], as well as [29].

A continued fraction is an expression which looks like

$$\alpha = a_0 + \frac{b_0}{a_1 + \frac{b_1}{a_2 + \frac{b_2}{a_3 + \ddots}}}$$

A simple continued fraction is one where all the b_i 's are 1. We will only be interested in simple continued fractions and henceforth drop the adjective “simple”. One immediately recognizes that it is far more convenient to represent a continued fraction as

$$\alpha = [a_0, a_1, a_2, \dots]$$

where the a_i are called the *partial quotients*. A finite continued fraction is one with only finitely many partial quotients. Of course, these are precisely the rational numbers; while an infinite continued fraction has infinitely many partial quotients and must represent an irrational. Truncations of a continued fraction,

$$[a_0, a_1] = \frac{x_1}{y_1}, \quad [a_0, a_1, a_2] = \frac{x_2}{y_2}, \quad \dots,$$

are known as its *convergents*. The individual terms, x_h and y_h are known as the *continuants* since they continue the continued fraction via (2.2) given below. The point is that the convergents x_h/y_h provide good rational approximations to the quantity α .

Thankfully, there is an easier method to compute the x_h, y_h . It is,

$$\begin{pmatrix} a_0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} a_1 & 1 \\ 1 & 0 \end{pmatrix} \cdots \begin{pmatrix} a_h & 1 \\ 1 & 0 \end{pmatrix} = \begin{pmatrix} x_h & x_{h-1} \\ y_h & y_{h-1} \end{pmatrix}, \quad (2.1)$$

which is easily confirmed via induction. Also, since we have

$$\begin{pmatrix} x_h & x_{h-1} \\ y_h & y_{h-1} \end{pmatrix} \begin{pmatrix} a_{h+1} & 1 \\ 1 & 0 \end{pmatrix} = \begin{pmatrix} x_{h+1} & x_h \\ y_{h+1} & y_h \end{pmatrix},$$

the continuants are given by the recurrence relations:

$$x_{h+1} = a_{h+1} x_h + x_{h-1}, \quad (2.2a)$$

$$y_{h+1} = a_{h+1} y_h + y_{h-1}, \quad (2.2b)$$

with the definition that

$$\begin{pmatrix} x_{-1} & x_{-2} \\ y_{-1} & y_{-2} \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

Various formulae can now be obtained regarding continued fractions. For example, by transposing (2.1), we find

$$\begin{pmatrix} x_h & y_h \\ x_{h-1} & y_{h-1} \end{pmatrix} = \begin{pmatrix} a_h & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} a_{h-1} & 1 \\ 1 & 0 \end{pmatrix} \cdots \begin{pmatrix} a_0 & 1 \\ 1 & 0 \end{pmatrix} \longleftrightarrow [a_h, a_{h-1}, \dots, a_0],$$

where the ‘ $\longleftrightarrow$ ’ notation is used to represent the correspondence between continued fraction expansions and matrix products. Then,

$$[a_h, a_{h-1}, \dots, a_0] = \frac{x_h}{x_{h-1}}$$

and

$$[a_h, a_{h-1}, \dots, a_1] = \frac{y_h}{y_{h-1}}.$$

In other words, two consecutive denominators or numerators contain all the prior information. It is also useful to sometimes drop the subscripts and denote x_h, y_h by x, y and x_{h-1}, y_{h-1} by x', y' .

Moreover, if

$$\alpha = [a_0, a_1, \dots, a_h, \alpha_{h+1}] \longleftrightarrow \begin{pmatrix} x_h & x_{h-1} \\ y_h & y_{h-1} \end{pmatrix} \begin{pmatrix} \alpha_{h+1} & 1 \\ 1 & 0 \end{pmatrix} \quad (2.3)$$

then we have

$$\alpha = \frac{x_h \alpha_{h+1} + x_{h-1}}{y_h \alpha_{h+1} + y_{h-1}}, \text{ which gives } \alpha_{h+1} = -\frac{y_{h-1} \alpha - x_{h-1}}{y_h \alpha - x_h}.$$

The α_{h+1} are called *complete quotients*.

These results so far have been formal but traditionally one likes to make the partial quotients positive integers. The *regular* continued fraction expansion of α is determined by the following algorithm,

$$a_0 = [\alpha], \alpha_0 = \alpha, \quad \alpha_{i+1} = \frac{1}{\alpha_i - a_i}, \quad a_{i+1} = [\alpha_{i+1}], \quad i = 0, 1, 2, \dots$$

so $\alpha = [a_0, a_1, \dots, a_h, \alpha_{h+1}]$. This procedure terminates when $\alpha_i \in \mathbb{Z}$.

The following results (each of which is easily proved by the matrix correspondence) allow an easy taming of nonpositive partial quotients:

$$\begin{aligned} [\dots, a, 0, b, \dots] &= [\dots, a + b, \dots], \\ [\dots, a, -b, c, \dots] &= [\dots, a, 0, -1, 1, -1, 0, b, -c, \dots] \quad \text{if } b \neq 1, \\ [\dots, a, -1, b, \dots] &= [\dots, a - 2, 1, b - 2, \dots]. \end{aligned}$$

For example, the first result is nothing other than

$$\begin{pmatrix} a & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} b & 1 \\ 1 & 0 \end{pmatrix} = \begin{pmatrix} a + b & 1 \\ 1 & 0 \end{pmatrix}.$$

The ease of these transformations is important to us since we will frequently encounter irregular continued fractions. It is also worth noting that there is always an odd as well as even length continued fraction expansion for any rational number. Merely observe that

$$[a, \dots, y, z] = [a, \dots, y, z - 1, 1];$$

thus $[a, \dots, x, y + 1]$ if $z = 1$.

Fractional partial quotients are not so easy to deal with. While we can guarantee the removal of a nonpositive integer partial quotient the same does not hold for fractional partial quotients. However, the following few results do provide some assistance. Multiplication can be accomplished via

$$x[a, b, c, d, \dots] = [ax, b/x, xc, d/x, \dots]$$

which leads to the Folding lemma,

Lemma 1 (Folding Lemma). *Let $x/y = [a_0, a_1, \dots, a_h]$ and denote $a_1, \dots, a_h$ by $\vec{w}$ ($\vec{w}$ corresponds to $a_h, \dots, a_1$). Then*

$$\frac{x}{y} + \frac{(-1)^h}{cy^2} = [a_0, \vec{w}, c - y'/y] = [a_0, \vec{w}, c, -\overleftarrow{w}].$$

This result, due to Mendés France [44], is more than just a novelty. Besides our use of it here, in [76] it is used to rediscover the symmetry formulas.

A result which will be pivotal to our expansions later on is the following,

Lemma 2. *If $x/y = [a_0, \vec{w}]$ where $\vec{w}$ is defined as above then*

$$x/y + \gamma = \left[a_0, \vec{w}, \frac{(-1)^h}{\gamma y^2} - \frac{b}{y} \right]$$

where b is equal to $(-1)^{h+1}/x$ modulo y .

Proof. Using the Folding lemma we obtain,

$$\frac{x}{y} + \gamma = \frac{x}{y} + \frac{(-1)^h}{y^2} (-1)^h \gamma y^2 = \left[a_0, \vec{w}, \frac{(-1)^h}{\gamma y^2} - \frac{b}{y} \right]$$

where b satisfies $xb - cy = (-1)^{h+1}$ which implies $b \equiv (-1)^{h+1}/x \pmod{y}$. ■

This lemma allows us to systematically remove fractional partial quotients. However, unlike negative integers, it is possible that the new element found may in fact be even “worse” and so propagate through the expansion.

2.1. Multiplying a continued fraction by a rational

Raney [59] provides a method by which one can easily determine any rational multiple of a continued fraction (in fact [59] does a little more, it provides a procedure for determining any linear fractional transformation of an element, but we will only be interested in using it for multiplying and dividing). We first need some notation, we put:

$$R = \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix}, \quad L = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}, \quad J = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix},$$

so that

$$\begin{pmatrix} a & 1 \\ 1 & 0 \end{pmatrix} J = R^a, \quad J \begin{pmatrix} a & 1 \\ 1 & 0 \end{pmatrix} = L^a, \quad J^2 = I.$$

Then matrix products can be represented as R – L sequences,

$$\begin{pmatrix} a & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} b & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} c & 1 \\ 1 & 0 \end{pmatrix} \cdots = R^a L^b R^c \cdots$$

In order to perform the transformation one needs to know all the 2×2 doubly balanced matrices of the determinant in question. (For this purpose, a 2×2 matrix of positive determinant is called doubly balanced if the entries on the main diagonal are greater than the entries off it.) From there one produces a transition table (see [59] for the details) which prescribes the exact transformation. For example, suppose that we wish to determine 3 times $\alpha = [7, 1, 4, 23, 9, 1, 1, 17]$. Then the set of doubly balanced matrices of determinant 3 is

$$A = \begin{pmatrix} 3 & 0 \\ 0 & 1 \end{pmatrix}, \quad B = \begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix}, \quad A' = \begin{pmatrix} 1 & 0 \\ 0 & 3 \end{pmatrix},$$

where the third matrix is called A' rather than C since it acts like the opposite of A . The transition table for these matrices is

	A	B	A'
A	$L^3 : L$	$R : R^3$	$L^2 R : RL^2$
B	$L : LR$		$R : RL$
A'	$R^2 L : LR^2$	$RL : L$	$L : L^3$
			$R^3 : R$

Then the R – L sequence for α is $R^7 L^1 R^4 L^{23} R^9 L^1 R^1 L^{17} J$ and its transitions proceed as:

Input seq.	R^7	LR	R	$R^2 L$	L^{21}	LR	R	R^6	RL	R	L^{17}	J
State	A	A	B	A'	A	A	B	A'	A'	B	A'	A'
Output seq.	R^{21}	R	RL	LR^2	L^7	R	RL	R^2	L	RL	L^{51}	J

Compacting the output sequence, we find

$$\begin{aligned} 3\alpha &= R^{23}L^2R^2L^7R^2L^1R^2L^1R^1L^{52} \\ &= [23, 2, 2, 7, 2, 1, 2, 1, 1, 52]. \end{aligned}$$

On the other hand, we may also use Lemma 2 to carry out this multiplication. Namely,

$$\begin{aligned} 3\alpha &= [21, \frac{1}{3}, 12, \frac{23}{3}, 27, \frac{1}{3}, 3, \frac{17}{3}] \\ &= [21, 0, 2, 1, \frac{1}{9}[12, \frac{23}{3}, 27, \frac{1}{3}, 3, \frac{17}{3}] - \frac{2}{3}] \\ &= [23, 1, \frac{2}{3}, 69, 3, 3, \frac{1}{3}, 51] \\ &= [23, 1, 0, 1, 2, \frac{1}{9}[69, 3, 3, \frac{1}{3}, 51] - \frac{1}{3}] \\ &\quad \vdots \\ &= [23, 2, 2, 7, 2, 1, 2, 2, 1, -1, 1, 53] \\ &= [23, 2, 2, 7, 2, 1, 2, 1, 1, 52] \end{aligned}$$

Needless to say this is not a particularly efficient method. Since the set of linear fractional transformations is generated by inversion and translation, both of which are easy to do, Lemma 2 can be used to determine any linear fractional transformation of α .

2.2. Continued fractions of quadratic irrationals

Since continued fractions provide good approximations (in fact if α is irrational then its convergents provide the locally best rational approximations), it makes sense to consider the continued fraction expansion of quadratic irrationals. As an example, here is the expansion of $\sqrt{67}$,

$$\begin{aligned} \sqrt{67} &= 8 - (-\sqrt{67} + 8) \\ \frac{\sqrt{67} + 8}{3} &= 5 - \frac{(-\sqrt{67} + 7)}{3} \\ \frac{\sqrt{67} + 7}{6} &= 2 - \frac{(-\sqrt{67} + 5)}{6} \\ \frac{\sqrt{67} + 5}{7} &= 1 - \frac{(-\sqrt{67} + 2)}{7} \\ \frac{\sqrt{67} + 2}{9} &= 1 - \frac{(-\sqrt{67} + 7)}{9} \\ \frac{\sqrt{67} + 7}{2} &= 7 - \frac{(-\sqrt{67} + 7)}{2}. \end{aligned}$$

There is no need to continue further since conjugation of the above tableau yields the remaining portion.

We have determined $\sqrt{67} = [8, 5, 2, 1, 1, 7, 1, 1, 2, 5, 16, 5, 2, 1, 1, 7, \dots]$ which is abbreviated as $\sqrt{67} = [8, \overline{5, 2, 1, 1, 7, 1, 1, 2, 5, 16}]$. The length of the overline is called the period length. For brevity this is expressed as $lp(\sqrt{67}) = 10$. When speaking of period length, we mean in the *regular* continued fraction expansion, thus period length is well-defined. The smallest possible period of α is called the *primitive* period of α .

Given a rational expansion $[a_0, \dots, a_h] = a/b$, we write $l(a/b) = h + 1$ which is dependent on the parity chosen for the expansion. If the parity has not been

determined then we call the *depth* of a/b , usually represented by $\delta(a/b)$, to be the number of partial quotients in the *even* length expansion of a/b .

The above tableau suggests an exact procedure for determining the continued fraction expansion of $\sqrt{D}$ for any $D \in \mathbb{N}$. A typical line in the expansion of $\sqrt{D}$ appears as

$$\frac{\sqrt{D} + P_h}{Q_h} = a_h - \frac{(-\sqrt{D} + P_{h+1})}{Q_h},$$

so that the complete quotients of $\sqrt{D}$ are $(\sqrt{D} + P_h)/Q_h$. Writing the expansion in the above manner also makes it easy to determine successive P 's and Q 's,

$$P_{h+1} = a_h Q_h - P_h, \quad (2.4a)$$

$$Q_{h+1} = (D - P_{h+1}^2)/Q_h. \quad (2.4b)$$

The end of a period corresponds to the occurrence of $Q_h = 1$, therefore we can say that $lp(\sqrt{D}) = \min\{h : h > 0 \text{ and } Q_h = 1\}$.

Deeper examination of the above tableau would demonstrate that

$$x_h^2 - Dy_h^2 = (-1)^{h+1} Q_{h+1} \quad (2.5)$$

(which like most observations is best proved by the matrix relation for continued fractions). At the end of a period we obtain a solution of the Pell equation

$$x^2 - Dy^2 = \pm 1$$

with the minus sign applying if and only if the period length is odd.

Equally, the equation $x^3 - 5$ allows one to readily describe the continued fraction expansion of $\sqrt[3]{5}$ to arbitrary precision. But the property which makes quadratic irrationals special is that of periodicity. The following, [81], is a well known result of Lagrange,

Theorem 3. *A continued fraction expansion is periodic if and only if it is the expansion of a quadratic irrational.*

There is no need to merely consider $\sqrt{D}$. Looking at the more general quadratic irrational $\gamma = (a + b\sqrt{D})/c$ is just as applicable. When we examine such an element we insist that $c \mid a^2 - b^2 D$; there is no loss in generality in doing so. As an initial warning, it may be the case that there is a preperiod. Consider

$$\gamma = \frac{7 - \sqrt{71}}{2} = [-1, 3, 2, \overline{16, 2, 2, 1, 7, 1, 2, 2}]$$

and

$$\zeta = 8 + \sqrt{71} = [\overline{16, 2, 2, 1, 7, 1, 2, 2}].$$

Notice that γ and ζ both have the same “tail”. This means that the two elements are equivalent. More precisely, two elements $\gamma, \zeta \in \mathbb{Q}(\sqrt{D})$ are called *equivalent* if there exists some transformation $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$ of determinant ± 1 such that

$$\gamma = \frac{a\zeta + b}{c\zeta + d}.$$

By taking the decomposition of the matrix $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$ this is the same as saying that γ and ζ have the same “tail”, that is

$$\gamma = [a_0, \dots, a_r, \overline{b_1, b_2, \dots, b_h}] \quad \text{and} \quad \zeta = [c_0, \dots, c_s, \overline{b_1, b_2, \dots, b_h}].$$

Thus any element appearing in the continued fraction expansion of γ will be equivalent to γ . In the above example, γ and ζ are related via the transformation $\begin{pmatrix} -5 & -2 \\ 7 & 3 \end{pmatrix}$ which follows from

$$\begin{pmatrix} -5 & -2 \\ 7 & 3 \end{pmatrix} = \begin{pmatrix} -1 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 3 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 2 & 1 \\ 1 & 0 \end{pmatrix}.$$

If the expansion of a quadratic irrational α has no preperiod, that is $\alpha = [\overline{a_0, \dots, a_h}]$, then α is called *purely periodic*. It is a result of Galois that a quadratic irrational α has a purely periodic continued fraction expansion if and only if it is *reduced*, where reduced means

$$\alpha > 1 \quad \text{and} \quad -1 < \bar{\alpha} < 0.$$

As an aside, note that $\sqrt{71}$ is not reduced because the continued fraction expansion is not purely periodic (and $-\sqrt{71} < -1$) but $8 + \sqrt{71}$ is reduced. If a complete quotient, $(\sqrt{D} + P_h)/Q_h$ is reduced then

$$0 < P_h < \sqrt{D} \quad \text{and} \quad 0 < Q_h < 2\sqrt{D}.$$

Recall that an algebraic integer is a root of a monic equation, thus a quadratic integer γ is a root of

$$x^2 - tx + n = 0 \quad \text{for some } t, n \in \mathbb{Z}.$$

The number t is called the *trace* of γ , and n the *norm* of γ ; since $\gamma + \bar{\gamma} = t$ and $\gamma\bar{\gamma} = n$. The bar, $\bar{\gamma}$, is used to denote the nontrivial automorphism of $\mathbb{Q}(\gamma)$.

Thus $(11 + 3\sqrt{31})/5$ is an algebraic number since it satisfies $25x^2 - 110x - 158$, but $5 + 3\sqrt{51}$ is a quadratic integer since it satisfies $x^2 - 10x - 434$. As a further example the expansion of $\gamma = 5 + 3\sqrt{51}$ is

$$\begin{aligned} \gamma &= 26 - (\bar{\gamma} + 16) \\ (\gamma + 16)/18 &= 2 - (\bar{\gamma} + 10)/18 \\ (\gamma + 10)/13 &= 2 - (\bar{\gamma} + 6)/13 \\ (\gamma + 6)/26 &= 1 - (\bar{\gamma} + 10)/26 \\ (\gamma + 10)/9 &= 4 - (\bar{\gamma} + 16)/9 \\ (\gamma + 16)/2 &= 21 - (\bar{\gamma} + 16)/2 \\ &\vdots \end{aligned}$$

And so $5 + 3\sqrt{51} = [26, \overline{2, 2, 1, 4, 21, 4, 1, 2, 2, 42}]$. The observant reader may have noticed that both of the detailed expansions have had $Q_h = 2$ occur at the halfway point of the period. When expanding $\sqrt{D}$ it is the case that $Q_h = 2$ implies that the period length is $2h$. However, for other quadratic irrationals this need not be the case, for example $(1 + \sqrt{153})/2$ has $Q_3 = 2$ yet the period length is 8. Furthermore, for the expansion of $\sqrt{D}$, a $Q_h = 4$ occurs at roughly one third of the primitive period of $\sqrt{D}$. This concept is generalized in [54] where the existence of two complete quotients of $\sqrt{D_n}$ satisfying $Q_h = 3W$ and $Q_{h'} = W$ occur halfway to $Q_{h''} = 3$, as well as revealing a symmetry within the partial quotients.

In our simpler case, the symmetry in the expansion of γ can be summarized as

$$\begin{aligned} P_h &= P_{h+1} \quad \text{if and only if} \quad lp(\gamma) = 2h, \\ Q_h &= Q_{h+1} \quad \text{if and only if} \quad lp(\gamma) = 2h + 1. \end{aligned}$$

Writing the remainders as $-(\bar{\gamma} + P_h)/Q_h$ again makes it easy to determine the reciprocal $(\gamma + P_{h+1})/Q_{h+1}$ namely

$$\begin{aligned} P_{h+1} &= a_h Q_h - P_h - (\gamma + \bar{\gamma}), \\ Q_{h+1} &= -\frac{\gamma\bar{\gamma} + P_h^2 + P_h(\gamma + \bar{\gamma})}{Q_h}. \end{aligned}$$

If the minimal polynomial of γ is $x^2 - tx + n$ then these can be rewritten as

$$\begin{aligned} P_{h+1} &= a_h Q_h - P_h - t, \\ Q_{h+1} &= -\frac{n + P_h^2 + tP_h}{Q_h}. \end{aligned}$$

which generalise (2.4). The bounds for P_h and Q_h of a reduced complete quotient are now given by

$$0 < 2P_h + t < \gamma - \bar{\gamma} \quad \text{and} \quad 0 < Q_h < \gamma - \bar{\gamma}. \quad (2.6)$$

The generalisation of (2.5) is

$$x_h^2 - (\gamma + \bar{\gamma})x_h y_h + \gamma \bar{\gamma} y_h^2 = (-1)^{h+1} Q_{h+1}.$$

Of some importance is the following result,

Proposition 4. *Given a solution (x, y) of Pell's equation*

$$(x - \gamma y)(x - \bar{\gamma} y) = \pm 1,$$

each decomposition

$$\begin{pmatrix} x & -\gamma \bar{\gamma} y \\ y & x - (\gamma + \bar{\gamma})y \end{pmatrix} = \begin{pmatrix} a_0 & 1 \\ 1 & 0 \end{pmatrix} \cdots \begin{pmatrix} a_s & 1 \\ 1 & 0 \end{pmatrix}$$

with integers a_i yields a purely-periodic continued fraction expansion

$$\gamma = [\overline{a_0, \dots, a_s}]$$

Proof. Since the cited matrix is a unimodular matrix with integer entries it may be decomposed as a product of elementary row transformations with integers a_i . Given the decomposition, suppose that

$$\alpha = [\overline{a_0, a_1, \dots, a_s}].$$

Then by the matrix correspondence and the meaning of periodicity, we have

$$\begin{aligned} \alpha &= [\overline{a_0, a_1, \dots, a_s}] = [a_0, a_1, \dots, a_s, \alpha] \\ &\longleftrightarrow \begin{pmatrix} x & -\gamma \bar{\gamma} y \\ y & x - (\gamma + \bar{\gamma})y \end{pmatrix} \begin{pmatrix} \alpha & 1 \\ 1 & 0 \end{pmatrix} = \begin{pmatrix} x\alpha - \gamma \bar{\gamma} y & x \\ y\alpha + x - (\gamma + \bar{\gamma})y & y \end{pmatrix} \\ &\longleftrightarrow \alpha = (x\alpha - \gamma \bar{\gamma} y) / (y\alpha + x - (\gamma + \bar{\gamma})y) \end{aligned}$$

Hence,

$$y(\alpha^2 - (\gamma + \bar{\gamma})\alpha + \gamma \bar{\gamma}) = 0.$$

■

Note that the a_i need not be positive, so this doesn't contradict the lack of pure-periodicity in the regular continued fraction expansion of $\sqrt{D}$ since $\sqrt{D} = [a_0, a_1, \dots, a_{h-1}, a_0, 0]$.

3. UNITS AND ORDERS

For a quadratic field $K = \mathbb{Q}(\sqrt{D})$ the ring of algebraic integers, denoted by $\mathcal{O}_K$, can be described as $[1, \omega'] = \mathbb{Z} + \omega' \mathbb{Z}$ where d_K is the squarefree kernel of D and

$$\omega' = \begin{cases} \sqrt{d_K} & \text{if } d_K \not\equiv 1 \pmod{4} \\ (1 + \sqrt{d_K})/2 & \text{if } d_K \equiv 1 \pmod{4} \end{cases}$$

The *discriminant*, D_K , of $\mathcal{O}_K$ is equal to d_K if d_K is congruent to 1 modulo 4 and $4d_K$ otherwise.

By defining an *order* $\mathcal{O}$ of K to be any free $\mathbb{Z}$ -module of rank 2 which is a subring of K and contains 1, one observes, [13], that the maximal order of K is none other than $\mathcal{O}_K$ and that any other order may be represented as

$$\mathcal{O} = [1, f\omega'] = \mathbb{Z} + f\omega'\mathbb{Z}.$$

The number f is called the *conductor* of $\mathcal{O}$. The *discriminant* of an order $\mathcal{O} \subset \mathcal{O}_K$ is equal to $D = f^2 D_K$. Thus, the discriminant of an order is always congruent to 0 or 1 modulo 4. The discriminant of the maximal order is called *fundamental*.

For any discriminant, $D \equiv t \pmod{4}$, the element $\omega = (t + \sqrt{D})/2$ is an algebraic integer since $t \equiv 0, 1 \pmod{4}$. In the sequel ω shall be strictly reserved for these purposes. If we know that D is fundamental then we usually write ω' .

The *norm* of $\alpha \in \mathbb{Q}(\sqrt{D})$ is defined as $N(\alpha) = \alpha\bar{\alpha}$ where the overline represents the nontrivial automorphism of the field. Moreover, the norm is clearly multiplicative. A *unit*, ε , of $\mathcal{O}$ is an algebraic integer which divides 1, hence necessarily, $N(\varepsilon) = \pm 1$. The *fundamental unit* is the smallest nontrivial positive unit. Dirichlet's unit theorem states that for any real quadratic number field the set of units of $\mathcal{O}$ is an abelian group under multiplication generated by -1 and the fundamental unit. That is, any other nontrivial unit η , satisfies $\eta = \pm \varepsilon^r$ for some $r \in \mathbb{Z}$. Furthermore the norm of a unit $\varepsilon = x + \omega y$ must satisfy

$$N(\varepsilon) = \begin{cases} x^2 - \frac{D}{4}y^2 & \text{for } D \equiv 0 \pmod{4}, \\ x^2 - xy + \frac{1-D}{4}y^2 & \text{for } D \equiv 1 \pmod{4}. \end{cases}$$

But such a solution corresponds directly to the convergents of the primitive period of ω . Thus the continued fraction expansion provides an excellent method for determining the fundamental unit. There are of course other methods, such as Gauss' cyclotomic approach, see [16], but we won't consider any of them here.

If we wish to determine the fundamental unit (as we do) then keeping track of the convergents is rather excessive. Instead it will suffice to note

Proposition 5. *If $\mathcal{O}$ is an order of $\mathbb{Q}(\sqrt{D})$ and*

$$\alpha_i = \frac{\alpha + P_i}{Q_i} \quad i = 0, \dots, h$$

represents a system of reduced elements in any cycle of quadratic irrationals in $\mathcal{O}$, which has period length $h + 1$, then

$$\varepsilon = \prod_{i=1}^{h+1} \alpha_i$$

is the fundamental unit of $\mathcal{O}$.

Proof. From the matrix product representation, if

$$\alpha_0 = [a_0, a_1, \dots, a_i, \alpha_{i+1}]$$

then

$$\alpha_{i+1} = -\frac{y_{i-1}\alpha_0 - x_{i-1}}{y_i\alpha_0 - x_i} \quad \text{for } i \geq 0.$$

From this we find

$$\begin{aligned} \alpha_1 \alpha_2 \dots \alpha_{h+1} &= (-1)^{h+1} (x_h - y_h \alpha_0)^{-1} \\ &= \frac{(-1)^{h+1} (x_h - y_h \bar{\alpha}_0)}{(x_h - y_h \alpha_0)(x_h - y_h \bar{\alpha}_0)}. \end{aligned}$$

Since α_0 is reduced, we have $\alpha_{h+1} = \alpha_0$, so $Q_{h+1} = 1$ and $(x_h - y_h \alpha_0)(x_h - y_h \bar{\alpha}_0) = (-1)^{h+1}$. Thus $\alpha_1 \dots \alpha_{h+1}$ is a unit of $\mathcal{O}$. Furthermore, if $h + 1$ is a primitive period, this unit is also fundamental in $\mathcal{O}$. ■

As a corollary we obtain the well-known result that the product of all reduced elements in $\mathcal{O}$ is equal to ε^h , where ε is the fundamental unit of $\mathcal{O}$ and h is the number of inequivalent cycles of $\mathcal{O}$.

Since the growth of the convergents is quite furious the size of the fundamental unit will in general be exceedingly large. In order to circumvent this the *regulator* of the order $\mathcal{O}$, denoted by $R(\mathcal{O})$, is defined as the logarithm of the fundamental unit. If $\mathcal{O}$ is the maximal order of $\mathbb{Q}(\sqrt{D})$ then we also denote the regulator by $R(D)$.

Once the continuants of the primitive period have been computed, all future continuants can be found in terms of these, that is

Proposition 6. *If α has period length $h + 1$ then*

$$x_{h+k} - y_{h+k}\bar{\alpha} = (x_h - y_h\bar{\alpha})(x_{k-1} - y_{k-1}\bar{\alpha}).$$

Proof. From above, we have

$$\alpha_1 \dots \alpha_{h+1+k} = \frac{(-1)^{h+1+k}(x_{h+k} - y_{h+k}\bar{\alpha})}{N(x_{h+k} - y_{h+k}\alpha)}.$$

Using periodicity, we may also write,

$$\begin{aligned} \alpha_1 \dots \alpha_{h+1+k} &= (\alpha_1 \dots \alpha_{h+1})(\alpha_1 \dots \alpha_k) \\ &= \frac{(-1)^{h+1}(x_h - y_h\bar{\alpha})}{N(x_h - y_h\alpha)} \cdot \frac{(-1)^k(x_{k-1} - y_{k-1}\bar{\alpha})}{N(x_{k-1} - y_{k-1}\alpha)} \end{aligned}$$

and since $N(x_{k-1} - y_{k-1}\alpha) = (-1)^{h+1}N(x_{k+h} - y_{k+h}\alpha)$ we obtain the desired result. $\blacksquare$

3.1. Quadratic forms and continued fractions

To any real quadratic form $ax^2 + bxy + cy^2$ of discriminant D , with $D \equiv t \pmod{4}$, we can associate the quadratic irrational $\gamma = \alpha + (b-t)/2/|a|$ where α is a root of $x^2 - tx + (t-D)/4$. Also note that $D \equiv t \equiv b^2 \pmod{4}$ so $(b-t)/2$ is an integer. However, to each quadratic irrational $(\alpha + P)/Q$ we can associate two quadratic forms: $(Q, 2P+t, -Q')$ and $(-Q, 2P+t, Q')$. If these two forms are equivalent then the number of forms in the cycle of $(Q, 2P+t, -Q')$ will be twice that of the period length of $(\alpha + P)/Q$. Note that this condition is equivalent to being able to solve $(x - \alpha y)(x - \bar{\alpha} y) = -1$. In this case the number of distinct cycles of reduced forms will equal the number of distinct cycles of quadratic irrationals. On the other hand, if $(x - \alpha y)(x - \bar{\alpha} y) = -1$ is unsolvable then the two forms (a, b, c) and $(-a, b, -c)$ are not equivalent yet the two corresponding cycles are absorbed by the one cycle of the quadratic irrational $(\alpha + (b-t)/2)/|a|$. In other words the number of distinct reduced cycles of forms will be twice the number of distinct cycles of quadratic irrationals. In conclusion, using continued fractions we can determine the form class number.

As an example, consider $\alpha = \sqrt{229}$. On the left is the expansion of α and on the right are the corresponding principal forms:

α	$=$	15	$-$	$(\bar{\alpha} + 15)$	f_0	$=$	$(1, 0, -229)$
$(\alpha + 15)/4$	$=$	7	$-$	$(\bar{\alpha} + 13)/4$	f_1	$=$	$(-4, 30, 1)$
$(\alpha + 13)/15$	$=$	1	$-$	$(\bar{\alpha} + 2)/15$	f_2	$=$	$(15, 26, -4)$
$(\alpha + 2)/15$	$=$	1	$-$	$(\bar{\alpha} + 13)/15$	f_3	$=$	$(-15, 4, 15)$
$(\alpha + 13)/4$	$=$	7	$-$	$(\bar{\alpha} + 15)/4$	f_4	$=$	$(4, 26, -15)$
$(\alpha + 15)/1$	$=$	30	$-$	$(\bar{\alpha} + 15)/1$	f_5	$=$	$(-1, 30, 4)$
$(\alpha + 15)/4$	$=$	7	$-$	$(\bar{\alpha} + 13)/4$	f_6	$=$	$(4, 30, -1)$

Thus, the length of the cycle of reduced principal forms is twice that of the period length of the expansion of $\sqrt{229}$. Note that the discriminant of the forms is $4 \cdot 229$. Also observe that as expected, $f_1 = (-4, 30, 1)$ is equivalent to $f_6 = (4, 30, -1)$. Moreover, since this can only occur when the period length of the continued fraction is odd we must have a form $(c, b, -c)$ occurring at halfway, which we do, that is f_3 . In conclusion, the form class number of 916 will be equal to the number of inequivalent reduced cycles of quadratic irrationals in $\mathbb{Q}(\sqrt{229})$. Also note that by Proposition 5, a unit is given by

$$\begin{aligned} \varepsilon &= \frac{\alpha + 15}{4} \cdot \frac{\alpha + 13}{15} \cdot \frac{\alpha + 2}{15} \cdot \frac{\alpha + 13}{4} \cdot \frac{\alpha + 15}{1} \\ &= 1710 + 113\sqrt{229} \end{aligned}$$

However, this is *not* the fundamental unit of the ring of integers in $\mathbb{Q}(\sqrt{229})$. In this case the maximal order is generated by 1 and $(1 + \sqrt{229})/2$, since $229 \equiv 1 \pmod{4}$. Thus a proper examination gives,

$$\omega = (1 + \sqrt{229})/2 = [8, \overline{15}]$$

so that

$$\varepsilon_1 = 7 + \omega$$

is a fundamental unit of the ring of integers of $\mathbb{Q}(\sqrt{229})$ with norm -1 . Furthermore, we have $\varepsilon = (\varepsilon_1)^3$, a feature which is always true in these cases, see for example [12]. If we wished to determine the form class number of 229 then we would need to examine the inequivalent reduced cycles of $\mathbb{Q}(\omega)$.

One might also notice that the expansion of ω is exceptionally short, and this is what leads to the class number being somewhat large (229 is the first prime congruent to 1 modulo 4 which has a class number greater than 1). In Chapter II we will explain more about such examples. Another cycle of $\mathbb{Q}(\alpha)$ is:

$$\begin{array}{rclcl} (\alpha + 14)/3 & = & 9 & - & (\overline{\alpha} + 13)/3 \\ (\alpha + 13)/20 & = & 1 & - & (\overline{\alpha} + 7)/20 \\ (\alpha + 7)/9 & = & 2 & - & (\overline{\alpha} + 11)/9 \\ (\alpha + 11)/12 & = & 2 & - & (\overline{\alpha} + 13)/12 \\ (\alpha + 13)/5 & = & 5 & - & (\overline{\alpha} + 12)/5 \\ (\alpha + 12)/17 & = & 1 & - & (\overline{\alpha} + 5)/17 \\ (\alpha + 5)/12 & = & 1 & - & (\overline{\alpha} + 7)/12 \\ (\alpha + 7)/15 & = & 1 & - & (\overline{\alpha} + 8)/15 \\ (\alpha + 8)/11 & = & 2 & - & (\overline{\alpha} + 14)/11 \\ (\alpha + 14)/3 & = & 9 & - & (\overline{\alpha} + 13)/3 \end{array}$$

Note that this expansion incorporates two inequivalent classes. Namely the one containing $(\alpha + 14)/3$ and the one which by conjugation, contains $(\alpha + 13)/3$. That the two are inequivalent follows from the non-existence of $(\alpha + 13)/3$ in the list of elements in this expansion. By using the bounds on P_h and Q_h given in (2.6), we see that there are no further reduced quadratic irrationals in $\mathbb{Q}(\sqrt{229})$. Thus the form class number of 916 is 3. Similarly, one finds that the form class number of 229 is 3.

Clearly, the above procedure is completely impractical for large discriminants, but fortunately it happens that there are analytic means which allow for easier computation. One can compute the class number from Dirichlet's class number formula, [12], but we shall have no need to use it here.

Proposition 5 would suggest that the number of forms in each cycle should not vary too much and in fact, utilising the bounds on the distance between forms one can find, [33], that

$$\frac{l_1}{l_2} < \frac{\log D}{\log 2}$$

where l_1 and l_2 represent the number of reduced forms in any two cycles of discriminant D .

Symmetry (which as exemplified as above is not a general characteristic) follows simply from whether or not a form in the particular class is equivalent to its conjugate or not. Forms which are equivalent to their conjugates are called *ambiguous*. These forms play a vital role in factoring algorithms based on quadratic orders since they correspond to squarefree divisors of D . For more on ambiguity and symmetry see [50].

Given two orders, $\mathbb{Z}[f\omega]$ and $\mathbb{Z}[g\omega]$, where $f|g$, a natural question is to relate the fundamental unit of $\mathbb{Z}[f\omega]$, ε_f , with that of $\mathbb{Z}[g\omega]$, ε_g . First, if p is a rational prime and $F(X) \in \mathbb{Z}[X]$ then $F(X)^p \equiv F(X^p) \pmod{p}$. Hence, if F is the minimal polynomial of ε_f then $\varepsilon_f^p \equiv \varepsilon_f$ or $\bar{\varepsilon}_f \pmod{p}$. More generally, [85], for $\omega = \sqrt{D}$ and $g/f = p^k$, by taking

$$\psi_D(p^k) = \begin{cases} 2^k & \text{if } p = 2 \\ p^k & \text{if } p \text{ is odd and } p|D \\ p^{k-1} \left(p - \left(\frac{D}{p} \right) \right) / 2 & \text{otherwise} \end{cases}$$

then $\varepsilon_g = \varepsilon_f^{\psi_D(p^k)}$. This can be extended to all numbers via

$$\psi_D(p_1^{e_1} \dots p_k^{e_k}) = \text{LCM}[\psi_D(p_1^{e_1}), \dots, \psi_D(p_k^{e_k})]$$

and then $\varepsilon_f^{\psi_D}$ is a unit of $\mathbb{Z}[g\omega]$, not necessarily fundamental. On the other hand, this is equivalent to comparing the continued fraction expansions of $g\omega$ with $f\omega$, i.e. a multiplication of g/f . While it is easy to compare the units, there is no general procedure for comparing the two expansions and in most cases, one is left to physically compute the two.

For most practical purposes quadratic forms are more than sufficient, however they are somewhat limited algebraically. It turns out that a more useful concept is that of ideals.

4. IDEALS, INFRASTRUCTURE AND COMPOSITION

Gauss was certainly aware of identities like

$$2 \cdot 3 = (1 + \sqrt{-5})(1 - \sqrt{-5})$$

which exemplify the failure of unique prime factorization in $\mathbb{Z}[\sqrt{-5}]$. But it was not until the latter part of the 19th Century that the general mathematical community came to realize that there was a serious issue here. The first resolution was provided by Kummer, but the one which caught on and became prevalent was Dedekind's concept of ideals. The following results are also described in [51], [26], [54]. Throughout, K represents a real quadratic field, $\mathcal{O}_K$ its maximal order and $\mathcal{O}$ some order of K .

An *integral ideal* $\mathfrak{a}$ of $\mathcal{O}$ is any subset such that whenever $\alpha, \beta \in \mathfrak{a}$ then $\alpha \pm \beta \in \mathfrak{a}$ and for every $\gamma \in \mathcal{O}$ we have $\gamma\alpha \in \mathfrak{a}$. A *fractional ideal* of $\mathcal{O}$ is a subset $\mathfrak{a}$ of K such that $d\mathfrak{a}$ is an integral ideal of $\mathcal{O}$ for some $d \in K^*$. The norm of an integral ideal $\mathfrak{a}$ is defined as $|\mathcal{O}/\mathfrak{a}|$; it follows that the norm is finite and multiplicative.

If $\alpha_1, \dots, \alpha_j \in K^*$ then $\mathfrak{a} = \left\{ \sum_{i=1}^j \gamma_i \alpha_i : \gamma_i \in \mathcal{O} \right\}$ is an ideal of $\mathcal{O}$ which we denote by $(\alpha_1, \dots, \alpha_j)$ and say that it is generated by $\alpha_1, \dots, \alpha_j$. If an ideal is

generated by a single element it is called *principal*, the set of all principal ideals of $\mathcal{O}$ is denoted by $\mathcal{P}$. If $\mathfrak{a} = (\alpha_1, \dots, \alpha_n)$ and $\mathfrak{b} = (\beta_1, \dots, \beta_m)$ then the product $\mathfrak{a}\mathfrak{b}$ is given by $\mathfrak{a}\mathfrak{b} = (\alpha_1\beta_1, \alpha_1\beta_2, \dots, \alpha_n\beta_{m-1}, \alpha_n\beta_m)$ with nm generators. We say $\mathfrak{a}|\mathfrak{b}$ if $\mathfrak{a}\mathfrak{c} = \mathfrak{b}$ for some $\mathfrak{c} \in \mathcal{O}$; this is equivalent to $\mathfrak{a} \supset \mathfrak{b}$. Then a prime ideal $\mathfrak{p}$ of $\mathcal{O}$ is one which is not divisible by any ideals other than (1) and $\mathfrak{p}$. Equivalently, $\mathcal{O}/\mathfrak{p}$ is an integral domain. An ideal $\mathfrak{a}$ is called *primitive* if it has no rational integer divisors.

The key feature of ideals in a maximal order is that they possess unique factorisation. Consequently, they also possess analogues of important results from integers. For example, the Chinese Remainder Theorem,

Theorem 7. *Let $\mathfrak{a} = \mathfrak{p}_1^{e_1} \dots \mathfrak{p}_j^{e_j}$ be an ideal of $\mathcal{O}_K$. Then the natural map, $\phi : \mathcal{O}_K \rightarrow \prod (\mathfrak{p}_i^{e_i})$ is onto and induces an isomorphism,*

$$\mathcal{O}_K/\mathfrak{a} \approx \prod (\mathcal{O}_K/\mathfrak{p}_i^{e_i})$$

A proof can be found in any introduction to algebraic number theory, for example [32], [68]. As a corollary,

Corollary 8. *Any integral ideal $\mathfrak{a}$ of $\mathcal{O}_K$ can be represented as $\mathfrak{a} = (N(\mathfrak{a}), \alpha)$ for some $\alpha \in \mathcal{O}_K$.*

Proof. Let $\mathfrak{q}_1, \dots, \mathfrak{q}_j$ be the prime factors of $(N(\mathfrak{a}))$. For $i = 1, \dots, j$ let e_i satisfy $\mathfrak{q}_i^{e_i} \parallel \mathfrak{a}$ and pick some $\alpha_i \in \mathfrak{q}_i^{e_i}$ which is not in $\mathfrak{q}_i^{e_i+1}$. By the Chinese Remainder Theorem there exists an $\alpha \in \mathcal{O}_K$ such that

$$\alpha \equiv \alpha_i \pmod{\mathfrak{q}_i^{e_i+1}} \quad i = 1, \dots, j \quad \text{and} \quad \alpha \equiv 0 \pmod{\mathfrak{a} / \prod \mathfrak{q}_i^{e_i}}$$

Hence, $(N(\mathfrak{a}))/\mathfrak{a}$ and (α) are coprime and thus $\mathfrak{a} = (N(\mathfrak{a}), \alpha)$. ■

However, if the conductor of an order is greater than 1 then $\mathcal{O}$ is *not* integrally closed in K and thus, unique factorization of ideals may fail to hold. This can be remedied by considering ideals prime to the conductor¹. Any integral ideal $\mathfrak{a}$ of $\mathcal{O}$ may be written as

$$\mathfrak{a} = c\langle Q, \omega + P \rangle = cQ\mathbb{Z} + c(\omega + P)\mathbb{Z}, \quad c \in \mathbb{Z}.$$

Such a representation is called a *normal $\mathbb{Z}$ -basis* of $\mathfrak{a}$. Also note that $\mathfrak{a}$ is primitive if and only if $c = 1$. The norm of $\mathfrak{a}$ is equal to c^2Q . The conjugate of $\mathfrak{a}$, which we denote by $\overline{\mathfrak{a}}$, is given by $\langle cQ, c(\overline{\omega} + P) \rangle$; this gives $\mathfrak{a}\overline{\mathfrak{a}} = (N(\mathfrak{a}))$. The procedure of taking the normal $\mathbb{Z}$ -bases for two ideals, $\mathfrak{a}$, $\mathfrak{b}$, and giving the normal $\mathbb{Z}$ -basis of their product $\mathfrak{a}\mathfrak{b}$ is known as *composition*, which is detailed on page 20.

An ideal $\mathfrak{a}$ of $\mathcal{O}$ is called *invertible* if it divides the unit ideal (the ideal generated by 1), that is $\mathfrak{a}\mathfrak{a}' = \mathcal{O}$ for some ideal $\mathfrak{a}'$ of $\mathcal{O}$. The set of all invertible ideals of $\mathcal{O}$ is denoted by $\mathcal{I}$. Two ideals $\mathfrak{a}$, $\mathfrak{b}$ are called *equivalent* if $\mathfrak{a} = (\alpha)\mathfrak{b}$ for some $\alpha \in K^*$. If $N(\alpha) > 0$ then they are called *properly equivalent*. This is an equivalence relation and the group $\mathcal{I}/\mathcal{P}$ is called the *ideal class group* of $\mathcal{O}$. Typically, $h(D)$ is used to represent the *ideal class number* which is the order of the group. Using *narrow* equivalence, that is $N(\alpha) > 0$, one finds the *narrow* class group. Note that there can only be a distinction between the two when the norm of the fundamental unit is 1. The connection to the form class group is given by

Theorem 9. *There is a 1-to-1 correspondence between the representatives of the form class group and representatives of the narrow ideal class group.*

Observing that if $N(\varepsilon) = 1$ then an ideal class in the ordinary sense comprises two ideal classes in the narrow sense and this shows that in this case the form class number is twice the ideal class number. See either [13] or [7] for more details on this correspondence.

¹An ideal $\mathfrak{a}$ is called prime to the conductor if $(N(\mathfrak{a}), f) = 1$

An integral ideal $\mathfrak{a}$ is *reduced* if there does not exist any non-zero $\alpha \in \mathfrak{a}$ which satisfies

$$|\alpha| < N(\mathfrak{a}) \quad \text{and} \quad |\bar{\alpha}| < N(\mathfrak{a}).$$

The class number and regulator are combined in Siegel's formula [32],

$$\log(h(D)R(D)) \sim \log(\sqrt{D})$$

for any infinite sequence of fundamental discriminants. For real quadratic fields, this says that $h(D) = 1$ infinitely often is very much like having regulators near to $\sqrt{D}$ infinitely often. Thus, the class number 1 problem for real quadratic fields is akin to finding large regulators which is the same as finding long period lengths.

Unique factorisation of ideals leads us to want to be able to characterize the prime ideals. The prime ideals $\mathfrak{p}$ of $\mathcal{O}$ satisfy $\mathfrak{p} \cap \mathbb{Z} = p\mathbb{Z}$ for some prime rational integer p , and we say that $\mathfrak{p}$ *lies over* p . The prime ideals can be described by,

$$\left(\frac{D}{p}\right) = \begin{cases} -1 \\ 0 \\ 1 \end{cases} \iff (p) = \begin{cases} (p) & p \text{ is inert} \\ \mathfrak{p}^2 & p \text{ ramifies} \\ \mathfrak{p}\bar{\mathfrak{p}} & p \text{ splits} \end{cases}$$

where D is the discriminant of the order $\mathcal{O}$ and $\left(\frac{D}{p}\right)$ is the Kronecker symbol. Recall that in the case of $p = 2$, the Kronecker symbol is defined as

$$\left(\frac{D}{2}\right) = \begin{cases} 0 & \text{if } 2 \mid D \\ -1 & \text{if } D \equiv \pm 3 \pmod{8} \\ 1 & \text{if } D \equiv \pm 1 \pmod{8} \end{cases}$$

Also note that the only time an ideal is not invertible is when one of its underlying primes ramifies and divides the conductor of the order, in other words $N(\mathfrak{p}) \mid f$. An ideal $\mathfrak{a}$ which is prime to the conductor may be written as $\mathfrak{a} = (t)\mathfrak{r}\mathfrak{s}$ where the primes dividing each of $\mathfrak{r}$ and $\mathfrak{s}$ ramify and split respectively.

4.1. Infrastructure

As we have already seen there is a structure involved in the continued fraction of ω in that there is a certain gap between two elements. This was termed the *infrastructure* by Shanks, [64] and it is utilized by

Definition. Let $\mathfrak{a}$ and $\mathfrak{b}$ be equivalent ideals of $\mathcal{O}$, that is $\mathfrak{a} = (\alpha)\mathfrak{b}$, for some $\alpha \in K^*$, then the *distance* from $\mathfrak{a}$ to $\mathfrak{b}$, denoted by $\delta(\mathfrak{a}, \mathfrak{b})$, is defined as

$$\partial(\mathfrak{a}, \mathfrak{b}) = \frac{1}{2} \log |\alpha/\bar{\alpha}|$$

which is defined modulo the least unit of norm 1.

Actually, this is Lenstra's definition of distance [33], Shanks originally considered $\delta(\mathfrak{a}, \mathfrak{b}) = \log |\alpha|$, which is defined modulo the regulator. We also define for a principal ideal $\mathfrak{a}$, $\partial(\mathfrak{a}) := \partial(1, \mathfrak{a})$.

We would like this operation to be a homomorphism, but unfortunately the best one can say is that for $\mathfrak{c}' = \mathfrak{a}\mathfrak{b} = (\alpha\beta)$ we can find a reduced ideal $\mathfrak{c} = (\alpha\beta\gamma)$ and so

$$\partial(\mathfrak{c}) = \frac{1}{2} \log |\alpha/\bar{\alpha}| + \frac{1}{2} \log |\beta/\bar{\beta}| + \frac{1}{2} \log |\gamma/\bar{\gamma}| \approx \partial(\mathfrak{a}) + \partial(\mathfrak{b}),$$

since $\log |\gamma/\bar{\gamma}|$ is small.

Returning now to the story of composition which was introduced earlier, the formula for composing ideals is given by

Theorem 10. If $\mathfrak{a} = \langle Q, \omega + P \rangle$ and $\mathfrak{b} = \langle Q', \omega + P' \rangle$ then their composite is given by $\mathfrak{ab} = \langle q, \omega + p \rangle$ where

$$\begin{aligned} G &= \gcd(Q, Q', P + P' + t) \\ q &= QQ'/G^2 \\ p &\equiv P \pmod{Q/G} \quad \text{and} \quad p \equiv P' \pmod{Q'/G} \\ (P - p)(P' - p) &\equiv n + tp + p^2 \pmod{q} \end{aligned}$$

Note that even if $\mathfrak{a}$ and $\mathfrak{b}$ are reduced it is possible that $\mathfrak{ab}$ will not be reduced, which means that one normally needs to reduce the ideal $\mathfrak{ab}$. Shanks provided an alternative for slightly alleviating this burden called NUCOMP [65]; it works by reducing while in the middle of composing. The works [77], [27] contain an exposition of this algorithm.

By applying the correspondence between ideals and forms, the above formulas provide a procedure for multiplying binary quadratic forms. We can also express composition in terms of matrices. Consider the matrices M and N ,

$$\begin{aligned} M &= \begin{pmatrix} x_h & -ny_h \\ y_h & x_h - ty_h \end{pmatrix} = \begin{pmatrix} x_h & x_{h-1} \\ y_h & y_{h-1} \end{pmatrix} \begin{pmatrix} 1 & P_{h+1} \\ 0 & Q_{h+1} \end{pmatrix} \\ &= \begin{pmatrix} a_0 & 1 \\ 1 & 0 \end{pmatrix} \cdots \begin{pmatrix} a_h & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & P_{h+1} \\ 0 & Q_{h+1} \end{pmatrix} \\ N &= \begin{pmatrix} x'_i & -ny'_i \\ y'_i & x'_i - ty'_i \end{pmatrix} = \begin{pmatrix} x'_i & x'_{i-1} \\ y'_i & y'_{i-1} \end{pmatrix} \begin{pmatrix} 1 & P'_{i+1} \\ 0 & Q'_{i+1} \end{pmatrix} \\ &= \begin{pmatrix} b_0 & 1 \\ 1 & 0 \end{pmatrix} \cdots \begin{pmatrix} b_i & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & P'_{i+1} \\ 0 & Q'_{i+1} \end{pmatrix} \end{aligned}$$

associated to two sequences in the continued fraction expansion of γ , where γ has trace t and norm n . The essence is that the product of two matrices of the ‘shape’ M and N is again of the same shape. Namely,

$$MN = \begin{pmatrix} X & -nY \\ Y & X - tY \end{pmatrix}. \quad (4.1)$$

This is a simple consequence of Fermat’s observation (1.2), specifically,

$$(x_h - \gamma y_h)(x'_i - \gamma y'_i) = (X - \gamma Y)$$

where $X = x_h x'_i - ny_h y'_i$ and $Y = x_h y'_i + x'_i y_h - ty_h y'_i$. By taking the decomposition of (4.1) we find

$$MN = \begin{pmatrix} c_0 & 1 \\ 1 & 0 \end{pmatrix} \cdots \begin{pmatrix} c_r & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & p \\ 0 & q \end{pmatrix} G$$

with p, q and G given by the rules above.

Example 1. Consider $D = 4 \cdot 271$, $\omega = \sqrt{271}$, $\mathcal{O}_K = [1, \omega]$, then

$$\begin{array}{l|l} 0 & \omega = 16 - (\overline{\omega} + 16) \\ 1 & (\omega + 16)/15 = 2 - (\overline{\omega} + 14)/15 \\ 2 & (\omega + 14)/5 = 6 - (\overline{\omega} + 16)/5 \\ 3 & (\omega + 16)/3 = 10 - (\overline{\omega} + 14)/3 \\ 4 & (\omega + 14)/25 = 1 - (\overline{\omega} + 11)/25 \\ 5 & (\omega + 11)/6 = 4 - (\overline{\omega} + 13)/6 \\ 6 & (\omega + 13)/7 = 1 - (\overline{\omega} + 4)/17 \end{array} \quad \begin{array}{l|l} 7 & (\omega + 4)/15 = 1 - (\overline{\omega} + 11)/15 \\ 8 & (\omega + 11)/10 = 2 - (\overline{\omega} + 9)/10 \\ 9 & (\omega + 9)/19 = 1 - (\overline{\omega} + 10)/19 \\ 10 & (\omega + 10)/9 = 2 - (\overline{\omega} + 8)/9 \\ 11 & (\omega + 8)/23 = 1 - (\overline{\omega} + 15)/23 \\ 12 & (\omega + 15)/2 = 15 - (\overline{\omega} + 15)/2 \\ & \vdots \end{array}$$

where the numbers on the left side of the line indicate which line of the expansion is being calculated. From lines 3 and 5 respectively we find the quadratic forms and ideals,

$$\mathcal{A} = (-3, 32, 5), \quad \mathfrak{a} = \langle 3, \omega + 16 \rangle \quad \text{and} \quad \mathcal{B} = (-6, 22, 25), \quad \mathfrak{b} = \langle 6, \omega + 11 \rangle.$$

Applying the composition formulas given in Theorem 10, we find the composites

$$\mathcal{AB} = (2, 22, -75) \quad \text{and} \quad \mathfrak{ab} = \langle 2, \omega + 11 \rangle$$

By reducing $\mathcal{AB}$ we find the form $\mathcal{C} = (2, 30, -23)$, whereas the ideal $\mathfrak{ab}$ is also equal to $\langle 2, \omega + 15 \rangle$ and so is already reduced. In term of matrices, the form $\mathcal{A}$ is associated to the product

$$\begin{pmatrix} 16 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 2 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 6 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 16 \\ 0 & 3 \end{pmatrix} = \begin{pmatrix} 214 & 3523 \\ 13 & 214 \end{pmatrix} = A$$

and the form $\mathcal{B}$ is associated to the product

$$\begin{pmatrix} 16 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 2 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 6 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 10 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 11 \\ 0 & 6 \end{pmatrix} = \begin{pmatrix} 2387 & 39295 \\ 145 & 2387 \end{pmatrix} = B.$$

The product AB is given by

$$\begin{aligned} AB &= \begin{pmatrix} 1021653 & 16818531 \\ 62061 & 1021653 \end{pmatrix} \\ &= \begin{pmatrix} 16 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 2 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 6 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 10 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 4 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix} \\ &\quad \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 2 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 3 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 17 \\ 0 & -2 \end{pmatrix} \cdot 3 \end{aligned}$$

Since 2 ramifies in $\mathcal{O}_K$, we have that the line 12 in the expansion is halfway and that the square of $\mathfrak{ab}$ has norm 1. In terms of the matrix product, this means

$$\frac{(AB/3)^2}{2} = \begin{pmatrix} x & 271y \\ y & x \end{pmatrix}, \quad x = 115974983600, \quad y = 7044978537$$

has determinant 1. Thus, $x + \omega y$ is the fundamental unit of $\mathcal{O}$. Explicitly, we have

$$\varepsilon = \frac{(214 + 13\omega)^2(2387 + 145\omega)^2}{18}.$$

The distance of the ideals is given by

$$\partial(\mathfrak{a}) = \log \left| \frac{214 + 13\omega}{214 - 13\omega} \right| \quad \text{and} \quad \partial(\mathfrak{b}) = \log \left| \frac{2387 + 145\omega}{2387 - 145\omega} \right|.$$

Since $\mathfrak{ab}$ is reduced, $\partial(\mathfrak{ab}) = \partial(\mathfrak{a}) + \partial(\mathfrak{b})$. Also note

$$\varepsilon = \frac{214 + 13\omega}{214 - 13\omega} \cdot \frac{2387 + 145\omega}{2387 - 145\omega}$$

so that $R(\mathcal{O}) = \partial(\mathfrak{a}) + \partial(\mathfrak{b})$.

5. FUNCTION FIELDS

It has long been known that there is an analogy between the cases of number fields and function fields. Apparently, Kronecker was aware of this as early as 1862, but it wasn't until Dedekind and Weber published their results in 1882 that it became commonly known. Modern references include [40], [30] both of which strive to deal with function fields and number fields as closely as possible.

Formally, for any field k , if L is a field of transcendence degree 1 over k then L is called a *function field over k* . In particular, if $Y^2 = f(X) \in k[X]$ where $f(X)$ is a squarefree polynomial then $L = k(X)(Y)$ is the field of functions of the curve given by the equation $g(X, Y) = Y^2 - f(X)$. Curves of the form $Y^2 = f(X)$ are called *hyperelliptic* if the degree of $f(X)$ is greater than 4. These (and the elliptic ones) will be of most interest herein.

When the degree of $f(X)$ is odd or its leading coefficient is not square then $k(X, \sqrt{f(X)})$ is called an imaginary quadratic function field and when the degree of $f(X)$ is even with a square as its leading coefficient then $k(X, \sqrt{f(X)})$ is called

a *real quadratic function field* (this will be the case we normally consider). This definition is in complete analogy with that for imaginary and real quadratic fields. Moreover, [41] proves that only finitely many quadratic imaginary extensions of $\mathbb{F}_q[X]$ have class number 1, in analogy to the number field case.

For a function f in the function field K over k , let $D_0 = \sum P_i$ where the sum is taken over all zeroes, P_i , of f in k . Similarly, $D_\infty = \sum Q_j$ where Q_j runs over the poles of f . Then the divisor of f , denoted by (f) is the formal sum $D_0 - D_\infty$. Likewise any formal sum of points, $P_1 + \dots + P_i - Q_1 - \dots - Q_j$ is called a *divisor*. A divisor of a function is called *principal*. The group of principal divisors of K is denoted by $P(K)$. For a divisor $D = \sum m_P P$ the *degree* of D is defined as $\sum m_P$. We denote the group of divisors of degree 0 by $\text{Div}_0(K)$. The group of principal divisors is a subgroup of $\text{Div}_0(K)$. The factor group given by $\text{Div}_0(K)/P(K)$ is called the *divisor class group* (of degree 0) of K .

This correspondence was utilized by Cantor [9] to demonstrate that adding divisors on hyperelliptic curves can be achieved by the same procedure as composition of ideals in quadratic fields, other references to this topic are [45], [57]. This correspondence would also suggest that there might exist a relationship between continued fractions and divisor class groups. Before embarking upon this let us examine continued fractions of functions, for more details see [79], [75].

Given a formal Laurent series $f(X) = \sum_{h=-m}^{\infty} a_h X^{-h}$ in a variable X^{-1} , its integer part is nothing other than its polynomial part, that is $a_{-m} X^m + \dots + a_{-1} X + a_0$. Then the continued fraction expansion of $f(X)$, namely $[a_0(X), a_1(X), \dots]$, can be found by using the same algorithm as that for numbers, we merely need to take

$$a_h(X) = \lfloor f_h(X) \rfloor$$

$$f_{h+1}(X) = \frac{1}{f_h(X) - a_h(X)} \quad \text{where } f_0(X) = f(X).$$

In other words, $f(x) = [a_0(X), \dots, a_h(X), f_{h+1}(X)]$, consequently the $f_i(X)$'s are the *complete quotients* of $f(X)$. All the formal identities given in §2 will also hold. Truncations, $[a_0(X), \dots, a_h(X)]$ yield rational functions $\frac{x(X)}{y(X)}$ and

$$\deg(y_{h+1}(X)) = \deg(a_{h+1}(X)) + \deg(y_h(X)).$$

Furthermore,

Proposition 11. *If $x(X)$ and $y(X)$ are relatively prime polynomials then*

$$\deg(y(X)f(X) - x(X)) < -\deg(y(X)) \quad (5.1)$$

if and only if $x(X)/y(X)$ is a convergent to $f(X)$.

Proof. (All letters, except f , given here will represent polynomials in X). We already know that for a convergent x/y of f we have,

$$\deg(y_h f - x_h) = -\deg y_{h+1} = -(\deg a_{h+1} + \deg y_h) < -\deg y_h.$$

Conversely, suppose that x/y is not a convergent and that (5.1) is satisfied. Then there is some h such that $\deg y_{h-1} \leq \deg y < \deg y_h$ with y not a constant multiple of y_{h-1} . Then,

$$y = ay_h + by_{h-1}$$

$$x = ax_h + bx_{h-1}$$

is solvable for some nonzero polynomials $a(X)$, $b(X)$ since $x_h y_{h-1} - x_{h-1} y_h = \pm 1$. Hence

$$yf - x = a(fy_h - x_h) + b(fy_{h-1} - x_{h-1})$$

of degrees $(\deg a - \deg y_{h+1})$ and $(\deg b - \deg y_h)$ respectively. If these two degrees were equal we would have $\deg ay_h = \deg by_{h-1}$ and so

$$\deg b - \deg y_h = \deg a - \deg y_{h-1} > \deg a - \deg y_{h+1}.$$

This impossibility means that we must have $\deg(yf - x) = \deg b - \deg y_h$ and $\deg b - \deg y_h < -\deg y$, by assumption. On the other hand, $\deg b \geq \deg y_h - \deg y_{h-1}$ implies that $\deg b - \deg y_h \geq -\deg y$, which is a contradiction. ■

The case we are primarily interested in is when $f(X)$ is the square root of a polynomial. In analogy with the number field case, our interest will be solely on real quadratic function fields, that is $\deg(f(X))$ is even and its leading coefficient is a square. The main feature now is that periodicity of $f(X)$ is only guaranteed if the base field is finite, namely

Proposition 12. *If k is a finite field and $f(X) \in k[X]$ is of even degree then $\sqrt{f(X)}$ is periodic.*

Proof. By Proposition 11 there exists infinitely many distinct rational functions $x(X)/y(X)$ which satisfy $\deg(y(X)\sqrt{f(X)} - x(X)) < -\deg y(X)$. If $f(X)$ has degree $2(g+1)$ then this yields

$$\deg(x(X)^2 - f(X)y(X)^2) \leq g.$$

Since k is a finite field, there only finitely many polynomials $r(X) \in k[X]$ which satisfy $\deg r(X) \leq g$. By the box principle there exist distinct polynomials in $k[X]$ satisfying

$$x(X)^2 - f(X)y(X)^2 = x'(X)^2 - f(X)y'(X)^2 = m(X),$$

and

$$x(X) \equiv x'(X) \pmod{m(X)}, \quad y(X) \equiv y'(X) \pmod{m(X)}.$$

Hence, $p(X) = (xx' - fyy')/m^2$ and $q(X) = (xy' - x'y)/m^2$ are polynomials in $k[X]$ which satisfy

$$p(X)^2 - f(X)q(X)^2 = \kappa, \quad \kappa \in k.$$

In other words, Pell's equation is satisfied, and $p(X) + \sqrt{f(X)}q(X)$ is a unit of $k[X, \sqrt{f(X)}]$. ■

When considering formal power series over a field of characteristic 0 the partial quotients will typically be of degree 1 and furthermore the heights of these polynomials are expected to increase at an exponential rate. Since such an expansion does not expect to be periodic we say that if $\sqrt{f(X)}$ is periodic then $k(X, \sqrt{f(X)})$ is an *exceptional* function field. The results on quadratic irrationals given earlier now apply. We merely note that an element $f(X, Y) \in k(X, Y)$ is *reduced* if $\deg f > 0$ and $\deg \bar{f} < 0$, where conjugation represents the mapping $f(X) \mapsto -f(X)$.

5.1. Some examples

Now suppose that $Y^2 = f(X)$ has degree $2(g+1)$, then a typical complete quotient of Y may be represented as $(Y + P_h)/Q_h$ where we have

$$\deg P_h = g+1 \quad \text{and} \quad \deg Q_h \leq g.$$

Thus the end of a period is equivalent to $\deg a_h = g+1$ and $\deg Q_h = 0$.

If the expansion is periodic then there exists a non-trivial unit in the function field, but now the regulator is taken to be the degree of the fundamental unit, in other words it is discrete. Furthermore, Proposition 5 implies that the regulator must be the sum of the degrees of the partial quotients of a primitive period.

For example,

$$Y^2 = X^6 + 2X^2 + X + 1$$

is in $\mathbb{Q}[X]$ and the continued fraction expansion of Y is

$$\left[x^3, x - \frac{1}{2}, -4x - 6, \frac{-1}{16}x + \frac{1}{8}, \frac{128}{5}x + \frac{512}{25}, \frac{125}{1488}x - \frac{6025}{69192}, \right. \\ \left. \frac{-3217428}{8125}x - \frac{812642742}{105625}, \frac{-1373125}{203694562323}x + \frac{20593600625}{158474369487294}, \right. \\ \left. \frac{7991216816924103}{1570855000}x + \frac{279952799594094347}{77757322500}, \dots \right] \quad (5.2)$$

On the other hand, we can consider Y to be defined over $\mathbb{F}_p[X]$ for various primes p , and these expansions must be periodic. For example, the continued fraction expansion of Y over $\mathbb{F}_5[X]$ is given by

h	$a_h(X)$	$P_h(X)$	$Q_h(X)$
0	X^3	0	1
1	$X + 2$	X^3	$2X^2 + X + 1$
2	$X + 4$	$X^3 + 3X + 2$	$2X^2 + 2X + 2$
3	$4X + 2$	$X^3 + 2X + 1$	$3X^2 + X$
4	$3X^2 + 3X + 3$	$X^3 + 4$	$4X + 1$
5	$4X + 2$	$X^3 + 4$	$3X^2 + X$
6	$X + 4$	$X^3 + 2X + 1$	$2X^2 + 2X + 2$
7	$X + 2$	$X^3 + 3X + 2$	$2X^2 + X + 1$
8	$2X^3$	X^3	1

demonstrating that $(4X + 1)|(X^6 + 2X^2 + X + 1)$ over $\mathbb{F}_5[X]$ and we also observe that the regulator is 11. Over $\mathbb{F}_7[X]$ one finds,

h	$a_h(X)$	$P_h(X)$	$Q_h(X)$
0	X^3	0	1
1	$X + 3$	X^3	$2X^2 + X + 1$
2	$3X + 1$	$X^3 + 4X + 3$	$3X^2 + 6X + 6$
3	$3X + 1$	$X^3 + 6X + 3$	$3X^2 + 6X + 1$
4	$6X + 2$	$X^3 + 3X + 5$	$5X^2 + 3X + 4$
5	$5X + 4$	$X^3 + 6X + 3$	$6X^2 + 5X + 5$
6	$2X + 6$	$X^3 + 4X + 3$	$X^2 + 4X + 4$
7	X^3	X^3	2
8	$2X + 6$	X^3	$X^2 + 4X + 4$
9	$5X + 4$	$X^3 + 4X + 3$	$6X^2 + 5X + 5$
10	$6X + 2$	$X^3 + 6X + 3$	$5X^2 + 3X + 4$
11	$3X + 1$	$X^3 + 3X + 5$	$3X^2 + 6X + 1$
12	$3X + 1$	$X^3 + 6X + 3$	$3X^2 + 6X + 6$
13	$X + 3$	$X^3 + 4X + 3$	$2X^2 + X + 1$
14	$2X^3$	X^3	1

which has a regulator 9. There is a result due to Yu [87] and van der Poorten [74] which states that if $\sqrt{f(X)}$ is periodic over $\mathbb{Q}$ then $R_q q^a = R_p p^b$, where R_i is the regulator over $\mathbb{F}_i[X]$ and a, b are some integers. Thus, in the example given above, if $\mathbb{Q}[X, Y]$ were to have a nontrivial unit its regulator would need to satisfy $5^a 11 = 7^b 9$ which is clearly impossible and this shows that the ... in (5.2) does continue forever.

The last expansion demonstrates the concept of a ‘quasi-period’. Observe that a unit is produced at line 7 but the expansion is not fully periodic until line 14, in fact the lines 8-14 are a twist of lines 1-7. More precisely, if the continued fraction

expansion of $\sqrt{f(X)}$ is periodic with $Q_h = \kappa$ a unit then

$$\sqrt{f(X)} = [a_0(X), \overline{a_1(X), \dots, a_h(X), \kappa a_1(X), \kappa^{-1} a_2(X), \dots, \kappa a_h(X)}],$$

it is also evident that in such a case h must be odd. For more on quasi-periodicity, see [6] and [79].

5.2. Continued fractions and torsion divisors

The relationship between divisors and continued fractions can be stated as: the divisor $(\infty^+ - \infty^-)$ on the Jacobian of the curve $\mathcal{C} : Y^2 = f(X)$ is torsion if and only if the continued fraction expansion of $\sqrt{f(X)}$ is periodic. In the case of elliptic curves, every quartic can be transformed into a cubic which enables one to give a precise relationship between regulators of quartics and torsion on elliptic curves. Adams and Razar [1] showed that under the transformation,

$$X = \frac{w+b}{v-a}, \quad Y = 2v + a - \left(\frac{w+b}{v-a}\right)^2,$$

the curve

$$E : w^2 = v^3 + Av + b$$

with rational point $P = (a, b)$ is birationally equivalent to

$$Q : Y^2 = X^4 - 6aX^2 - 8bX + c$$

with P being sent to ∞ (where $c = -4A - 3a^2$). Furthermore, the regulator of Y is equal to the order of the point P on E .

Details of elliptic curves can be found in [67], [24], but all we need to know is that any elliptic curve defined over a field k of characteristic not equal to 2 or 3 can be represented in the so-called Weierstrass form,

$$E : y^2 = x^3 + ax + b, \quad a, b \in k$$

and that the addition of points on the curve defines a group law where the point at infinity acts as the identity. We use the notation $E(k)$ to denote the group of k -rational points on E . Furthermore, an admissible change of variables is any transformation

$$x = s^2x' + r \quad y = s^3y' + ts^2x' + u, \quad r, s, t, u \in k,$$

with u invertible in k . Moreover, Mazur [43] has completely determined all possible torsion structures of elliptic curves over the rationals.

Theorem 13 (Mazur's theorem). *For a nonsingular elliptic curve over $\mathbb{Q}$ the torsion subgroup $E(\mathbb{Q})_{\text{tors}}$ of the group of rational points $E(\mathbb{Q})$ is one of*

$$\mathbb{Z}/m\mathbb{Z} \quad \text{for } 1 \leq m \leq 10 \quad \text{or} \quad m = 12$$

or

$$\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2m\mathbb{Z} \quad \text{for } 1 \leq m \leq 4.$$

Each of these cases was already known to occur. Kubert [31] gives the following table in which Tate-normal form is used to describe an elliptic curve. Namely, if

$$y^2 + (1-c)xy - by = x^3 - bx^2$$

then the point $(0, 0)$ is a torsion point of maximal order where b and c are given in the following table.

$E(\mathbb{Q})_{tors}$	b	c
$\mathbb{Z}_4$	t	0
$\mathbb{Z}_5$	t	t
$\mathbb{Z}_6$	$t(t+1)$	t
$\mathbb{Z}_7$	$t^2(t-1)$	$t(t-1)$
$\mathbb{Z}_8$	$(t-1)(2t-1)$	$\frac{(t-1)(2t-1)}{t}$
$\mathbb{Z}_9$	$t^2(t-1)(t^2-t+1)$	$t^2(t-1)$
$\mathbb{Z}_{10}$	$\frac{t^2(t-1)(2t-1)}{(t^2-3t+1)^2}$	$\frac{-t(t-1)(2t-1)}{t^2-3t+1}$
$\mathbb{Z}_{12}$	$\frac{t(2t-1)(2t^2-2t+1)(3t^2-3t+1)}{(t-1)^4}$	$\frac{-t(2t-1)(3t^2-3t+1)}{(t-1)^3}$
$\mathbb{Z}_2 \times \mathbb{Z}_4$	$\frac{\frac{1}{16}(4t-1)(4t+1)}{-(t-1)^2(t-5)}$	0
$\mathbb{Z}_2 \times \mathbb{Z}_6$	$\frac{-(t-1)^2(t-5)}{(t^2-9)^2}$	$\frac{-2(t-5)}{t^2-9}$
$\mathbb{Z}_2 \times \mathbb{Z}_8$	$\frac{(2t_1)(8t^2+4t+1)}{(8t^2-1)^2}$	$\frac{(2t-1)(8t^2+4t+1)}{2t(4t+1)(8t^2-1)^2}$

(We exclude the cases of torsion 2 and 3 since they are trivial and of little interest to us here). Tran [69] used the transformation of Adams and Razar to list all quartics whose continued fraction expansion is periodic. Unfortunately, this situation is unique to curves of genus less than 2; there is no nice equivalence for curves of degree greater than 4. Essentially the problem is that a sextic (or more) can only be transformed into a quintic (or more) if it possesses a Weierstrass point (in other words it must have a rational linear factor).

In summary, we have:

Proposition 14. *Let C be the curve $Y^2 = f(X)$ where $f(X) \in k[X]$ is of degree $2g+2$ and is squarefree and let J be its Jacobian. Then the following are equivalent:*

- (i) *The continued fraction expansion of Y is periodic,*
- (ii) *$k[X, Y]$ contains a non-trivial unit,*
- (iii) *$(\infty^+ - \infty^-)$ has finite order in J ,*
- (iv) *The continued fraction expansion of Y contains more than one partial quotient of degree $g+1$.*

Lastly, we will also use the big oh notation. Namely, if f and g are functions of some continuous or integral variable, with g being a positive function, then $f = O(g)$ means that $|f| \leq cg$ for some c , this is also sometimes denoted by $f \ll g$. Likewise, $|f| \geq cg$ is denoted by $f \gg g$. If we have the situation where $f \ll g$ and $f \gg g$ then we write $f = \Theta(g)$, in other words f and g are of the same order of magnitude.

Chapter II. History of sleepers, beepers and creepers

Examples like $\sqrt{x^2 + 1} = [x, \overline{2x}]$ where the continued fraction expansion can be explicitly stated have long been known. Certainly, at least, they are present in Perron's book [58] with various results known. The first in-depth study of such matters was conducted by Schinzel in the 1960s, [60], [61]. In [60] he uses the two results

$$\limsup_{n \rightarrow \infty} lp(\sqrt{n^2 + h}) < \infty \iff h \mid 4n^2$$

and that every quadratic is birationally equivalent to $n^2 + h$ to show that

Theorem 15 (Quadratic Sleepers). *If $f(n) = an^2 + bn + c$ with $a, b, c \in \mathbb{Z}$ and $\Delta = b^2 - 4ac \neq 0$ then*

$$\limsup_{n \rightarrow \infty} lp(\sqrt{f(n)}) < \infty$$

if and only if $a = \square$ and $\Delta \mid 4(2a, b)^2$.

Kaplansky, [28], called such functions *sleepers*. In other words, a sleeper is a function $f(x) \in \mathbb{Q}[x]$ which satisfies

$$\limsup_{n \rightarrow \infty} lp(\sqrt{f(n)}) < \infty.$$

As notation, we will let $\sqrt{f(X)} = [a_0(X), a_1(X), \dots]$ with $a_i(X) \in \mathbb{Q}[X]$ represent the continued fraction expansion of $\sqrt{f(X)}$ over $\mathbb{Q}[X]$, and $\sqrt{f(x)} = [a_0, a_1, \dots]$ with $a_i \in \mathbb{Z}$ represent the continued fraction expansion of $\sqrt{f(x)}$ over $\mathbb{Z}$. In short, capital indeterminates represent expansions over function fields while small letters denote expansions over the integers. Even for functions possessing periodic expansions the two can be distinct (in fact it is rare for them to coincide). For example

$$f(x) = 9x^2 + 8x + 2$$

has over the integers the expansion

$$\sqrt{f(x)} = [3x + 1, \overline{2, 1, 3x, 1, 2, 2(3x + 1)}].$$

While over $\mathbb{Q}[X]$, $f(X)$ has the expansion

$$\sqrt{f(X)} = [3X + \frac{4}{3}, \overline{9(3X + \frac{4}{3}), 2(3X + \frac{4}{3})}].$$

We shall say that any real quadratic function field possessing a nontrivial unit is an *exceptional function field*. The matter of sleepers was dealt with by Schinzel [61],

Theorem 16 (Schinzel's theorem). *For $f(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_0$ we have*

$$\limsup_{x \rightarrow \infty} lp(\sqrt{f(x)}) < \infty$$

if and only if $\mathbb{Q}(\sqrt{f(X)})$ is an exceptional function field and if

$$\varepsilon = p_k + q_k \sqrt{f(X)}$$

represents the unit corresponding to the primitive period (note that this unit is not fundamental in the case of a quasi-period) then $p_k \in \frac{1}{2}\mathbb{Z}$.

It is not necessary that $q_k \in \frac{1}{2}\mathbb{Z}$ but if m is the denominator of q_k then $m^2 \mid f(X)$, whereupon $\varepsilon \in \frac{1}{2}\mathbb{Z}[\sqrt{f(X)}]$.

Some further observations are that the condition $x \rightarrow \infty$ shouldn't be weakened to $x \rightarrow \infty$ for all $x \in S$ for some set S since this would forfeit the condition that n was even and certainly is not providing any generalization.

The condition that $2p_k \in \mathbb{Z}$ is called the *Schinzel condition*. Thus finding sleepers consists of finding exceptional real quadratic function fields which have a unit satisfying the Schinzel condition.

A modern rendition of quadratic sleepers is given in [80]. Nowadays all the possible torsion groups of elliptic curves are known by Mazur's theorem, but back in the 60's this was only a conjecture. Consequently, Schinzel was only able to remark

... in particular that $LP\sqrt{f}$ can take the values 1, 2, 3, 4, 6, 8, 10, 14, 18, 22 and possibly also 5, 7, 9, 11 (I have not verified this) and, if the conjecture of Nagell [55] is true, no other values.

As it happens, Tran [69], was able to show that all of the possible values occur except 9 and 11. But this doesn't finish the story of sleepers for quartics. This merely determines the curves possessing a unit, it still remains to be decided whether it is possible that a particular congruence class of X exists where this unit is located in $\frac{1}{2}\mathbb{Z}[X]$ or not. We do this in §7.

There is also the question of finding functions of degree greater than 4 possessing a non-trivial unit. At the time Schinzel remarked

For polynomials f of degree > 4 I do not know any method which would always lead to the solution of problem P_1

(P_1 refers to determining functions possessing a periodic continued fraction expansion.) Indeed, even now such a method still eludes us. The only known practical method follows from Flynn. In §8 we review this method (in the language of continued fractions) since it will come in handy later on in the work on creepers.

6. CREEPERS AND BEEPERS

It has long been known that there exist parametric families of discriminants which yield continued fraction expansions of linear length. The earliest example seems to arise from Nyberg [56] who examined

$$D_n = (x^n + (x \pm 1)/2)^2 \mp x^n.$$

Being written in Norwegian seemed to restrict the entering of this result into the collective knowledge of western mathematicians. Consequently, it was not until Shanks [62] noticed the so-called Shanks' sequence that others learned of these sequences. In [62], Shanks examines the class numbers of $n^2 - 2^{2k+1}$. For $n^2 - 8$ he finds,

$n^2 - 8$	$h(n^2 - 8)$
17	1
41	1
73	1
113	1
$\vdots$	
4217	1
4481	3
6553	1
7561	1

$$\begin{array}{|c|c|} \hline 8273 & 1 \\ \hline 8641 & 1 \\ \hline \end{array}$$

Naturally, 4481 stood out to him and he realized that it was an instance of

$$S_n = (2^n + 3)^2 - 8.$$

Furthermore, he showed that one expects $h(S_n) \rightarrow \infty$, and so creepers were born. A quick method for calculating the class number of S_n is given in [63]. The continued fraction expansion of Shanks' sequence, where $\omega = (1 + \sqrt{D_n})/2$, is

$$\begin{array}{rcl} \omega & = & 2^{n-1} + 1 - (\bar{\omega} + 2^{n-1}) \\ \frac{\omega + 2^{n-1}}{2^n} & = & 1 - \frac{(\bar{\omega} + 2^{n-1} - 1)}{2^n} \\ \frac{\omega + 2^{n-1} - 1}{2} & = & 2^{n-1} - \frac{(\bar{\omega} + 2^{n-1})}{2} \\ \frac{\omega + 2^{n-1}}{2^{n-1}} & = & 2 - \frac{(\bar{\omega} + 2^{n-1} - 1)}{2^{n-1}} \\ & \vdots & \\ \frac{\omega + 2^{n-1}}{2} & = & 2^{n-1} - \frac{(\bar{\omega} + 2^{n-1} - 1)}{2} \\ \frac{\omega + 2^{n-1} - 1}{2^n} & = & 1 - \frac{(\bar{\omega} + 2^{n-1})}{2^n} \\ \frac{\omega + 2^{n-1}}{1} & = & 2^n + 1 - \frac{(\bar{\omega} + 2^{n-1} - 1)}{1} \end{array}$$

Then $\omega = [2^{n-1} + 1, 1, 2^{n-1}, 2, 2^{n-2}, \dots, 2^{n-j}, 2^{j+1}, \dots, 2^{n-1}, 1, 2^n + 1]$, which means that $lp(\omega) = 2n + 1$. Taking $\varepsilon_n = \prod_{h \in I} (\omega + P_h)/Q_h$ gives the fundamental unit

$$\varepsilon_n = \left(\frac{2^n + 1 + \sqrt{D}}{2} \right) \left(\frac{2^n + 3 + \sqrt{D}}{4} \right)^n.$$

A vital point is then noted by Shanks. He was interested in Gauss' class number problem for real quadratic fields, so he wants to be able to find large class numbers in order to exclude them. He remarks

If these S_n were the only primes $n^2 - 8$ with $h > 1$ we could conclude, heuristically, that "almost all" primes $n^2 - 8$ had $h = 1$ and therefore Gauss's idea of a fixed limiting ratio < 1 would be inapplicable to this subset of positive determinants. However, there are other counterexamples. The examples $d = 8713, 9281$ and 2089 mentioned above have periods for their principal forms of a much more intricate character than those for the S_n and, a priori, there is no reason why similar examples cannot occur for $n^2 - 8$.

It just so happens that we can explain two of the other occurrences as creepers (these are the generalizations of Shanks sequence which we will be studying). Namely,

$$D_n = (3 \cdot 2^n + 1)^2 - 4 \cdot 2^n \quad \text{and} \quad E_n = (3 \cdot 2^n - 3)^2 + 4 \cdot 2^n$$

and $D_5 = 9281, E_4 = 2089$. By the results to be presented we see shall discover that each of the sequences of discriminants D_n and E_n also have period length

linear in n just like Shanks sequence S_n . However, the discriminant 8713 is a little different. The sequence of discriminants given by

$$F_n = (2^n - 3^3 \cdot 2^k - 1)^2 + 4 \cdot 2^n$$

is a creeper. However, to be part of this family it is necessary that $n \equiv 3 \pmod{6}$, whereas $F_4 = 8713$. It does not seem that 8713 belongs to any special sequence of discriminants.

Several people have since generalized Shanks' sequence. However the next development was from the opposite direction, constructing units which can't be small.

In [86], Yamamoto demonstrated that for

$$D_n = (rx^n + x - 1)^2 + 4rx^n$$

with $r \nmid x - 1$ then $R(D_n) \gg (\log D_n)^3$ which is the current record (Halter-Koch, [21], attempted to extend Yamamoto's method to demonstrate the existence of $R(D_n) \gg (\log D_n)^4$, for more about this see page 60). This time distribution of journals rather than language seems to be the problem as Hendy produced [22] unaware of Yamamoto's paper. Hendy wished to construct discriminants such that the sequence Q_h/Q_0 satisfies

$$Q_{2r} = c^r Q_0, \quad Q_{2r+1} = c^{n-r} Q_0.$$

Such discriminants are given by

$$D_n = \left(\frac{Q_0 m c^n}{2} + k \right)^2 + Q_0^2 c^n$$

The story continues with L. Bernstein, [4], [5], who was aware of the work of Yamamoto but apparently not of Shanks nor Hendy. He uses brute force to one-by-one run through a series of cases. Some examples include:

$$\begin{aligned} D_k &= (A^k + a)^2 + A, & A &= 2a + 1, \\ D_k &= (A^k - a)^2 + A, & A &= 2a + 1, \\ D_k &= (A^k + a + 1)^2 - A, & A &= 2a + 1, \\ D_k &= (A^k + A - 1)^2 + 4A, & A &= 2^d b, \quad b \text{ odd}, \quad d \geq 1, \quad k \geq 2. \end{aligned}$$

Bernstein's results are then generalized by Azuhata, [2], [3], and Levesque and Rhin in [39] (this seems to be the first paper which is aware of the work of Nyberg). These results are synthesised in a systematic manner by H. Williams in [82]. Later more general results, indeed the most general known prior to this work, were given by H. Williams in [83] and by Van der Poorten in [72] who uses the automata introduced by Raney [59] to ease the pain. This generalization is

$$D_n = \left(qrx^n + \frac{\mu(x^k - \lambda)}{q} \right)^2 + 4\lambda rx^n \quad (6.1)$$

with $\mu, \lambda \in \{-1, 1\}$ and $rq \mid x^k - \lambda$. Prior to this present work, (6.1) represented the edge of the knowledge.

Mollin and Williams in [52], [53] generalize the idea of Hendy and demonstrate that if D_n possesses 3 consecutive Q_h 's which are a power of x for every n then D_n must be of the form (6.1).

6.1. Beepers

The principle feature of these families is that $lp(\omega_n) = an + b$ for some $a, b \in \mathbb{Z}$. But these are not the only families possessing linear period lengths. Following the bounds produced in [25] regarding the difference in period lengths of $\sqrt{D}$ and $\frac{1+\sqrt{D}}{2}$, K. Williams and Buck wanted to show that the bounds were sharp, which

they did in [8]. To do this they required infinite families with a determinable period length. For all the cases required, bar one, they found a suitable creeper. But in the other case they used the family

$$D_n = (2F_{6n+1} + 1)^2 + 8F_{6n} + 4$$

where F_n is the n th Fibonacci number. It has the period length $lp(\omega_n) = 6n + 1$.

This example is vastly different to all the preceding ones because all the partial quotients (excluding the obvious) are bounded. Namely,

$$\omega_n = [F_{6n+1} + 1, \overline{1}, \dots, 1, \overline{2F_{6n+1} + 1}]$$

Thus $R(D_n) = \Theta(\log D_n)$. However, all the previous examples have $R(D_n) = \Theta((\log D_n)^2)$, (because there exists a $Q_h = x^\nu$ with ν independent of n).

Following a challenge from K. Williams, Van der Poorten was able to “produce such examples at will” in [73]. These families have been called *beepers* in honour of the beer won by Van der Poorten. Further work on beepers has been completed by Mollin and Cheng in [46], [47], [48], [49]. The point is to notice that a beeper is essentially an infinite family of selected sleepers.

6.2. Kreepers

We define a *creeper* to be an infinite family of discriminants D_n , polynomially parametrized, which satisfy

$$lp(\omega_n) = an + b \quad \text{with } a, b \in \mathbb{Q} \quad \text{and} \quad R(D_n) = \Theta(n^2).$$

A letter from Kaplansky [28] to Mollin, K. Williams and H. Williams attempted to provide a complete description of quadratic creepers with some conjectures. It was this letter which prompted the current investigation, a copy of this letter may be found as an appendix. Kaplansky coined the terms “sleepers”, a sequence of discriminants whose period lengths are bounded, and “creepers”, a sequence whose lengths gently go to infinity, and “leapers”, the generic discriminants whose period lengths increase exponentially. First, he restricted his interest to discriminants of the form $am^{2n} + bm^n + c$, where a, b, c , and m are fixed integers; denoting this sequence by $(a, b, c; m)$ or just (a, b, c) . By giving several conjectures he hoped to provide a complete analysis of creepers, much as Schinzel had done with sleepers. Among his conjectures were,

1. If (a, b, c) is a creeper then so is $(a, -b, c)$.
2. If (a, b, c) is a creeper then a and c are squares.

The second conjecture is a vital starting point, and it helps to assume its truth. Following this, we call an infinite sequence of discriminants $\{D_n\}$ a *kreeper* if

- $D_n = A^2x^{2n} + Bx^n + C^2$ where A, B, C are rational integers, and x is a positive integer.
- $lp(\omega_n) = an + b$ where a and b are rational numbers.
- In the principal cycle there exists an element whose norm is x^g for some g fixed independently of n .

Note that the existence of some $Q_h = x^g$ implies $R(D_n) = \Theta(n^2)$. In other words,

Theorem 17. *Every kreeper is also a creeper.*

Proof. In any kreeper we have reduced, principal ideals with norms:

$$x^g, x^{2g}, \dots, x^{hg} \quad \text{where } h \gg n.$$

By Proposition 5,

$$\begin{aligned}\varepsilon &\geq \frac{\omega + P}{x^g} \cdot \frac{\omega + P''}{x^{2g}} \cdots \frac{\omega + P^{(h)}}{x^{hg}}, \\ &\geq cx^{n-g} \cdot cx^{n-2g} \cdots cx^{n-hg} \quad \text{for some } c \in \mathbb{Q},\end{aligned}$$

thus, $R(D_n) \gg n^2$. Conversely,

$$\begin{aligned}\varepsilon &= \frac{\omega + P_1}{Q_1} \cdots \frac{\omega + P_{an+b}}{Q_{an+b}} < (dx^n)^{an+b} \quad \text{for some } d \in \mathbb{Q}, \\ R(D_n) &\ll n^2.\end{aligned}$$

Hence, $R(D_n) = \Theta(n^2)$ and so every creeper is also a creeper. ■

The principle object of this thesis is to determine all keepers. This will be done in Chapters V and VII.

As a concluding note, the concepts of creepers and beepers have been combined by Madden. In [42], he explicitly constructs continued fraction expansions of the shape

$$\begin{aligned}\sqrt{D_n} &= \left[a_0, A_0, \vec{N}, B_{k-1}, A_1, \vec{N}, B_{k-2}, A_2, \vec{N}, \dots, \right. \\ &\quad \left. A_{k-1}, \vec{N}, B_0, a_0, \dots, 2a_0 \right] \quad (6.2)\end{aligned}$$

where $\vec{N} = [a_1, a_2, \dots, a_{f(n)}]$ is a beeper and

$$\sqrt{D_n} = [a_0, A_0, B_{k-1}, A_1, B_{k-2}, A_2, \dots, 2a_0]$$

is a creeper from the family (6.1). We won't delve into these matters here since we want to examine discriminants given by polynomials and the beeper aspect prevents this from occurring. In effect, Madden's construction are to keepers what beepers are to sleepers. Namely, Madden selects a sequence of discriminants from different keepers so that (6.2) is satisfied. In other words, these examples are incorporated within the entire set of keepers and so do not provide larger regulators. This does not detract from their interest.

Chapter III. Sleepers

In Chapter II we saw that Schinzel demonstrated that finding sleepers is the same as finding exceptional function fields which satisfy the Schinzel condition. Now let us suppose that we have a function, $f(X)$, such that $\sqrt{f(X)}$ is quasi-periodic. By Proposition 6 of Chapter I we know that if

$$p_k^2 - q_k^2 f(X) = -\kappa,$$

then

$$p_{2k+1}^2 - q_{2k+1}^2 f(X) = 1,$$

and

$$p_{2k+1} + q_{2k+1} \sqrt{f(X)} = \frac{(p_k + q_k \sqrt{f(X)})^2}{\kappa}.$$

Hence,

$$2p_{2k+1} = \frac{2p_k^2 + 2q_k^2 f(X)}{\kappa} = \frac{4p_k^2 + 2\kappa}{\kappa}$$

The Schinzel condition is that $2p_{2k+1} \in \mathbb{Z}$. From above, this is clearly equivalent to $\kappa | 4p_k^2$. Thus we have demonstrated,

Proposition 18. *If $\sqrt{f(X)}$ is quasi-periodic with a fundamental unit of norm $-\kappa$ then the Schinzel condition is met if and only if $\kappa \in \mathbb{Q}^2$.*

From the work of Tran [69] we know that the quartics with quasi-period 9 and 11 can never satisfy the Schinzel condition and so don't possess any numerical sleeper subfamilies. To review why this is so, given the continued fraction expansion we have either $Q_9 = \kappa$ or $Q_{11} = \kappa$. In either case, to find a situation where there is no quasi-period we need to determine the cases of $\kappa \in \mathbb{Q}^2$. For if $\kappa = s^2$ then the continued fraction expansion of $\sqrt{f(X)}/s$ will have $Q_{9,11} = 1$ and hence the primitive period becomes 9 or 11 respectively. But it turns out that in these two cases it is impossible to find $\kappa = s^2$. In each case respectively, we have

$$Q_9 = -4t(t-1)(t^2 - 3t + 1) \quad \text{or} \quad Q_{11} = \frac{4t(2t-1)^2(3t^2 - 3t + 1)^3}{(t-1)^4}$$

so we need to solve the equations,

$$s^2 = -t^4 + 4t^3 - 4t^2 + t \quad \text{or} \quad s^2 = t(t-1)(3t^2 - 3t + 1).$$

These can be easily transformed into

$$y^2 = x^3 - x^2 - x \quad \text{or} \quad y^2 = x^3 + x^2 + x$$

both of which contain one rational point, that being $(0, 0)$ of order 2 which leads to values of $t = 0, 1$ which yield singular curves.

This shows that in these two cases there is no general solution, but there will still exist some values of t such that the Schinzel condition can be satisfied. For example, for quasi-period 9,

$$Y^2 = (X^2 - (4t^6 - 16t^5 + 8t^4 + 8t^3 - 4t + 1))^2 + 32t^3(t-1)(2t-1)(t^2 - 3t + 1)(X + 2t^3 - 2t^2 - 2 + 1)$$

we have that by the methods yet to be presented that the Schinzel condition can be met when $|t^2 - 3t + 1| = 1$, that is $t = 2, 3$ (the values $t = 0, 1$ yield singular curves).

7. SLEEPERS FROM CURVES OF GENUS 1

Given Schinzel's theorem and Mazur's theorem, one may ask which elliptic curves with a torsion divisor at infinity yield a family of sleepers. Quadratics were easily dealt with by Schinzel, but at the time Mazur's theorem was merely a conjecture and he was therefore unable to provide complete results for genus 1 curves.

However, [79] and [69] explicitly list all quartics with an exceptional real quadratic function field. The form in which we shall write our quartics will normally be as

$$Y^2 = (X^2 + u(t))^2 + 2v(t)(X + w(t)) \quad u(t), v(t), w(t) \in \mathbb{Z}[t]. \quad (7.1)$$

The easiest curves to begin with are the ones corresponding to torsion 5 (torsion 3 and 4 are somewhat trivial). Namely,

$$Y^2 = (X^2 - (t^2 - 6t + 1))^2 + 32t(X - (t - 1)).$$

The partial quotients are then

h	$a_h(X)$
0	$X^2 - t^2 + 6t - 1$
1	$\frac{1}{16t}(X + t - 1)$
2	$4(X - (t + 1))$
3	$\frac{1}{16t}(X + t - 1)$
4	$2(X^2 - t^2 + 6t - 1)$

The simple approach is to note that if all the partial quotients are integers then necessarily the unit is also integral and so by Schinzel's theorem it must be a sleeper, (in fact we could also allow a single $\frac{1}{2}$ to float about without getting into trouble).

Clearly, as long as $X \equiv 1 - t \pmod{16t}$ then all the partial quotients will be integers and hence so must the unit and therefore the family will be a set of sleepers. In detail, when we take $X \equiv 1 - t \pmod{16t}$, that is $X = 1 - t + 16tZ$, we get

h	$a_h(Z)$
0	$256t^2Z^2 + (-32t^2 + 32t)Z + 4t$
1	Z
2	$64tZ - 8t$
3	Z
4	$512t^2Z^2 + (-64t^2 + 64t)Z + 8t$

and this is the actual expansion of $\sqrt{g(Z)}$ over $\mathbb{Z}$ where

$$g(Z) = 2^{16}t^4Z^4 + 2^{14}t^3(t - 1)Z^3 + 2^{10}t^2(t^2 + 1)Z^2 + 2^8t^2(t - 3)Z - 2^4t(3t - 4).$$

Using this idea to attack the other cases, the continued fraction expansion of Y as given in (7.1), commences as

$$\begin{aligned} Y &= X^2 + u(t) - (\overline{Y} + X^2 + u(t)) \\ \frac{Y + X^2 + u(t)}{2v(t)(X + w(t))} &= \frac{X - w(t)}{2v(t)} - \frac{(\overline{Y} + X^2 + u_2(t))}{2v(t)(X + w(t))} \end{aligned}$$

Detailing more of the expansion becomes tricky, but not impossible see [78], but for our purposes we will work with the first 2 steps and cross our fingers.

Since we only need $\varepsilon \in \frac{1}{2}\mathbb{Z}[X]$, $\frac{1}{2}$'s occurring in partial quotients are not overly frightening. Hence the above expansion suggests considering

$$X \equiv w(t) \pmod{v(t)}.$$

It turns out that this will suffice in most cases. In fact the only cases which fail to satisfy this are:

- Family 6 ($\mathbb{Z}_8$)
- Family 6 ($\mathbb{Z}_{12}$)
- Family 18 ($\mathbb{Z}_{10}$)
- Family 22 ($\mathbb{Z}_{12}$)

(Here we have adopted the notation used in [69] where Family $m(\mathbb{Z}_n)$ represents the function whose expansion has a primitive period of length m , arising from the torsion group of order n).

The last two we know, from Proposition 18, can never satisfy the Schinzel condition. The first two can be relatively easily explained. If we examine Family 6($\mathbb{Z}_8$)

$$Y^2 = \left(X^2 - (20t^4 - 40t^3 + 28t^2 - 8t + 1)\right)^2 - 32t^2(t-1)^2(2t-1)^2(X + (2t-1)^2) \quad (7.2)$$

then we see that the expansion is

h	$a_h(X)$
0	$X^2 - (20t^4 - 40t^3 + 28t^2 - 8t + 1)$
1	$\frac{1}{-16t^2(t-1)^2(2t-1)^2} (X - (2t-1)^2)$
2	$4(2t-1)^2 (X - (2t-1)^2)$
3	$\frac{1}{-32t^2(t-1)^2(2t-1)^4} (X^2 - (20t^4 - 40t^3 + 28t^2 - 8t + 1))$

The difference between this curve and the others is the presence of a common factor between $v(X)$ and $w(X)$. This causes only a slight problem.

Evidently, from above, trying to make each partial quotient an integer will immediately fail, so a little compromise is needed. First, it eases things somewhat to divide the above expansion by $4t(t-1)(2t-1)^2$. This does not alter the unit but lessens the requirements upon the partial quotients. Now we are looking at

h	$a_h(X)$
0	$\frac{1}{4t(t-1)(2t-1)^2} (X^2 - (20t^4 - 40t^3 + 28t^2 - 8t + 1))$
1	$\frac{1}{-4t(t-1)} (X - (2t-1)^2)$
2	$\frac{1}{t(t-1)} (X - (2t-1)^2)$
3	$\frac{1}{-8t(t-1)(2t-1)^2} (X^2 - (20t^4 - 40t^3 + 28t^2 - 8t + 1))$

In other words, we seek a solution of

$$X \equiv (2t-1)^2 \pmod{4t(t-1)}$$

and

$$X^2 \equiv 20t^4 - 40t^3 + 28t^2 - 8t + 1 \pmod{8t(t-1)(2t-1)^2}$$

One such solution is $X \equiv 1 + 2t(t-1) \pmod{4t(t-1)(2t-1)}$.

The curve of Family 6($\mathbb{Z}_{12}$) is given by,

$$Y^2 = \left(X^2 + 12t^8 - 48t^7 + 12t^6 + 132t^5 - 216t^4 + 156t^3 - 60t^2 + 12t - 1\right)^2 + 32t^3(t-1)^3(2t-1)^3(3t^2 - 3t + 1)(X + (2t-1)^3).$$

It is treated in a similar manner to (7.2) with a solution being

$$X \equiv (2t-1)^3 + 32t^3(t-1)^3(3t^2 - 3t + 1) \pmod{16t^3(t-1)^3(2t-1)^3(3t^2 - 3t + 1)}.$$

7.1. *Smaller solutions*

As an aside, we remark that considering $X \equiv w(t) \pmod{v(t)}$ is far from best possible. While we are only interested in showing whether a family of sleepers exists or not, one can sometimes do better. For example, consider the curve with torsion 9,

$$Y^2 = (X^2 - (t^6 - 6t^5 + 9t^4 - 10t^3 + 6t^2 + 1))^2 + 32t^2(t-1)(t^2-t+1)(X - (t^3 - t^2 - 1)).$$

The expansion of $Y/64t^2$ is

h	$a_h(X)$
0	$\frac{X^2 - (t^6 - 6t^5 + 9t^4 - 10t^3 + 6t^2 + 1)}{8t}$
1	$\frac{(X + t^3 - t^2 - 1)}{2t(t-1)(t^2-t+1)}$
2	$\frac{(X - (t^3 - t^2 + 1))}{2t}$
3	$\frac{(X + t^3 - 3t^2 + 2t - 1)}{2t(t-1)}$
4	$\frac{(X - (t^3 - 3t^2 + 4t - 1))}{2(t^2 - t + 1)}$
5	$\frac{(X + t^3 - 3t^2 + 2t - 1)}{2t(t-1)}$
6	$\frac{(X - (t^3 - t^2 + 1))}{2t}$
7	$\frac{(X + t^3 - t^2 - 1)}{2t(t-1)(t^2-t+1)}$
8	$\frac{(X^2 - (t^6 - 6t^5 + 9t^4 - 10t^3 + 6t^2 + 1))}{4t}$

If we let $C = -(t^3 - t^2 - 1)$, then we have

$$\begin{aligned} C - (t^3 - t^2 + 1) &= -2t^2(t-1) \\ C + t^3 - 3t^2 + 2t - 1 &= -2t(t-1) \\ C - (t^3 - 3t^2 + 4t - 1) &= 2(t-1)(t^2 - t + 1) \\ C^2 - (t^6 - 6t^5 + 9t^4 - 10t^3 + 6t^2 + 1) &= 4t^2(t-1)(t^2 - t + 1) \end{aligned}$$

which means that all the partial quotients are integers, except a_0 which is in $\frac{1}{2}\mathbb{Z}[X]$, under the restriction $X \equiv C \pmod{2t(t-1)(t^2-t+1)}$ and hence the unit is also in $\frac{1}{2}\mathbb{Z}[X]$. Consequently,

$$X \equiv -(t^3 - t^2 - 1) \pmod{2t(t-1)(t^2-t+1)}$$

will provide a numerical sleeper (which is a little more general than the earlier solution).

7.2. *A detailed expansion*

Next we note that although our method of attack was to make the partial quotients integers we can still find an acceptable unit even though some of the partial

quotients are not integers (and worse than just $\frac{1}{2}$'s). For example the curve with torsion 8 and even period length,

$$Y^2 = (X^2 + 4t^4 + 4t^3 - 16t^2 + 8t - 1)^2 + 32t^3(t-1)(2t-1)(X - (2t^2 - 4t + 1))$$

has the expansion,

h	$a_h(X)$
0	$X^2 + 4t^4 + 4t^3 - 16t^2 + 8t - 1$
1	$\frac{(X + 2t^2 - 4t + 1)}{16t^3(t-1)(2t-1)}$
2	$4t(X - (2t^2 - 2t + 1))$
3	$\frac{(X - (2t - 1))}{16t^2(t-1)(2t-1)}$
4	$\frac{4(2t-1)(X - (2t-1))}{t}$
5	$\frac{(X - (2t^2 - 2t + 1))}{16(t-1)(2t-1)^2}$
6	$\frac{4(2t-1)(X + 2t^2 - 4t + 1)}{t^2}$
7	$\frac{(X^2 + 4t^4 + 4t^3 - 16t^2 + 8t - 1)}{32t(t-1)(2t-1)^2}$

We discover that upon taking $X \equiv -(2t^2 - 4t + 1) \pmod{16t^2(t-1)(2t-1)}$ that some of the partial quotients are not integral. Even worse, these denominators can't be removed by judicious multiplication by rationals. However, it happens that although our method of attack has failed the unit is still in $\frac{1}{2}\mathbb{Z}[X]$. One can see this by determining the unit and taking the above congruence class but that is not particularly enlightening. If we make the substitution $X = -1 + 4t - 2t^2 + 16t^3(t-1)(2t-1)Z$ and then the dilations, $Z \mapsto Z/8$, $Y \mapsto Y/64$ and then consider $W^2 = Y^2/2^{18}t^2(t-1)^2$ (which doesn't affect the unit),

$$W^2 = \left(\frac{1}{2}t^5(t-1)^2(2t-1)Z^2 - \frac{1}{2}t^2(t-1)(2t^2 - 4t + 1)Z + \frac{1}{2}t(t-1)\right)^2 + t(t-1)(t^3(t-1)Z - (2t^2 - 4t + 1)/(2t-1)).$$

The continued fraction expansion of W over $\mathbb{Q}[Z]$ is

h	$a_h(Z)$
0	$\frac{1}{2}t^5(2t-1)(t-1)^2Z^2 - \frac{1}{2}t^2(t-1)(2t^2 - 4t + 1)Z + \frac{1}{2}t(t-1)$
1	$t(2t-1)Z$
2	$t^3(t-1)Z - (t-1)$
3	$t^2(2t-1)Z - 1$
4	$t(t-1)(2t-1)Z - \frac{(t-1)}{t}$
5	$t^4Z - t$
6	$(t-1)(2t-1)Z$
7	$t^6(t-1)(2t-1)Z^2 - t^3(2t^2 - 4t + 1)Z + t^2$

Moreover, the periodic expansion of W is symmetric. The only non-integral partial quotient is $a_4(Z)$, where

$$P_4(Z) = \frac{1}{2}t^5(t-1)^2(2t-1)Z^2 - \frac{1}{2}t^2(t-1)(2t^2-4t+1)Z - \frac{t(t-1)}{2(2t-1)}$$

$$Q_4(Z) = t^4(t-1)Z - \frac{t(t^2-3t+1)}{2t-1}$$

Thus the fourth line of the expansion reads

$$\frac{W + P_4}{Q_4} = t(2t-1)(t-1)Z - \frac{(t-1)}{t} - \frac{(\overline{W} + P_5)}{Q_4}$$

By applying Lemma 2 from page 8 we can expand this as an element in the two variables Z, t rather than just Z . Specifically, the next partial quotients are the expansion of

$$\frac{(t-1)}{t}(t(2t-1)Z-1)$$

which equals

$$[(2t-1)(t-1)Z-1, t].$$

Note that we do not need to worry about the parity of the length of the expansion since we are working over function fields and not rationals. Thus the next element in the expansion is

$$\frac{-Q_4}{(-1)(\overline{W} + P_5)t^2} - \frac{j}{t} \quad \text{where} \quad j \equiv \frac{(-1)^2}{(t-1)[t(2t-1)Z-1]} \pmod{t}.$$

So $j = -1$ and this element becomes

$$\frac{W + P_5}{-Q_5 \cdot t^2} + \frac{1}{t} = -t^2Z + \frac{1}{t} - \frac{1}{t} - \frac{(\overline{W} + P_6)}{-Q_5 \cdot t^2}$$

and the expansion safely continues as

$$\frac{t^2(W + P_6)}{-Q_6} = -t^2(t-1)(2t-1)Z - \frac{(\overline{W} + P_7)t^2}{-Q_6}$$

$$\frac{(W + P_7)}{-Q_7 \cdot t^2} = -t^4(t-1)(2t-1)Z^2 + t(2t^2-4t+1)Z - 1 - \frac{(\overline{W} + P_8)}{-Q_7 \cdot t^2}.$$

Also, $P_7 = P_8$ still, so this is halfway and all the partial quotients are now integral, except $a_0 \in \frac{1}{2}\mathbb{Z}[Z]$. Moreover, by the preceding symmetry we also have that $a_9, \dots, a_{15}$ are also integral.

In summary, we take:

h	$a_h(Z)$
0	$\frac{1}{2}t^5(2t-1)(t-1)^2Z^2 - \frac{1}{2}t^2(t-1)(2t^2-4t+1)Z + \frac{1}{2}t(t-1)$
1,9	$t(2t-1)Z$
2,10	$t^3(t-1)Z - (t-1)$
3,11	$t^2(2t-1)Z - 1$
4,12	$(2t-1)(t-1)Z - 1$
5,13	t
6,14	$-t^2Z$
7,15	$-t^2(t-1)(2t-1)Z$
8	$-t^4(t-1)(2t-1)Z^2 + t(2t^2-4t+1)Z - 1$

While this is not the regular expansion over $\mathbb{Q}[Z]$, we can still consider

$$\begin{pmatrix} a_0 & 1 \\ 1 & 0 \end{pmatrix} \cdots \begin{pmatrix} a_{15} & 1 \\ 1 & 0 \end{pmatrix} = \begin{pmatrix} x & x' \\ y & y' \end{pmatrix}$$

and find a solution of $x^2 - W^2 y^2 = 1$ with $x \in \frac{1}{2}\mathbb{Z}[Z]$, so that Schinzel's condition is met.

In conclusion,

Theorem 19. *For a quartic $f(X) = (X^2 + u(t))^2 + 2v(t)(X + w(t))$, with a periodic continued fraction $\sqrt{f(X)}$, the congruence class $x \equiv w(t) \pmod{v(t)}$ yields a numerical sleeper $\sqrt{f(x)}$ unless (i) the period length of $\sqrt{f(X)}$ is 18 or 22, in which case there is no general solution, or (ii) the period length of $\sqrt{f(X)}$ is 6 and the regulator is either 8 or 12. In this case, a solution is given by*

$$x \equiv 1 + 2t(t-1) \pmod{4t(t-1)(2t-1)}$$

when the regulator is 8 and

$$x \equiv (2t-1)^3 + 32t^3(t-1)^3(3t^2-3t+1) \pmod{16t^3(t-1)^3(2t-1)^3(3t^2-3t+1)}$$

when the regulator is 12.

8. FINDING EXCEPTIONAL FUNCTION FIELDS

Knowing Schinzel's theorem, if we wish to determine all numerical sleepers we need to determine all functions whose square root has a periodic continued fraction expansion. Having dealt with the cases of genus 0 and 1 our attention turns to genus 2 and higher. Unfortunately, no results like Mazur's are known for these cases. In fact there is not even an upper bound for the size of torsion (some bounds are known in special cases, like Silverberg's [66] which says that when the Jacobian of a genus 2 curve is of CM-type then the order of a torsion point is no more than $2^3 \cdot 3 \cdot 5 \cdot 7 \cdot 13 \cdot 17$) nor is there a single $m \in \mathbb{N}$ for which it is known that this is not the order of a torsion divisor on a genus 2 curve.

In the genus 1 case, the torsion groups were originally found by using the addition formulae for adding points on elliptic curves. However, when the genus is greater than 1 such formulae are not nearly as simple, leaving this approach unfeasible. Van der Poorten [78] shows that the theory of continued fractions can be used to explicitly construct all curves of any genus with a specified regulator. However, beyond genus 1 the complexity of the computation required is quite intimidating once the regulator becomes large.

The only other known method originates from Flynn ([18], [19] and also Chapter 8 of [10]) and was used by Leprévost [38], [37], [36], [35], [34] and Elkies [15] successively to determine all currently known rational torsion divisors of simple genus 2 Jacobians. A word here is needed to distinguish between simple and split Jacobians of genus 2 curves. A split Jacobian of genus 2 is one which is isogenous to the product of two Jacobians of genus 1. The state of the art for these matters is [23], where a torsion divisor of order 63 on a curve of genus 2 is found. The idea there basically is to use the torsion groups of the two underlying elliptic curves.

Leprévost, [37], gives the following condition for simplicity,

Proposition 20. *The Jacobian of a curve of genus 2 is simple if there exists a prime l of good reduction such that the Galois group of the characteristic polynomial of Frobenius of the reduced curve is the 8-element dihedral group D_4 .*

We shall not be perturbed by whether our curves are simple or not (it turns out that they all are). We are now going to explain Flynn's procedure (and Leprévost's and Elkies extensions) for determining rational torsion divisors. There will be nothing new here, except for the interpretation in terms of continued fractions. But we include the explanation since it will become significant later on in our studies of creepers, in particular during Chapter VIII.

The idea is to write down a set of equations satisfied by the curve,

$$\begin{aligned} Y^2 &= (A_1(X))^2 + 4t_1X^{e_1}(X-1)^{f_1} \\ &= (A_2(X))^2 + 4t_2X^{e_2}(X-1)^{f_2} \\ &= (A_3(X))^2 + 4t_3X^{e_3}(X-1)^{f_3} \end{aligned}$$

If we set $D_1 = \{(0, A_1(0)), \infty^-\}$, $D_2 = \{(1, A_1(1)), \infty^-\}$ and $D_3 = \{\infty^-, \infty^-\}$ where the notation $\{(x, y), (u, v)\}$ means the divisor class containing $(x, y) + (u, v) - (\infty^+ + \infty^-)$. Then we find some φ_i for $i = 1, 2, 3$ as a linear combination of these 3 divisors, so that

$$\begin{pmatrix} (\varphi_1) \\ (\varphi_2) \\ (\varphi_3) \end{pmatrix} = \begin{pmatrix} A & B & C \\ D & E & F \\ G & H & I \end{pmatrix} \begin{pmatrix} D_1 \\ D_2 \\ D_3 \end{pmatrix} \quad (8.1)$$

Then the order of the class containing D_1, D_2, D_3 divides the determinant of the matrix in (8.1).

Following Flynn, Leprevost was able to determine examples of torsion of every order up to 29 (see [38]) and 30 in an unpublished paper with Everett Howe. The example of torsion 29 (at the time the record) is

$$\begin{aligned} Y^2 &= (2X^3 + X^2 - 4X + 2)^2 - 8X^3(X-1)^2 = A_1(X)^2 - 8X^3(X-1)^2 \quad (8.2) \\ &= (2X^3 - 3X^2 + 4X - 2)^2 + 8X^2(X-1)^3 = A_2(X)^2 + 8X^2(X-1)^3 \\ &= (2X^3 - X^2 - 2)^2 + 16X(X-1) = A_3(X)^2 + 16X(X-1) \end{aligned}$$

and by taking $\varphi_1 = Y + A_1(X)$, $\varphi_2 = (Y + A_2(X))/X^2$, and $\varphi_3 = Y - A_3(X)$ we find

$$\begin{pmatrix} (\varphi_1) \\ (\varphi_2) \\ (\varphi_3) \end{pmatrix} = \begin{pmatrix} 3 & 2 & -2 \\ -2 & 3 & 0 \\ 1 & 1 & -3 \end{pmatrix} \cdot \begin{pmatrix} D_1 \\ D_2 \\ D_3 \end{pmatrix}$$

Thus D_1, D_2, D_3 are divisors of order 29. (This determinantal procedure is not the most efficient since it only says the torsion order divides the determinant and so there is more work to do when the determinant is composite. It is in fact easier just to use Cantor's procedure and explicitly compute the multiples of the divisor).

The continued fraction expansion of (8.2) can be found in the Tables on page 137.

Flynn also noticed that there exist families of curves whose Jacobian possess torsion divisors of order g^2 as $g \rightarrow \infty$, [19]. Namely the curve,

$$Y_g^2 = (A(X))^2 - tX^g(X-1)^{g+1} \quad (8.3)$$

where

$$A(X) = \frac{1}{2}(X^{g+1} - t(X-1)^g - X^g(X-1))$$

and the divisor $\{(0, A(0)), \infty\}$ has torsion order of $2g^2 + 2g + 1$. We shall explain such examples later on, it turns out that these are examples of function field keepers.

A word of warning is useful here. The continued fraction expansion finds the order of the torsion of the divisor $(\infty^+ - \infty^-)$ so, for example, when we consider Leprevost's example of a curve with torsion 21,

$$Y^2 = 4X^6 - 12X^5 + 13X^4 - 6X^3 + 3X^2 - 2X + 1.$$

We find the expansion

h	$a_h(X)$	$P_h(X)$	$Q_h(X)$
0	$2X^3 - 3X^2 + X$	0	1
1	$2X - 1$	$2X^3 - 3X^2 + X$	$2X^2 - 2X + 1$
2	$-X + \frac{1}{2}$	$2X^3 - 3X^2 + 3X - 1$	$-4X(X - 1)$
3	$-2X + 1$	$2X^3 - 3X^2 - X + 1$	$-2X(X - 1)$
4	$X - \frac{1}{2}$	$2X^3 - 3X^2 + 3X - 1$	$4X^2 - 4X + 2$
5	$4(2X^3 - 3X^2 + X)$	$2X^3 - 3X^2 + X$	$\frac{1}{2}$

which means that $(\infty^+ - \infty^-)$ has order 7. The point of order 21 is in fact $(0, 1)$, so we send this point to ∞^+ and obtain the curve

$$w^2 = v^6 - 2v^5 + 3v^4 - 6v^3 + 13v^2 - 12v + 4$$

which yields the expansion

h	$a_h(v)$	$P_h(v)$	$Q_h(v)$
0	$v^3 - v^2 + v - 2$	0	1
1	$\frac{1}{4}v$	$v^3 - v^2 + v - 2$	$8v(v - 1)$
2	$4(v + 1)$	$v^3 - v^2 - v + 2$	$\frac{1}{2}v^2 - v + 1$
3	$\frac{-1}{8}(v^2 - v + 1)$	$v^3 - v^2 + v + 2$	$-16v$
4	$-4(v^2 + 1)$	$v^3 - v^2 + v - 2$	$\frac{-1}{2}(v - 1)$
5	$\frac{1}{4}(v + 1)$	$v^3 - v^2 + v$	$8(v - 1)^2$
6	$2(v - 1)$	$v^3 - v^2 - 3v + 2$	v^2
7	$\frac{-1}{2}v$	$v^3 - v^2 + 3v - 2$	$-4v(v - 1)$
8	$-v$	$v^3 - v^2 - 3v + 2$	$-2v(v - 1)$
9	$v - 1$	$v^3 - v^2 + 3v - 2$	$2v^2$
10	$\frac{1}{2}(v + 1)$	$v^3 - v^2 - 3v + 2$	$4(v - 1)^2$
11	$-2(v^2 + 1)$	$v^3 - v^2 + v$	$-(v - 1)$
12	$\frac{-1}{4}(v^2 - v + 1)$	$v^3 - v^2 + v - 2$	$-8v$
13	$2(v + 1)$	$v^3 - v^2 + v + 2$	$v^2 - 2v + 2$
14	$\frac{1}{2}v$	$v^3 - v^2 - v + 2$	$4v(v - 1)$
15	$v^3 - v^2 + v - 2$	$v^3 - v^2 + v - 2$	2

displaying a regulator of order 21.

8.1. Further examples

Also note that there is no loss in generality in taking our remainders to be X and $X - 1$ since we have two degrees of freedom in dilating and translating. However, there is no need for there to be only two remainders, i.e. there could well be a third term $X + r$ where the value of r can no longer be arbitrarily selected. The difficulty now is that there are a lot more possible combinations for our equations, and that these are harder to solve. While we could try to continue with this third, or even a fourth factor it is easier to take just two equations so that Y is not completely determined and then search over the possible curves. For example,

$$\begin{aligned} Y^2 &= A_1(X)^2 + 4t_1X(X - 1)(X + r) \\ &= A_2(X)^2 + 4t_2X(X - 1)(X + r)^2 \end{aligned}$$

In order to have the ideals compose appropriately we require

$$X(X - 1)(X + r) \mid (A_1(X) + A_2(X)).$$

This immediately entails that $r = 1$,

$$A_1(X) = X^3 + c_1X + d_1 \quad \text{and} \quad A_2(X) = X^3 + c_2X + d_2.$$

Equating coefficients then yields

$$\begin{aligned} Y^2 &= (X^3 + cX + d)^2 + 4(1 + c + d)X(X - 1)(X + 1) \\ &= (X^3 - (2 + c)X - d)^2 + 4(1 + c)X(X - 1)(X + 1)^2. \end{aligned}$$

Thus the reduced representation is

$$Y^2 = (X^3 + cX + 2(c + 1) - d)^2 - 4(c^2 + (2 - d)c + 1 - d)(X + 1).$$

From here one can either run through the values of c and d or impose further restrictions upon the curve such as

$$Y^2 = A_3(X)^2 + 4t_3X(X - 1)^3$$

Some examples are the case of regulator 39 for $c = 2$, $d = 1$; another is $c = -2$, $d = 1$ which has regulator 27; and $c = 8$, $d = 3$ with regulator 11. The case $c = 2$, $d = 1$, namely

$$Y^2 = (X^3 + 2X + 5)^2 - 24(X + 1)$$

is particularly interesting. The expansion of Y is

h	$a_h(X)$	$P_h(X)$	$Q_h(X)$
0	$X^3 + 2X + 5$	0	1
1	$\frac{-1}{12}X^2 + \frac{1}{12}X - \frac{1}{4}$	$X^3 + 2X + 5$	$-24(X + 1)$
2	$-6X - 6$	$X^3 + 2X + 1$	$\frac{-1}{3}X(X - 1)$
3	$\frac{-1}{18}X + \frac{1}{9}$	$X^3 - 4X - 1$	$-36(X + 1)^2$
4	$-9X$	$X^3 - 2X - 3$	$\frac{-2}{9}(X - 1)(X + 1)$
5	$\frac{-1}{9}X + \frac{1}{9}$	$X^3 + 3$	$-18(X^2 + X + 2)$
6	$\frac{-3}{2}X^2 - \frac{3}{2}X - \frac{9}{2}$	$X^3 + 2X - 7$	$\frac{-4}{3}(X - 1)$
7	$\frac{-1}{3}X + \frac{1}{3}$	$X^3 + 2X + 1$	$-6X(X + 1)$
8	$-X$	$X^3 - 4X - 1$	$-2(X + 1)(X - 1)$
9	$\frac{-1}{2}X^2 - 1$	$X^3 + 2X + 1$	$-4X$
10	$\frac{-2}{3}X$	$X^3 + 2X - 1$	$-3X^2$
11	$\frac{-3}{4}X + \frac{3}{4}$	$X^3 - 2X + 1$	$\frac{-8}{3}X(X + 1)$
12	$\frac{-4}{3}X + \frac{8}{3}$	$X^3 - 1$	$\frac{-3}{2}(X + 1)^2$
13	$\frac{-1}{8}X - \frac{1}{4}$	$X^3 - 10X + 5$	$-16(X - 1)^2$
14	$8X + 8$	$X^3 + 4X - 1$	$\frac{1}{4}X(X - 1)$
15	$\frac{1}{32}X - \frac{3}{64}$	$X^3 - 6X + 1$	$32(2X^2 + 3X - 1)$
16	$\frac{128}{3}X - 64$	$X^3 + \frac{1}{2}X + \frac{1}{2}$	$\frac{3}{128}(2X^2 + 3X - 1)$

This line is halfway in the expansion and so the regulator of Y is 39. This is currently the largest known regulator of any quartic.

Elkies was able to produce examples of torsion of orders 30, 32, 34, 39 and 40. The 40 example highlights the limitations of using continued fractions. The curve is given by

$$Y^2 = 3X^5 + 19X^4 + 44X^3 + \frac{185}{4}X^2 + 22X + 4$$

with the obvious rational non-Weierstrass points $(-2, 1)$, $(0, 2)$, $(-1, 1/2)$ and the rational Weierstrass point $(-4/3, 0)$. If we consider any of the rational points and their corresponding continued fraction expansions then we obtain a regulator of 10. But when we consider the divisor $P = (-2, 1) + (-4/3, 0)$ we find it has order 40. Using Cantor's notations, the multiples mP of P are:

m	A	B
1	$(3X + 4)(X + 2)$	$-3/2X - 2$
2	$9(X + 2)^2$	$-3/2X - 2$

3	$-(X+1)(X+2)/3$	$X/2$
4	$-(3X+1)(X+4)/3$	$-3/2X-2$
5	$-3(X+1)(X+2)$	$-3/2X-2$
6	$-X/3$	2
7	$-3(4X^2+9X+4)/4$	$-X/8-1/2$
8	$4(X+1)^2$	$-X/2-1$
9	$-(3X^2+9X+4)/3$	$-X/2-2/3$
10	$X(X+1)$	$-5X/2-2$
11	$-(3X^2+6X+2)$	$-X/2-3/2$
12	$X^2/4$	$11X/2+2$
13	$-3X(X+2)/4$	$-X/2-2$
14	$-3X(3X+4)/4$	$-3X/2-2$
15	$-27X(X+2)/4$	$-3X/2-2$
16	$-4(X+1)/27$	$1/2$
17	$-27(X^2-X-4)/4$	$-23X/2-18$
18	$X(X+1)/9$	$-3X/2-2$
19	$9(X+2)$	-1
20	$9(3X+4)$	0

and since this last line is 2-torsion we have that the entire divisor is 40 torsion.

This same approach can be used to find points of low height. Unlike Leprévost's technique, which guarantees a unit will be produced, we have not restricted the curve as much. Consequently, it is possible that the continued fraction expansion will commence as if it were going to find a unit (that is, the heights of the partial quotients are small) but if some further equation fails to be satisfied, the unit is not constructed; that is the expansion is not periodic. From this, Elkies was able to find the record for lowest heights on a genus 2 curve. Specifically,

$$Y^2 = X^6 - 2X^5 - 4X^4 + 2X^3 + \frac{37}{4}X^2 - \frac{15}{2}X + \frac{9}{4}.$$

The beginning of the expansion of Y can be found in the tables on page 137. By examining the regulators over $\mathbb{F}_p[X]$ we find that the continued fraction expansion is non-periodic. As was stated earlier one expects the partial quotients of a non-periodic expansion to be linear, but remarkably in this case one finds

h	$a_h(X)$	$P_h(X)$	$Q_h(X)$
0	$X^3 - X^2 - \frac{5}{2}X - \frac{3}{2}$	0	1
1	$\frac{-2}{15}X^2 + \frac{2}{15}X + \frac{1}{3}$	$X^3 - X^2 - \frac{5}{2}X - \frac{3}{2}$	$-15X$
4	$\frac{25}{3}X^2 - \frac{125}{6}$	$X^3 - X^2 - \frac{5}{2}X + \frac{3}{2}$	$\frac{6}{25}(X-1)$
11	$\frac{-81}{500}X^2 + \frac{81}{250}X + \frac{81}{1000}$	$X^3 - X^2 - \frac{5}{2}X + \frac{7}{2}$	$\frac{-1000}{81}(X+1)$
26	$\frac{-625}{3072}X^2 - \frac{625}{1536}X - \frac{4375}{6144}$	$X^3 - X^2 - \frac{5}{2}X - \frac{9}{2}$	$\frac{-6144}{625}(X-3)$
36	$\frac{625}{18}X^2 - \frac{625}{36}X - \frac{6875}{72}$	$X^3 - X^2 - \frac{5}{2}X + \frac{9}{4}$	$\frac{18}{625}(2X-1)$

Furthermore, while the heights of the partial quotients are still increasing their rate of increase is considerably slower than expected. Three of the partial quotients of degree 2 are a manifestation of the following equations

$$\begin{aligned}
Y^2 &= \left(X^3 - X^2 - \frac{5}{2}X - \frac{3}{2}\right)^2 - 15X \\
&= \left(X^3 - X^2 + \frac{5}{2}X - \frac{3}{2}\right)^2 - 10X^3(X-1) \\
&= \left(X^3 - X^2 + \frac{1}{2} - \frac{3}{2}\right)^2 - 6X(X+1)(X-1)^2 \\
&= \left(X^3 - X^2 - \frac{5}{2}X + \frac{3}{2}\right)^2 - 6X^2(X-1)
\end{aligned}$$

But the other two linear $Q_h(X)$'s which correspond to the rational points $(3, 6)$ and $(\frac{1}{2}, \frac{7}{8})$, are unexpected and this is what makes this example special. Elkies has shown using Coleman-Chabauty bounds that there are no more rational points on the curve.

8.2. The elliptic case

These ideas also apply in the elliptic case. In particular, suppose we have

$$\begin{aligned} Y^2 &= A_1(X)^2 + 4t_1(X - 1) \\ &= A_2(X)^2 + 4t_2X(X - 1) \\ &= A_3(X)^2 + 4t_3X^2 \end{aligned}$$

with the conditions $(X - 1) \mid (A_1(X) + A_2(X))$ and $X \mid (A_2(X) + A_3(X))$ so that the ideals compose nicely. Then easy algebra yields

$$Y^2 = (X^2 - tX + (2t - 1))^2 + 4t(t - 1)(X - 1)$$

and the continued fraction expansion of Y is

h	$a_h(X)$	$P_h(X)$	$Q_h(X)$
0	$X^2 - tX + (2t - 1)$	0	1
1	$\frac{1}{2t(t - 1)}(X - (t - 1))$	$X^2 - tX + (2t - 1)$	$4t(t - 1)(X - 1)$
2	$\frac{2(t - 1)(X - t)}{1}$	$X^2 - tX - 1$	$\frac{1}{t - 1}X$
3	$\frac{1}{2(t - 1)^2}(X - t)$	$X^2 - tX + 1$	$4(t - 1)^2X$
4	$\frac{2(t - 1)^2}{t}(X - (t - 1))$	$X^2 - tX - 1$	$\frac{t}{(t - 1)^2}(X - 1)$
5	$\frac{1}{2(t - 1)^3}(X^2 - tX + (2t - 1))$	$X^2 - tX + (2t - 1)$	$4(t - 1)^3$

and these are the quartics of torsion order 6. The elliptic case is very special however. Namely, any quartic can be written

$$Y^2 = A(X)^2 + 4t(X + r)$$

and what is more, from the continued fraction expansion we know that $\deg(Q_h(X))$ is 1 except at the end of a period, hence each step of the continued fraction expansion gives an equation,

$$Y^2 = A_{i+1}(X)^2 + 4t_{i+1}(X + r_i)(X + r_{i+1})$$

where $(X + r_i) \mid (A_i(X) + A_{i+1}(X))$, until we find some $r_i = r_{i+1}$ which indicates halfway in the expansion. Thus this method will find every quartic with a periodic continued fraction expansion.

The case of odd torsion (i.e. no quasi-period) corresponds not to $r_i = r_{i+1}$ but rather to $(X + r_{i+1}) \mid Y^2$, so that $Q_{i+1} \mid Y^2$ and this point must be halfway in the expansion. Note that since the expansion has no quasi-period such a place is necessary. For example

$$Y^2 = (X^2 + 39)^2 + 384(X - 3)$$

has torsion 9 and its expansion is

h	$a_h(X)$	$P_h(X)$	$Q_h(X)$
0	$X^2 + 39$	0	1
1	$(X + 3)/192$	$X^2 + 39$	$384(X - 3)$

2	$4(X-5)$	X^2-57	$(X+5)/2$
3	$(X-1)/64$	X^2+7	$128(X+1)$
4	$8(X-3)/3$	X^2-9	$3(X+3)/4$
5	$(X-1)/64$	X^2-9	$128(X+1)$
6	$4(X-5)$	X^2+7	$(X+5)/2$
7	$(X+3)/192$	X^2-57	$384(X-3)$
8	$2(X^2+39)$	X^2+39	1

with $(X+3) \mid Y^2$ representing halfway.

As a further illustration, the extreme case of torsion 12, is nothing more than the solution of the following equations, again there would be no loss in generality in supposing that $r_1 = 0$, $r_1 = -1$,

$$\begin{aligned}
Y^2 &= A_1(X)^2 + 4t_1(X+r_1) \\
&= A_2(X)^2 + 4t_2(X+r_1)(X+r_2) \\
&= A_3(X)^2 + 4t_3(X+r_2)(X+r_3) \\
&= A_4(X)^2 + 4t_4(X+r_3)(X+r_4) \\
&= A_5(X)^2 + 4t_5(X+r_4)(X+r_5) \\
&= A_6(X)^2 + 4t_6(X+r_5)^2
\end{aligned}$$

under the conditions, $(X+r_i) \mid (A_i(x) + A_{i+1}(X))$.

However, this does not generalize to higher genus since $\deg(Q_h(X))$ should be expected to be greater than 1, in fact the cases of $\deg(Q_h(X)) < g$ are degenerate in a sense and are special.

Not much is known about torsion on Jacobians of genus 3 and above. It would indeed be interesting to see whether the methods presented here would continue to hold. Would it be possible to find equations something like

$$\begin{aligned}
Y_g^2 &= A_1(X)^2 + 4t_1 X^{f_1(g)} (X-1)^{h_1(g)} \\
&= A_2(X)^2 + 4t_2 X^{f_2(g)} (X-1)^{h_2(g)} \\
&\vdots
\end{aligned}$$

where the f_i and h_i are each linearly independent? If so, it would suggest that we would be able to find regulators of order g^3 , equivalently torsion divisors of order g^3 . There is no evidence for either case so far.

9. MORE SLEEPERS

The results of the preceding sections raise the question of which hyperelliptic curves with periodic continued fractions satisfy the Schinzel condition. The main difference now is that many of these curves come to us as isolated examples and seem not to belong to a parametrized family as in the genus 1 case. It turns out that either they are trivially satisfied or it is impossible for them to satisfy the Schinzel condition since they possess a quasi-period which can not be eliminated. Since the results are not very encompassing we include only a few examples.

Example 2. Consider Leprévost's example of regulator 29,

$$Y^2 = 4X^6 - 4X^5 + X^4 - 8X^3 + 20X^2 - 16X + 4.$$

A quick glance at the partial quotients suggests that there is no obvious substitution $X \equiv a \pmod{b}$ which would make them integral. However, if we first dilate $X \mapsto \frac{1}{2}V$, $Y \mapsto \frac{1}{8}U$ we find that the unit is in $\frac{1}{2}\mathbb{Z}[V]$ for $V \equiv 1 \pmod{4}$.

Example 3. Elkies' example of torsion 39 does not belong to a parametrized family and since there is no transformation which can remove the quasi-period, Proposition 18 entails it can never satisfy the Schinzel condition.

A final remark is that we have been considering sleepers as being parametrized by x and therefore having regulators of order $\log D$. However, we have seen Flynn's hyperelliptic family of curves with regulators of order g^2 . If we were to find some value of X such that for this family the Schinzel condition was met for each g then we would have a sequence of discriminants parametrized by g and having regulators of order g^2 . Any such family would be a creeper. In fact, Shanks' original sequence of discriminants is nothing other than the selection $X = 2, t = -2$ in Flynn's family of curves (8.3).

Chapter IV. Elementary aspects of creepers

Ultimately, we intend to show that any creeper can be represented as

$$D_n = \frac{c^2}{d^2} \left[\left(qrx^n + \frac{mz^2x^k - ly^2}{q} \right)^2 + 4rly^2x^n \right]$$

with r, l, m squarefree and

$$rly^2mz^2 \mid d^2 D_n, \quad (qrx, lymz) = 1, \quad (yl, mz) = 1, \quad (qr, x) = 1, \quad q \mid mz^2x^k - ly^2$$

and moreover that any such representation is a creeper. Before doing this we are going to examine some simple cases in order to understand the features of creepers which are paramount to the investigation. In this section, we will not concern ourselves with any of the terms c, d, y or z . In other words, we are restricting our discussion to

$$D_n = (qrx^n + (mx^k - l)/q)^2 + 4lrx^n.$$

In determining the continued fraction expansion of ω some useful notation is: if $D \equiv t \pmod{4}$ (remembering that t must be either 0 or 1) then we will take

$$s_1 = qrx^n + (mx^k - l)/q \quad \text{and} \quad s_2 = qrx^n - (mx^k - l)/q$$

as well as

$$S_1 = (s_1 - t)/2 \quad \text{and} \quad S_2 = (s_2 - t)/2$$

so that

$$D_n = s_1^2 + 4lrx^n = s_2^2 + 4rmx^{n+k}.$$

Further abbreviations are:

$$\alpha = \omega + S_1, \quad \bar{\alpha} = \bar{\omega} + S_1 \quad \text{and} \quad \beta = \omega + S_2, \quad \bar{\beta} = \bar{\omega} + S_2$$

This notation will be maintained throughout. For later use, note that

$$\begin{aligned} \alpha\bar{\alpha} &= (\omega + S_1)(\bar{\omega} + S_1) = \omega\bar{\omega} + S_1(\omega + \bar{\omega}) + S_1^2 \\ &= \frac{(t^2 - D)}{4} + t\frac{(s_1 - t)}{2} + \frac{(s_1^2 - 2s_1t + t^2)}{4} \\ &= \frac{s_1^2 - D}{4} = -lrx^n \end{aligned}$$

Similarly,

$$\begin{aligned} \beta\bar{\beta} &= (\omega + S_2)(\bar{\omega} + S_2) = \frac{(t^2 - D)}{4} + \frac{t(s_2 - t)}{2} + \frac{(s_2^2 - 2s_2t + t^2)}{4} \\ &= \frac{s_2^2 - D}{4} = -rmx^{n+k} \end{aligned}$$

Notice that we may always suppose that $(n, k) = 1$; for if not, that is, if $(n, k) = j$ then we can attain $(n, k) = 1$ via the transformation $x \rightarrow x^j$ and $k \rightarrow k/j$.

10. EASY KREEPERS

Our first case under consideration is to take $m = 1, l > 0$ with l and r squarefree. This is the simplest extension of the expressions given by earlier investigations.

Remembering that the trace of ω is t , the continued fraction expansion of ω commences as

$$\begin{aligned} \omega &= S_1 + t - (\bar{\omega} + S_1) \\ \frac{\omega + S_1}{lrx^n} &= \frac{q}{l} - \frac{\bar{\beta}}{lrx^n}. \end{aligned}$$

In the earlier cases of creepers, q/l was an integer and the expansion proceeded happily enough. However, now in general q/l is not an integer. Fortunately, Lemma 2 tells us exactly how to deal with fractional partial quotients.

10.1. Handling fractional partial quotients

Applying Lemma 2 (see page 8) to $q/l - \bar{\beta}/lrx^n$ we find the next partial quotients consist of the expansion of q/l followed by the complete quotient

$$\theta_1 = \frac{lr x^n}{-\bar{\beta}l^2(-1)^{p_1+1}} - \frac{c_1}{l},$$

where $p_1 := l(q/l)$ and $c_1 \equiv (-1)^{p_1}/q \pmod{l}$. This is,

$$\theta_1 = \frac{rx^n(\omega + S_2)}{-\bar{\beta}\bar{\beta}l(-1)^{p_1+1}} - \frac{c_1}{l} = \frac{\omega + S_2}{lx^k(-1)^{p_1+1}} - \frac{c_1}{l},$$

recalling that $l(a)$ represents the length of the continued fraction expansion of the rational a , and that we are free to choose its parity. Since we would like the above element to be reduced, we need $(-1)^{p_1+1} = 1$. Therefore, we choose the *odd* expansion of q/l . Then the next line in the expansion is

$$\frac{qrx^{n-k} - c_1 - e_1}{l} - \frac{(\bar{\alpha} - e_1x^k)}{lx^k},$$

where $e_1 \equiv qrx^{n-k} - c_1 \pmod{l}$. In fact, exactly in the case $l|D$ we have that $e_1 = 0$. For, $l|D$ if and only if $s_1 \equiv 0 \pmod{l}$, thus if and only if $q^2rx^n + mx^k \equiv 0 \pmod{l}$ (because $(q, l) = 1$), so if and only if $qrx^{n-k} - 1/q \equiv 0 \pmod{l}$.

Hence, under the condition $l|D$, the remainder is $-\bar{\alpha}/lx^k$ and the continued fraction expansion continues as:

$$\begin{aligned} \frac{lx^k}{-\bar{\alpha}} &= \frac{lx^k\alpha}{-\alpha\bar{\alpha}} = \frac{lx^k(\omega + S_1)}{lr x^n} \\ &= \frac{\omega + S_1}{rx^{n-k}} = qx^k - \frac{\bar{\beta}}{rx^{n-k}} \quad (\text{actual line}) \\ \frac{rx^{n-k}}{-\bar{\beta}} &= \frac{rx^{n-k}\beta}{-\beta\bar{\beta}} = \frac{rx^{n-k}(\omega + S_2)}{rx^{n+k}} \\ &= \frac{(\omega + S_2)}{x^{2k}} = qrx^{n-2k} - \frac{\bar{\alpha}}{x^{2k}} \quad (\text{actual line}) \\ \frac{x^{2k}}{-\bar{\alpha}} &= \frac{x^{2k}\alpha}{-\alpha\bar{\alpha}} = \frac{x^{2k}(\omega + S_1)}{lr x^n} \\ &= \frac{(\omega + S_1)}{lr x^{n-2k}} = \frac{qx^{2k}}{l} - \frac{\bar{\beta}}{lr x^{n-2k}} \quad (\text{actual line}). \end{aligned}$$

Seemingly, trouble has struck again in that qx^{2k}/l is not an integer. Armed with our previous knowledge we know that this fraction can be comfortably handled just like the previous one. Proceeding as before using Lemma 2,

$$\frac{qx^{2k}}{l} - \frac{\bar{\beta}}{lr x^{n-2k}} = \left[\vec{w}, \frac{lr x^{n-2k}}{-l^2\bar{\beta}} - \frac{c_2}{l} \right],$$

where $\vec{w} = [qx^{2k}/l]_{\text{odd}}$ and $c_2 \equiv -(qx^{2k})^{-1} \pmod{l}$. Thus,

$$\frac{lr x^{n-2k}}{-l^2\bar{\beta}} - \frac{c_2}{l} = \frac{rx^{n-2k}(\omega + S_2)}{lr x^{n+k}} - \frac{c_2}{l}.$$

And we find

$$\frac{(\omega + S_2)}{lx^{3k}} - \frac{c_2}{l} = \frac{rqx^{n-3k} - c_2}{l} - \frac{\bar{\alpha}}{lx^{3k}}.$$

Furthermore, we make the happy discovery that the apparent fraction $(rqx^{n-3k} - c_2)/l$ is in fact an integer because

$$\begin{aligned} rqx^{n-3k} - c_2 &\equiv rqx^{n-3k} + (qx^{2k})^{-1} \pmod{l} \\ &\equiv rqx^{n-k} + q^{-1} \pmod{l}, \end{aligned}$$

which we know from before to be congruent to 0 modulo l (under the condition $l|D$). Note that if we tried to continue with the expansion of $(\bar{\alpha} - e_1x^k)/lx^k$ when l does not divide D then l would propagate through the denominators in arbitrarily large powers. Intuitively, we see that we could not then explicitly construct a unit. This intuition will be made explicit later.

The pattern for the continued fraction expansion is now apparent. Ignoring the terms arising from the expansion of a rational, the Q_h 's cycle for each $j \geq 0$ as:

$$lx^{(1+2j)k}, \quad rx^{n-(1+2j)k}, \quad x^{(2+2j)k}, \quad rlx^{n-(2+2j)k}.$$

Ultimately, either $n - (\{1, 2\} + 2j)k = 0$ or $n - (\{1, 2\} + 2j)k < 0$. In the first case we have reached halfway (either $Q_h = r$ or $Q_h = rl$, so either way, $(\frac{D_n}{Q_h}) = 0$).

In the second case there is some detail still remaining. (As an aside, note that if $x \equiv 1 \pmod{l}$ then the rational numbers appearing are always the same. In such a case, we only need to physically determine up to $Q_h = rlx^{n-2k}$).

10.2. When $k|n$

As an example for $k|n$ it suffices to take $k = 1$ because, as noted earlier, we may assume that $(k, n) = 1$.

Example 4. Consider the creeper given by: $l = 11, x = 67, q = 7, r = 2$. Here,

$$D_n = \left(7 \cdot 2 \cdot 67^n + \frac{(67 - 11)}{7}\right)^2 + 4 \cdot 2 \cdot 11 \cdot 67^n.$$

In this example, we have $l|D_n$ for all n so we obtain a creeper for every n . Also, $x \equiv 1 \pmod{l}$ means that the sequence of constant partial quotients remain the same for all n .

$$\begin{aligned} \omega &= S_1 + 1 - (\bar{\omega} + S_1) \\ \frac{\omega + S_1}{2 \cdot 11 \cdot 67^n} &= \frac{7}{11} - \frac{(\bar{\omega} + S_2)}{2 \cdot 11 \cdot 67^n} \end{aligned}$$

Since $[7/11]_{\text{odd}} = [0, 1, 1, 1, 3]$ the next partial quotients are $\{0, 1, 1, 1, 3\}$ followed by

$$\frac{7 \cdot 2 \cdot 67^{n-1} - 3}{11} \quad \text{because } 3 \equiv -1/7 \pmod{11}.$$

Then we have $\{7 \cdot 67, 7 \cdot 2 \cdot 67^{n-2}\}$ followed by the element

$$\frac{\omega + S_1}{2 \cdot 11 \cdot 67^{n-2}} = \frac{7 \cdot 67^2}{11} - \frac{(\bar{\omega} + S_2)}{2 \cdot 11 \cdot 67^{n-2}}.$$

The next partial quotients are then $\{7(67^2 - 1)/11, 1, 1, 1, 3\}$. This calculation is easy since $67 \equiv 1 \pmod{11}$ implies

$$[7 \cdot 67^s/11]_{\text{odd}} = [7(67^s - 1)/11, 1, 1, 1, 3].$$

And then we are back to $7 \cdot 67^3$ and the pattern is clear.

Hence,

$$\begin{aligned} \omega &= [S_1 + 2, 1, 1, 3, \frac{7 \cdot 2 \cdot 67^{n-1} - 3}{11}, 7 \cdot 67, 7 \cdot 2 \cdot 67^{n-2}, \frac{7(67^2 - 1)}{11}, 1, 1, 1, \\ &\quad 3, \frac{7 \cdot 2 \cdot 67^{n-3} - 3}{11}, 7 \cdot 67^3, 7 \cdot 2 \cdot 67^{n-4}, \frac{7(67^4 - 1)}{11}, 1, 1, 1, 3, \dots]. \end{aligned}$$

If n is even, then ultimately, $j = n - 1$,

$$\left[\dots, \frac{7 \cdot 2 \cdot 67 - 3}{11}, 7 \cdot 67^j, 7 \cdot 2, \frac{7(67^n - 1)}{11}, 7 \cdot 2, 7 \cdot 67^j, \dots \right]$$

because when $a_h = 7(67^n - 1)/11$ we also have $Q_h = rl = 22|D_n$, which implies that this point is halfway in the continued fraction expansion. Furthermore,

$$lp(\omega) = (l(p_1) + 3)n - 2 \quad \text{and} \quad p_1 = [7/11]_{\text{odd}} = [0, 1, 1, 1, 3]$$

implies that the period length of ω is $8n - 2$. (The 3 arises from the number of terms where $x|Q_h$ in a sequence, excluding $Q_h = lx^{ik}$). When n is odd, we get $j = n$ and

$$[\dots, 1, 1, 1, 3, (7 \cdot 2 - 3)/11, 7 \cdot 67^n, (7 \cdot 2 - 3)/11, \dots]$$

because when $a_h = 7 \cdot 67^n$ we have $Q_h = r = 2|D_n$ which implies this is halfway in the expansion. Also,

$$lp(\omega) = (l(p_1) + 3)n + 2 = 8n + 2.$$

A full, detailed expansion can be located in the Tables on page 139.

10.3. When k does not divide n

If $k \nmid n$ then the above process continues until a fractional partial quotient (with a power of x in the denominator) occurs. This can happen in two different ways, depending upon the parity of $j = \lfloor n/k \rfloor$. Either we have the situation

$$\begin{aligned} \frac{\omega + S_1}{rx^{n-jk}} &= qx^{jk} - \frac{\bar{\beta}}{rx^{n-jk}} \\ \frac{\omega + S_2}{x^{(j+1)k}} &= \frac{rq}{x^{(j+1)k-n}} - \frac{\bar{\alpha}}{x^{(j+1)k}} \end{aligned} \quad (10.1a)$$

or

$$\begin{aligned} \frac{\omega + S_1}{rlx^{n-jk}} &= \frac{qx^{jk}}{l} - \frac{\bar{\beta}}{rlx^{n-jk}} \\ &\vdots \\ \frac{\omega + S_2}{lx^{(j+1)k}} - \frac{b_{j+1}}{l} &= \frac{qr - b_{j+1}x^{(j+1)k-n}}{lx^{(j+1)k-n}} - \frac{\bar{\alpha}}{lx^{(j+1)k}}, \end{aligned} \quad (10.1b)$$

where $n < (j+1)k < n+k$ implies $0 < (j+1)k - n < k$.

Of course, fractional partial quotients are the same thing as saying that the element $(\omega + S_2)/x^{(j+1)k}$ or $\omega + S_2/lx^{(j+1)k}$ is not reduced (that is $x^{(j+1)k} > x^n$).

A fractional partial quotient can be tamed via Lemma 2 so these situations shouldn't provide extra heartache. Here are the details for (10.1a).

From the line

$$\frac{rq}{x^{(j+1)k-n}} - \frac{\bar{\alpha}}{x^{(j+1)k}}$$

we take the *odd* expansion of $rq/x^{(j+1)k-n}$ followed by

$$\frac{x^{(j+1)k}}{-\bar{\alpha}x^{2((j+1)k-n)}} - \frac{c_{j+1}}{x^{(j+1)k-n}}, \quad c_{j+1} \equiv -1/rq \pmod{x^{(j+1)k-n}}.$$

This yields

$$\frac{\omega + S_1}{rlx^{(j+1)k-n}} - \frac{c_{j+1}}{x^{(j+1)k-n}} = \frac{\omega + S_1 - c_{j+1}rl}{rlx^{(j+1)k-n}}.$$

Normally we would be inclined to write the next line in the expansion as

$$a_i - \frac{\bar{\beta}}{rlx^{(j+1)k-n}}.$$

However, we know that this would lead us astray. Instead, observe that, since $(j+1)k - n < k$, a more appropriate action would be to take

$$\frac{\omega + S_1 - c_{j+1}rl}{rlx^{(j+1)k-n}} = \frac{s_1 - c_{j+1}rl}{rlx^{(j+1)k-n}} - \frac{\bar{\alpha}}{rlx^{(j+1)k-n}}.$$

Note that

$$\begin{aligned} s_1 - c_{j+1}rl &\equiv (q^2rx^n + (mx^k - l) + l)/q \pmod{lx^{(j+1)k-n}} \\ &\equiv x^k(q^2rx^{n-k} + m)/q \pmod{lx^{(j+1)k-n}} \\ &\equiv 0 \pmod{lx^{(j+1)k-n}}, \end{aligned}$$

because $(j+1)k - n < k$ and $l|q^2rx^{n-k} + m$. And $r|s_1 - c_{j+1}rl$ since $r|s_1$. Also note that it was essential that $(j+1)k - n < k$ in order to have the m in the second line. Without it, divisibility would not hold and trouble would arise. This is also good since such a method could not apply to an exponent of x less than k . Hence the expansion continues as (here $(j+1)k - n$ has been replaced by e),

$$\begin{aligned} \frac{\omega + S_1}{x^{n-e}} &= qrx^e - \frac{\bar{\beta}}{x^{n-e}} \\ \frac{\omega + S_2}{rx^{k+e}} &= qx^{n-(k+e)} - \frac{\bar{\alpha}}{rx^{k+e}} \\ \frac{\omega + S_1}{lx^{n-(k+e)}} &= \frac{qrx^{k+e}}{l} - \frac{\bar{\beta}}{lx^{n-(k+e)}}. \end{aligned}$$

And now we have again come to a fractional partial quotient.

This process will continue until an element with no power of x occurs in the denominator; such a term must exist because of the coprimality of n and k . At this point, $Q_h = r$ or $Q_h = rl$ and this must be halfway in the expansion.

Also note that previously the powers of x increased in the Q_h 's containing r and decreased otherwise, but when we reach the power of x which is larger than n we find that this reverses itself. Indeed, each time we find a Q_h which is too large, the pattern reverses.

The second situation (which arises when $\lfloor n/k \rfloor \equiv 0 \pmod{2}$) runs along exactly the same lines.

Example 5. As an illustration of this process, consider

$$D_n = (2 \cdot 5 \cdot 43^n + (43^3 - 7)/2)^2 + 4 \cdot 5 \cdot 7 \cdot 43^n$$

In this example we will need to consider both qx^i/l and qrx^i/l . Fortunately, $x \equiv 1 \pmod{l}$ means that we do not need to evaluate every time. Looking at $n = 11$,

$$\begin{aligned} \omega &= S_1 + 1 & - & (\bar{\omega} + S_1) \\ \frac{\omega + S_1}{5 \cdot 7 \cdot 43^{11}} &= \frac{2}{7} & - & \frac{\bar{\beta}}{5 \cdot 7 \cdot 43^{11}}. \end{aligned}$$

And $\lfloor 2/7 \rfloor_{\text{odd}} = [0, 3, 2]$, $-1/2 \equiv 3 \pmod{7}$ means

$$\begin{aligned} \frac{\omega + S_2}{7 \cdot 43^3} - \frac{3}{7} &= \frac{2 \cdot 5 \cdot 43^8 - 3}{7} & - & \frac{\bar{\alpha}}{7 \cdot 43^3} \\ \frac{\omega + S_1}{5 \cdot 43^8} &= 2 \cdot 43^3 & - & \frac{\bar{\beta}}{5 \cdot 43^8} \\ \frac{\omega + S_2}{43^6} &= 2 \cdot 5 \cdot 43^5 & - & \frac{\bar{\alpha}}{43^6} \\ \frac{\omega + S_1}{5 \cdot 7 \cdot 43^5} &= \frac{2 \cdot 43^6}{7} & - & \frac{\bar{\beta}}{5 \cdot 7 \cdot 43^5}. \end{aligned}$$

And $[\frac{2 \cdot 43^6}{7}]_{odd} = [\frac{2(43^6-1)}{7}, 3, 2]$, means

$$\begin{aligned} \frac{\omega + S_2}{7 \cdot 43^9} - \frac{3}{7} &= \frac{2 \cdot 5 \cdot 43^2 - 3}{7} - \frac{\bar{\alpha}}{7 \cdot 43^9} \\ \frac{\omega + S_1}{5 \cdot 43^2} &= 2 \cdot 43^9 - \frac{\bar{\beta}}{5 \cdot 43^2} \\ \frac{\omega + S_2}{43^{12}} &= \frac{2 \cdot 5}{43} - \frac{\bar{\alpha}}{43^{12}}. \end{aligned}$$

Here we obtain a 43 in the denominator and apply the earlier procedure,

So $[\frac{10}{43}]_{odd} = [0, 4, 3, 2, 1]$, $-1/10 \equiv 30 \pmod{43}$, implies

$$\begin{aligned} \frac{\omega + S_1}{5 \cdot 7 \cdot 43} - \frac{30}{43} &= \frac{s_1 - 30 \cdot 5 \cdot 2}{5 \cdot 7 \cdot 43} - \frac{\bar{\alpha}}{5 \cdot 7 \cdot 43} \\ \frac{\omega + S_1}{43^{10}} &= 2 \cdot 5 \cdot 43 - \frac{\bar{\beta}}{43^{10}} \\ \frac{\omega + S_2}{5 \cdot 43^4} &= 2 \cdot 43^7 - \frac{\bar{\alpha}}{5 \cdot 43^4} \\ \frac{\omega + S_1}{7 \cdot 43^7} &= \frac{2 \cdot 5 \cdot 43^4}{7} - \frac{\bar{\beta}}{7 \cdot 43^7}. \end{aligned}$$

Now we need to expand qrx^i/l , that is

$[\frac{2 \cdot 5 \cdot 43^4}{7}]_{odd} = [\frac{2 \cdot 5 \cdot 43^4 - 3}{7}, 2, 3]$, $-1/(10 \cdot 43^4) \equiv 2 \pmod{7}$ and

$$\begin{aligned} \frac{\omega + S_2}{5 \cdot 7 \cdot 43^7} - \frac{2}{7} &= \frac{2 \cdot 43^4 - 2}{7} - \frac{\bar{\alpha}}{5 \cdot 7 \cdot 43^7} \\ \frac{\omega + S_1}{43^4} &= 2 \cdot 5 \cdot 43^7 - \frac{\bar{\beta}}{5 \cdot 7 \cdot 43^7} \\ \frac{\omega + S_2}{5 \cdot 43^{10}} &= 2 \cdot 43 - \frac{\bar{\alpha}}{5 \cdot 43^{10}} \\ \frac{\omega + S_1}{7 \cdot 43} &= \frac{2 \cdot 5 \cdot 43^{10}}{7} - \frac{\bar{\beta}}{7 \cdot 43} \\ &\vdots \end{aligned}$$

The full expansion of this example can be found in the Tables at page 140.

11. WHEN l IS NEGATIVE

In the preceding section the exclusion of $l < 0$ was not of paramount importance, merely aesthetic value. Allowing negative l 's, whilst keeping $m = 1$, does not drastically alter the previous method. The only changes are that now there are negative partial quotients which need to be dealt with.

There is a slight change in the partial quotients appearing, due to the change in sign of l . The best way of seeing this is through a simple example.

Example 6. Consider

$$D_n = \left(11 \cdot 7 \cdot 131^n + \frac{(131 - (-23))}{11} \right)^2 + 4 \cdot 7(-23)131^n.$$

We shall let $n = 6$. Also note that $x \not\equiv 1 \pmod{l}$ but rather $x^{11} \equiv 1 \pmod{l}$. Thus normally, when finding rational expansions we would only be interested in the exponent of x modulo 11. Since in our example n is less than 11, none of the rational expansions will be repeated.

Noting that $2 \equiv -11^{-1} \pmod{23}$, the continued fraction expansion is

$$\begin{aligned}
\omega_n &= S_1 + 1 && - \bar{\alpha} \\
&\vdots && \{ \text{partial quotients of } 11/-23 \} \\
&&& [11/-23]_{\text{odd}} = [-1, 1, 1, 10, 1] \\
\frac{\beta - 2 \cdot 131}{23 \cdot 131} &= \frac{11 \cdot 7 \cdot 131^{n-1} - 2}{23} && - \frac{\bar{\alpha}}{23 \cdot 131} \\
\frac{\alpha}{-7 \cdot 131^{n-1}} &= -11 \cdot 131 && - \frac{\bar{\beta}}{-7 \cdot 131^{n-1}} \\
\frac{\beta}{-131^2} &= -11 \cdot 7 \cdot 131^{n-2} && - \frac{\bar{\alpha}}{-131^2} \\
\frac{\alpha}{7 \cdot 23 \cdot 131^{n-2}} &= \frac{11 \cdot 131^2}{23} && - \frac{\bar{\beta}}{7 \cdot 23 \cdot 131^{n-2}} \\
&\vdots && \{ \text{partial quotients of } [11 \cdot 131^2/23] \} \\
&&& [11 \cdot 131^2/23]_{\text{odd}} = [8207, 2, 3, 2, 1] \\
\frac{\beta - 16 \cdot 131^3}{23 \cdot 131^3} &= \frac{11 \cdot 7 \cdot 131^{n-3} - 16}{23} && - \frac{\bar{\alpha}}{23 \cdot 131^3} \\
\frac{\alpha}{-7 \cdot 131^{n-3}} &= -11 \cdot 131^3 && - \frac{\bar{\beta}}{23 \cdot 131^{n-3}} \\
\frac{\beta}{-131^4} &= -11 \cdot 7 \cdot 131^{n-4} && - \frac{\bar{\alpha}}{-131^4} \\
\frac{\alpha}{7 \cdot 23 \cdot 131^{n-4}} &= \frac{11 \cdot 131^4}{23} && - \frac{\bar{\beta}}{7 \cdot 23 \cdot 131^{n-4}} \\
&\vdots && \{ \text{partial quotients of } 11 \cdot 131^4/23 \} \\
&&& [11 \cdot 131^4/23]_{\text{odd}} = [140847788, 3, 3, 1, 1] \\
\frac{\beta - 13 \cdot 131^5}{23 \cdot 131^5} &= \frac{11 \cdot 7 \cdot 131^{n-5} - 16}{23} && - \frac{\bar{\alpha}}{23 \cdot 131^5} \\
\frac{\alpha}{-7 \cdot 131^{n-5}} &= -11 \cdot 131^5 && - \frac{\bar{\beta}}{-7 \cdot 131^{n-5}} \\
\frac{\beta}{-131^6} &= -11 \cdot 7 \cdot 131^{n-6} && - \frac{\bar{\alpha}}{-131^6} \\
\frac{\alpha}{7 \cdot 23 \cdot 131^{n-6}} &= \frac{11 \cdot 131^6}{23} && - \frac{\bar{\beta}}{7 \cdot 23 \cdot 131^{n-6}} \\
&\vdots &&
\end{aligned}$$

The Tables contain the detailed expansion for $n = 6$ which is

$$\omega_6 = \left[S_1 + 1, -1, 1, 10, 1, \frac{11 \cdot 7 \cdot 131^5 - 2}{23}, -11 \cdot 131, -11 \cdot 7 \cdot 131^4, \right. \\
\left. \frac{11 \cdot 131^2 - 10}{23}, 2, 3, 2, 1, \frac{11 \cdot 7 \cdot 131^3 - 16}{23}, -11 \cdot 131^3, -11 \cdot 7 \cdot 131^2, \right. \\
\left. \frac{11 \cdot 131^4 - 7}{23}, 3, 3, 1, 1, \frac{11 \cdot 7 \cdot 131 - 13}{23}, -11 \cdot 131^5, -11 \cdot 7, \frac{11 \cdot 131^6 - 21}{23}, \dots \right]$$

where the last term represents halfway of the expansion. After removing the negative partial quotients in the manner discussed earlier, we get

$$\omega_6 = [194575656054823, 1, 10, 129157421874, 1, 1440, 22676493916, 1, \\
8206, 2, 3, 2, 1, 7526216, 1, 24729000, 1321396, 1, 140847787, 3, \\
3, 1, 1, 437, 1, 424374386160, 76, 1, 2417088895089, 1, 76, \dots]$$

In this isolated example, removing the negative partial quotients is easy. However, in the expansion of a general creeper if there were $O(n)$ negative partial

quotients then it is not immediately obvious that the period length is still in an arithmetic progression after removing them. Consequently, after reaching the line

$$\frac{\beta - 2 \cdot 131}{23 \cdot 131} = \frac{11 \cdot 7 \cdot 131^{n-1} - 2}{23} - \frac{\bar{\alpha}}{23 \cdot 131}$$

even though $(11 \cdot 7 \cdot 131^{n-1} - 2)/23$ is an integer, we treat it as if it were a rational. In other words, applying Lemma 2, after the expansion of $(11 \cdot 7 \cdot 131^{n-1} - 2)/23$, the next complete quotient would be

$$\frac{-23 \cdot 131}{\bar{\alpha}(-1)^{p_2+1}} \quad \text{where } p_2 := l((11 \cdot 7 \cdot 131^{n-1} - 2)/23).$$

Thus if we want to ensure that $Q_h > 0$ we merely take $p_2 \equiv 0 \pmod{2}$. After the partial quotients $\{(11 \cdot 7 \cdot 131^{n-1} - 25)/23, 1\}$ the expansion continues,

$$\begin{aligned} \frac{\alpha}{7 \cdot 131^{n-1}} - 1 &= 11 \cdot 131 - 1 - \frac{\bar{\beta}}{7 \cdot 131^{n-1}} \\ \frac{\beta}{131^2} &= 11 \cdot 7 \cdot 131^{n-2} - \frac{\bar{\alpha}}{131^2} \end{aligned}$$

Again we take the *even* expansion of $11 \cdot 7 \cdot 131^{n-2}$, that is $[11 \cdot 7 \cdot 131^{n-2} - 1, 1]$, and this is followed by

$$\frac{\alpha}{7 \cdot 23 \cdot 131^{n-2}} - 1 = \frac{11 \cdot 131^2 - 23}{23} - \frac{\bar{\beta}}{7 \cdot 23 \cdot 131^{n-2}}$$

and so on $\dots$. Clearly, one can avoid negative partial quotients by applying Lemma 2 appropriately. When we come to the completely general expansion this is the route we will follow, in order to avoid removing negative partial quotients. But in practice, either method is just as effective.

Some might now be concerned about r, q being negative and providing negative partial quotients which can't easily be resolved. In Chapter V it is shown that q and r are positive, but here it suffices to note that without loss of generality $q > 0$ since the transformations

$$q \mapsto -q, \quad r \mapsto -r, \quad m \mapsto -m, \quad l \mapsto -l$$

preserve D_n . Similarly, without loss of generality $r > 0$ since

$$r \mapsto -r, \quad l \mapsto -l, \quad m \mapsto -m$$

also preserve D_n .

Alternatively, realizing that $rlx^n < 0$ for $l < 0$ is equivalent to saying $S_1 + t > \omega$ suggests that we find a different representation for D_n . In the case of $q = 1$ this can be done by examining

$$D_n = (rx^n + mx^k + l)^2 - 4mlx^k = s_3^2 - 4mlx^k. \quad (11.1)$$

And as usual, we take $S_3 = (s_3 - t)/2$, so then

$$-\omega\bar{\omega} = S_3^2 + tS_3 - mlx^k.$$

If $ml < 0$ this representation provides a “reduced” solution. It suggests we start by considering

$$\begin{aligned} \omega &= S_3 + t - (\bar{\omega} + S_3) \\ \frac{\omega + S_3}{mlx^k} &= \frac{rx^{n-k} + m}{lm} - \frac{\bar{\alpha}}{mlx^k} \end{aligned}$$

where this can now be continued in a similar manner to the earlier situations.

This is also perfectly good for $q \neq 1$, it just means more fractional partial quotients. Specifically, (11.1) becomes

$$D_n = \left(qrx^n + \frac{mx^k + l}{q} \right)^2 - \frac{4ml}{q^2} x^k$$

which now has rational terms and is not much of an improvement. This representation is however significant when expanding $\sqrt{D_n}$ as a function of x .

12. WHEN BOTH l AND m ARE NOT UNITS

The extra difficulty involved in allowing $m \neq 1$ is very similar to that introduced by allowing l to be something other than 1. Having treated the cases $l < 0$ and $l > 0$ separately, we saw the distinction between the two and should feel more than comfortable in not distinguishing between the two for m .

At this point it would seem wise to introduce some notations which will be convenient. It is already obvious that the points where x divides Q_h are vital. Consequently, we write

$$h_i := \text{the } i\text{-th occurrence of } x \text{ as a divisor of some } Q_h \text{ (} h = h_i \text{)}.$$

This occurrence means in “our” expansion which may not correspond with the regular continued fraction expansion of ω (we’ll see an example soon). Furthermore, when this complete quotient occurs

$$\frac{\omega + P_{h_i}}{Q_{h_i}} = \frac{A_i}{B_i} - \frac{(\overline{\omega} + P_{h_{i+1}})}{Q_{h_i}},$$

we will want to apply Lemma 2 to the right side. Thus we define:

$$p_i := l(A_i/B_i) \quad \text{and} \quad \Delta_i = (A_i, B_i).$$

(Remembering that we are free to choose the parity of p_i). Next, we need to abbreviate the exponents of x in Q_{h_i} . Take,

$$\lambda_1 := k$$

$$\lambda_{i+2} := \begin{cases} \lambda_i + k - n\varepsilon_i & \text{if } i \equiv 1 \pmod{2} \\ 0 & \text{if } i \equiv 0 \pmod{2} \end{cases}$$

where

$$\varepsilon_i = \begin{cases} 1 & \lambda_i \geq n - k \\ 0 & \lambda_i \leq n - k \text{ or } 2|i. \end{cases}$$

This means that $\varepsilon_i = 1$ represents those points in the expansion where $Q_{h_{i+1}} > x^n$. This notation will be exhibited explicitly in the later examples.

Starting from

$$\omega = S_1 + t - (\overline{\omega} + S_1),$$

it is desirable to ensure that the next element is positive. This can be done by taking

$$(-1)^{p_0+1} = \text{sign}(l) \quad (p_0 = l(S_1 + t)).$$

After the appropriate expansion of $S_1 + t$, the following complete quotient is

$$\frac{\omega + S_1}{|l|rx^n} - \frac{c_0}{1}$$

where

$$c_0 = \begin{cases} 1 & \text{when } l < 0 \\ 0 & \text{when } l > 0 \end{cases}$$

In fact, it is clear that whenever

$$\gamma = z + \frac{a}{b} \quad (z \in \mathbb{Z}),$$

we get $\gamma = [z, \frac{b}{a}] = [z - 1, 1, -1 + \frac{b}{-a}]$. Anyway, the next line in the continued fraction expansion is

$$\frac{q}{|l|} - c_0 - \frac{\overline{\beta}}{|l|rx^n}.$$

The next partial quotients are the expansion of $(q - c_0|l|)/|l|$ followed by the complete quotient,

$$\frac{|l|rx^n}{-\bar{\beta}|l|^2(-1)^{p_1+1}} - \frac{c_1}{|l|} = \frac{(\omega + S_2)}{|lm|x^k} - \frac{c_1}{|l|}, \quad c_1 \equiv (-1)^{p_1}/q \pmod{|l|},$$

$$p_1 := l((q - c_0|l|)/|l|),$$

provided $(-1)^{p_1+1} = \text{sign}(m)$. In other words, the expansion of $(q - c_0|l|)/|l|$ needs to be odd or even according to m being positive or negative respectively. The next line in the expansion is

$$\frac{qrx^{n-k} - c_1|m|}{|lm|} - \frac{\bar{\alpha}}{|lm|x^k}.$$

Now, observe that

$$qrx^{n-k} - c_1|m| \equiv (q^2rx^{n-k} + m)/q \equiv 0 \pmod{l};$$

because $l|s_1$. In the notation of the previous page,

$$A_2 = qrx^{n-k} - c_1|m|, \quad B_2 = |lm|, \quad \Delta_2 = |l|.$$

Thus the next complete quotient (after the expansion of $(qrx^{n-k} - c_1|m|)/|lm|$) is

$$\frac{-|lm|x^k}{\bar{\alpha}m^2(-1)^{p_2+1}} - \frac{c_2}{|m|} = \frac{\omega + S_1}{r|m|x^{n-k}} - \frac{c_2}{|m|}, \quad c_2 \equiv \frac{(-1)^{p_2}l}{qrx^{n-k} - c_1|m|} \pmod{m}$$

$$p_2 := l\left(\frac{qrx^{n-k} - c_1|m|}{|lm|}\right),$$

provided $(-1)^{p_2+1} = \text{sign}(l)$. And the next line in the expansion is

$$\frac{qx^k - c_2}{|m|} - \frac{\bar{\beta}}{r|m|x^{n-k}}.$$

Also,

$$qx^k - c_2 \equiv \frac{1}{qrx^{n-k} - c_1|m|}(q^2rx^n - c_1|m|qx^k + l) \equiv s_1/j \equiv 0 \pmod{m}.$$

Even though $(qx^k - c_2)/|m|$ is an integer, we still apply Lemma 2 to ensure that the next complete quotient is positive. After the appropriate expansion of $(qx^k - c_2)/|m|$, the next complete quotient is

$$\frac{-r|m|x^{n-k}}{\bar{\beta}(-1)^{p_3+1}} - c_3 = \frac{\omega + S_2}{x^{2k}} - c_3$$

where $p_3 = 1$, $c_3 = 0$ if m is positive, and $p_3 = 2$, $c_3 = 1$ if m is negative. The next line is

$$qrx^{n-2k} - c_3 - \frac{\bar{\alpha}}{x^{2k}}.$$

Again, this is an integer, but we still apply Lemma 2 and find the complete quotient,

$$\frac{-x^{2k}}{\bar{\alpha}(-1)^{p_4+1}} - c_4 = \frac{\omega + S_1}{|l|rx^{n-2k}} - c_4 = \frac{qx^{2k}}{|l|} - c_4 - \frac{\bar{\beta}}{|l|rx^{n-2k}}$$

where $p_4 = 1$, $c_4 = 0$ if l is positive and $p_4 = 2$, $c_4 = 1$ if l is negative. At this point, the pattern of the expansion is evident.

Example 7. As an example, consider

$$D_n = \left(172 \cdot 14145^n + \frac{(13 \cdot 14145 - 17)}{172}\right)^2 + 4 \cdot 17 \cdot 14145^n$$

In this example, $x \equiv 1 \pmod{ml}$ entails all the constant partial quotients are the same throughout the entire expansion. Rather than placing its expansion in the tables, we give it in full so as to point out the notations h_i , λ_i which are essential

when dealing with the general expansion, as in Chapters V and VII. (Actually, in this example the λ_i are particularly boring, they are more relevant when we have $k \nmid n$).

h	a_h	factors of Q_h	h_i	λ_i
0	688836516802358507512594285			
1	10	$17x^6$	h_0	
2	8			
3	2			
4	440707997998978586368	$(13)(17)x$	h_1	$\lambda_1 = k = 1$
5	3			
6	4			
7	187149	$13x^5$	h_2	
8	6885575649188707500	x^2	h_3	$\lambda_3 = 2k = 2$
9	2024349194	$17x^4$	h_4	
10	8			
11	2			
12	2202647642368	$(13)(17)x^3$	h_5	$\lambda_5 = 3k = 3$
13	3			
14	4			
15	37445009920269	$13x^3$	h_6	
16	34413936300	x^4	h_7	$\lambda_7 = 4k = 4$
17	405033861716982794	$17x^2$	h_8	
18	8			
19	2			
20	11008	$(13)(17)x^5$	h_9	$\lambda_9 = 5k = 5$
21	3			
22	4			
23	7492035965982635968269	$13x$	h_{10}	
24	172	x^6	h_{11}	$\lambda_{11} = 6k = 6$
25	81039590212042177354422857	17	h_{12}	
26	172	x^6	h_{13}	
27	7492035965982635968269	$13x$	h_{14}	
	$\vdots$			

The next example shows that the expansion which the preceding procedure creates might not correspond to the regular continued fraction expansion (in other words it may contain unreduced elements).

Example 8. Consider $D_n = (2^n - 5 \cdot 2^2 - 11)^2 + 4 \cdot 11 \cdot 2^n$. For D_n to be a kreeper, it is necessary that $n \equiv 6 \pmod{20}$. Here, we consider the case $n = 26$. The numerical expansion is given in the Tables on page 142 and upon checking, one will notice that it does not exactly correspond to the expansion detailed here. Applying the procedure laid out before,

$$\begin{aligned}
\frac{\omega + P_{h_0}}{Q_{h_0}} &= \frac{\omega + S_1}{11 \cdot 2^{26}} \\
\frac{\omega + P_{h_1}}{Q_{h_1}} &= \frac{\omega + S_2}{5 \cdot 11 \cdot 2^2} - \frac{c_1}{11} \\
\frac{\omega + P_{h_2}}{Q_{h_2}} &= \frac{\omega + S_1}{5 \cdot 2^{24}} - \frac{c_2}{5}.
\end{aligned}
\qquad \lambda_1 = k = 2$$

The complete quotient $\theta_{h_2} = (\omega + P_{h_2})/Q_{h_2}$ is unreduced because $5 \cdot 2^{24} > \omega - \bar{\omega}$.

$$\begin{aligned} \frac{\omega + P_{h_3}}{Q_{h_3}} &= \frac{\omega + S_2}{2^4} - c_3 & \lambda_3 = 2k = 4 \\ \frac{\omega + P_{h_4}}{Q_{h_4}} &= \frac{\omega + S_1}{11 \cdot 2^{22}} \\ &\vdots \\ \frac{\omega + P_{h_{19}}}{Q_{h_{19}}} &= \frac{\omega + S_2}{2^{20}} - c_{19} \\ \frac{\omega + P_{h_{20}}}{Q_{h_{20}}} &= \frac{\omega + S_1}{11 \cdot 2^6} \\ \frac{\omega + P_{h_{21}}}{Q_{h_{21}}} &= \frac{\omega + S_2}{5 \cdot 11 \cdot 2^{22}} - \frac{c_{21}}{11} & \lambda_{21} = 11k = 22. \end{aligned}$$

The complete quotient $\theta_{h_{21}} = (\omega + P_{h_{21}})/Q_{h_{21}}$ is unreduced because $5 \cdot 11 \cdot 2^{22} > \omega - \bar{\omega}$.

$$\begin{aligned} \frac{\omega + P_{h_{22}}}{Q_{h_{22}}} &= \frac{\omega + S_1}{5 \cdot 2^4} - \frac{c_{22}}{5} \\ \frac{\omega + P_{h_{23}}}{Q_{h_{23}}} &= \frac{\omega + S_2}{2^{24}} - c_{23} & \lambda_{23} = 12k = 24 \\ \frac{\omega + P_{h_{24}}}{Q_{h_{24}}} &= \frac{\omega + S_1}{11 \cdot 2^2} \\ \frac{\omega + P_{h_{25}}}{Q_{h_{25}}} &= \frac{\omega + S_2}{5 \cdot 11 \cdot 2^{26}} - \frac{c_{25}}{11} & \lambda_{25} = 13k = 26. \end{aligned}$$

The complete quotient $\theta_{h_{25}}$ is obviously unreduced because $Q_{h_{25}} > Q_{h_{21}}$. Furthermore,

$$\frac{\omega + P_{h_{26}}}{Q_{h_{26}}} = \frac{\omega + S_1}{5 \cdot 2^0} - \frac{c_{26}}{5}$$

and this is halfway through the expansion.

Specifically, the elements:

$$\frac{\omega + P_{h_0}}{Q_{h_0}}, \quad \frac{\omega + P_{h_2}}{Q_{h_2}}, \quad \frac{\omega + P_{h_{21}}}{Q_{h_{21}}}, \quad \frac{\omega + P_{h_{25}}}{Q_{h_{25}}},$$

fail to appear. This just means that for each of these complete quotients θ_{h_i} , we have $A_i/B_i < 1$. In other words, there is a nonpositive partial quotient. This is no problem because in Chapter I we detailed how to remove negative partial quotients. The vital point is that the number of such occurrences is bounded independent of n . Consequently the difference in period lengths between “our” expansion and the regular expansion is a number bounded independently of n . Specifically, such points can only occur when Q_{h_i} is roughly x^n . In other words, if λ_i is almost zero or n . Equivalently, $\varepsilon_j = 1$ is about to occur. These remarks will all be made precise when we come to expanding a general keeper in Chapter VII.

Chapter V. Determining keepers

We wish to determine all families of discriminants $\{D_n\}$ which satisfy

1. $d^2 D_n = A^2 x^{2n} + Bx^n + C^2$ with $A, B, C, d \in \mathbb{Z}$
2. there exists an infinite set I such that for every $n \in I$, in the continued fraction expansion of ω_n there exists a $Q_h = x^g$ with g fixed independently of n
3. $lp(\omega_n) = an + b$ with $a, b \in \mathbb{Q}$ for every $n \in I$.

such a family is called a *keeper*. We commence with some introductory results.

13. PRELIMINARY RESULTS

Later we will need to transfer results from one order to another. Given a discriminant D , let D' represent the fundamental part of D , $K = \mathbb{Q}(\sqrt{D})$, and $\omega' = (t + \sqrt{D'})/2$ where $D' \equiv t \pmod{4}$ (so $\mathbb{Z} + \omega'\mathbb{Z}$ is the maximal order of K). Some simple results are:

Proposition 21. *If $\mathcal{O}_1 = \mathbb{Z} + f\omega'\mathbb{Z}$, $\mathcal{O}_2 = \mathbb{Z} + g\omega'\mathbb{Z}$ and $f|g$ then $\mathcal{O}_2 \subseteq \mathcal{O}_1$.*

Proposition 22. *If $R(\mathcal{O})$ denotes the regulator of the order $\mathcal{O}$ and $\mathcal{O}_2 \subseteq \mathcal{O}_1$ where $\mathcal{O}_1$ and $\mathcal{O}_2$ are orders of K then $R(\mathcal{O}_2) \geq R(\mathcal{O}_1)$.*

Proof. If $\varepsilon = (x + \sqrt{Dg^2y})/2$ is the fundamental unit of $R(\mathcal{O}_2)$ then $\varepsilon = (x + \sqrt{Df^2\frac{g}{f}y})/2$ is a unit of $\mathcal{O}_1$. Hence $R(\mathcal{O}_1) \leq R(\mathcal{O}_2)$. ■

More specifically, we have

Theorem 23. *If $\mathcal{O}_1 = \mathbb{Z} + f\omega'\mathbb{Z}$, $\mathcal{O}_2 = \mathbb{Z} + g\omega'\mathbb{Z}$ are both orders of K where $f|g$, then $2gR(\mathcal{O}_1) > fR(\mathcal{O}_2)$.*

Proof. Let $\varepsilon = x + f\omega'y$ be the fundamental unit of $\mathcal{O}_1$,

$$\varepsilon = \frac{V + fU\sqrt{D'}}{2} \quad (V, U \in \mathbb{Z}).$$

Let η be the fundamental unit of the order $\mathcal{O}_2$. Then $\eta = \varepsilon^m$ for some $m \in \mathbb{N}$,

$$\eta = \frac{R + gS\sqrt{D'}}{2} \quad (R, S \in \mathbb{Z}).$$

If we define the Lucas functions $v_n := \varepsilon^n + \bar{\varepsilon}^n$, $u_n := (\varepsilon^n - \bar{\varepsilon}^n)/(\varepsilon - \bar{\varepsilon})$ then

$$\varepsilon^n = \frac{v_n + fUu_n\sqrt{D'}}{2}.$$

By classical results concerning Lucas functions (see for example [84]), we know that $k|u_s$ if $s = \Phi(k)$. Also, it is easy to show that $\Phi(k) < 2k$. Hence,

$$m \leq \Phi(g/f) < 2g/f \quad \text{which implies} \quad \frac{1}{m} > \frac{f}{2g}$$

and $\log \eta = m \log \varepsilon$. So, $R(\mathcal{O}_2) = mR(\mathcal{O}_1)$ and $R(\mathcal{O}_1) = R(\mathcal{O}_2)/m > fR(\mathcal{O}_2)/2g$. ■

Corollary 24. *If $\mathcal{O}_1$ and $\mathcal{O}_2$ are given as above, but f does not necessarily divide g , then*

$$R(\mathcal{O}_1) > \frac{(g, f)}{2g} R(\mathcal{O}_2).$$

Proof. Let $h = [g, f] = gf/(g, f)$. Put $\mathcal{O}_3 = \mathbb{Z} + h\omega'\mathbb{Z}$ so that $\mathcal{O}_3$ is contained in $\mathcal{O}_1$ and $\mathcal{O}_2$, thus $R(\mathcal{O}_3) \geq R(\mathcal{O}_2)$. And $R(\mathcal{O}_1) \geq fR(\mathcal{O}_3)/2h$ by the above Theorem. Hence,

$$R(\mathcal{O}_1) > \frac{f}{2h}R(\mathcal{O}_2) \text{ implies } R(\mathcal{O}_1) > \frac{(g, f)}{2g}R(\mathcal{O}_2).$$

■

13.1. Independent ideals in keepers

Let $\mathfrak{a}$ be any ideal of the order $\mathcal{O}$ which has conductor f in $\mathbb{Q}(\sqrt{D})$, and suppose $(N(\mathfrak{a}), f) = 1$. It follows that $\mathfrak{a} = (t)\mathfrak{r}\mathfrak{s}$ where $t \in \mathbb{Z}$ and any prime ideal divisor of $\mathfrak{r}$ lies over a prime p which ramifies and any prime ideal divisor of $\mathfrak{s}$ lies over a prime p which splits. Furthermore, $\mathfrak{r}$ and $\mathfrak{s}$ are primitive. We will denote by $s(\mathfrak{a})$ the ideal $\mathfrak{s}$. Note that if $t = 1$ (in which case $\mathfrak{a}$ is primitive) then $\mathfrak{a}^2 = (r)\mathfrak{s}^2$ where r is squarefree and $r \mid D$. Also, $N(s(\mathfrak{a})) = N(s(\overline{\mathfrak{a}}))$.

In [21], Halter-Koch attempts to extend the results of Yamamoto [86] by suggesting that if there exist $a_1, \dots, a_n \in \mathbb{N}$ and infinitely many discriminants $D_1, D_2, \dots$ such that

- (i) $a_1, \dots, a_n$ are norms of reduced principal ideals for each D_i
- (ii) $a_i = a'_i a''_i$ where
 - (a) all the prime factors of a'_i split
 - (b) a''_i is squarefree, divides D_j for each j , but not the conductor
- (iii) $a'_1, \dots, a'_n$ are multiplicatively independent

then $R(D_i) \gg (\log D_i)^{n+1}$. Unfortunately, this result is not true. For suppose that there existed 3 ideals with norms ab , ac and bc , which satisfy the above conditions, and write $(a) = \mathfrak{a}\overline{\mathfrak{a}}$, $(b) = \mathfrak{b}\overline{\mathfrak{b}}$, $(c) = \mathfrak{c}\overline{\mathfrak{c}}$. The norms are multiplicatively independent but the ideals lying over the norms are not. Halter-Koch forms the set of all products between the $\mathfrak{a}, \dots, \overline{\mathfrak{c}}$ which are reduced. But alas, not all of the products are principal and the ones which are principal are insufficient to provide the desired result. One way to overcome this is to use multiplicatively independent ideals instead of norms.

Definition. Let $\mathfrak{b}_1, \dots, \mathfrak{b}_n$ be integral invertible ideals of an order $\mathcal{O}$. We say that $\mathfrak{b}_1, \dots, \mathfrak{b}_n$ are *independent* in $\mathcal{O}$ if whenever there exist nonzero integers u, v and $\alpha_i, \beta_i \in \mathbb{Z}^+$ ($i = 1, \dots, n$) such that

$$(u) \prod_{i=1}^n \mathfrak{b}_i^{\alpha_i} = (v) \prod_{i=1}^n \mathfrak{b}_i^{\beta_i}$$

then $\alpha_i = \beta_i$ ($i = 1, \dots, n$).

A sufficient condition for the independence of two ideals $\mathfrak{b}$ and $\mathfrak{c}$ in $\mathcal{O}$ is given in Theorem 26, which needs the following,

Lemma 25. *If $\mathfrak{b}$ and $\mathfrak{c}$ are dependent in $\mathcal{O}$ then there exist nonzero integers u, v and nonnegative integers m, n , with $m + n > 0$, such that*

$$(u)\mathfrak{b}^m = (v)\mathfrak{d}^n$$

where $\mathfrak{d}$ is equal to $\mathfrak{c}$ or $\overline{\mathfrak{c}}$.

Proof. If $\mathfrak{b}, \mathfrak{c}$ are dependent there must exist integers u, v, i_1, i_2, j_1, j_2 such that

$$(u)\mathfrak{b}^{i_1}\mathfrak{c}^{i_2} = (v)\mathfrak{b}^{j_1}\mathfrak{c}^{j_2}$$

where i_1, i_2, j_1, j_2 are all positive and either $i_1 \neq j_1$ or $i_2 \neq j_2$. Without loss of generality, $i_1 > j_1$, whereupon

$$(u)\mathfrak{b}^{i_1-j_1}\mathfrak{c}^{i_2} = (v)\mathfrak{c}^{j_2}.$$

If $j_2 \geq i_2$ we get

$$(u)\mathfrak{b}^{i_1-j_1} = (v)\mathfrak{c}^{j_2-i_2}.$$

If $j_2 < i_2$ we get

$$\begin{aligned} (u)\mathfrak{b}^{i_1-j_1}\mathfrak{c}^{i_2-j_2} &= (v) \\ (u)\mathfrak{b}^{i_1-j_1}\mathfrak{c}^{i_2-j_2}(\bar{\mathfrak{c}})^{i_2-j_2} &= (v)(\bar{\mathfrak{c}})^{i_2-j_2} \\ ((u)N(\mathfrak{c})^{i_2-j_2})\mathfrak{b}^{i_1-j_1} &= (v)(\bar{\mathfrak{c}})^{i_2-j_2}. \end{aligned}$$

In each of these cases, u and v are nonzero integers; thus, in either case we obtain the desired result. $\blacksquare$

Theorem 26. *If $\mathfrak{b}$ and $\mathfrak{c}$ are dependent in $\mathcal{O}$ and $(N(\mathfrak{b})N(\mathfrak{c}), f) = 1$ then for some nonnegative integers m, n , with $m + n > 0$ we have*

$$N(s(\mathfrak{b}))^m = N(s(\mathfrak{c}))^n.$$

Proof. By the preceding Lemma, we know there must exist integers m, n, u, v such that

$$(u)\mathfrak{b}^m = (v)\mathfrak{d}^n,$$

where m, n are nonnegative and at least one of m, n is positive. Then we can write $(ut_1^m)\tilde{\mathfrak{b}}^m = (vt_2^n)\tilde{\mathfrak{d}}^n$ where $\tilde{\mathfrak{b}}$ and $\tilde{\mathfrak{d}}$ are primitive.

The condition $(f, N(\mathfrak{b})N(\mathfrak{c})) = 1$ and primitivity allow us to write

$$\begin{aligned} (ut_1^m)\tilde{\mathfrak{b}}^m &= (ut_1^m)r(\mathfrak{b})^m s(\mathfrak{b})^m, \\ (vt_2^n)\tilde{\mathfrak{d}}^n &= (vt_2^n)r(\mathfrak{d})^n s(\mathfrak{d})^n. \end{aligned}$$

Squaring these yields

$$(u^2 r_1^m t_1^{2m})s(\mathfrak{b})^{2m} = (v^2 r_2^n t_2^{2n})s(\mathfrak{d})^{2n}$$

where r_1 and r_2 divide D_n . Dividing out any common factors of $u^2 r_1^m t_1^{2m}$ and $v^2 r_2^n t_2^{2n}$ provides coprime integers u_1, v_1 such that

$$(u_1)(s(\mathfrak{b}))^{2m} = (v_1)(s(\mathfrak{d}))^{2n}.$$

Let $\mathfrak{p}^\alpha \parallel s(\mathfrak{b})$ and $\mathfrak{p}^\beta \parallel s(\mathfrak{d})$. If $\mathfrak{p}^{2m\alpha} \nmid s(\mathfrak{d})$ then $N(\mathfrak{p})|v_1$ implies that $\bar{\mathfrak{p}}|(v_1)$. The coprimality of u_1 and v_1 means $\bar{\mathfrak{p}}|(s(\mathfrak{b}))^{2m}$ which gives $N(\mathfrak{p})|s(\mathfrak{b})$ which is impossible because $s(\mathfrak{b})$ is primitive.

Hence, $\mathfrak{p}^{2m\alpha} \mid s(\mathfrak{d})$ which means that $2m\alpha \leq 2n\beta$. By symmetry, $2n\beta \leq 2m\alpha$ and so $m\alpha = n\beta$. Thus, $s(\mathfrak{b})^m = s(\mathfrak{d})^n$ and we find $N(s(\mathfrak{b}))^m = N(s(\mathfrak{d}))^n$. $\blacksquare$

13.2. Extending Yamamoto's result

Suppose that $\mathfrak{b}_1, \dots, \mathfrak{b}_n$ are independent ideals of $\mathcal{O}$ of discriminant D and let

$$S = \left\{ \prod_{i=1}^n \mathfrak{b}_i^{\alpha_i} : \alpha_i \geq 0 \ i = 1, \dots, n \text{ and } \prod_{i=1}^n N(\mathfrak{b}_i)^{\alpha_i} < \sqrt{D}/2 \right\}$$

The second condition being to ensure that each primitive ideal in S is reduced. We note that if $(v)\mathfrak{a}, (u)\mathfrak{c} \in S$ where $\mathfrak{a}$ and $\mathfrak{c}$ are primitive ideals of $\mathcal{O}$ then $\mathfrak{a} \neq \mathfrak{c}$ because of the definition of independence.

Also note that if $\mathfrak{a}_i = \langle Q_{i-1}, P_{i-1} + \omega \rangle$, then

$$\frac{P_{i-1} + \omega}{Q_{i-1}} > \frac{\sqrt{D}/2}{Q_{i-1}} = \frac{\sqrt{D}/2}{N(\mathfrak{a}_i)}.$$

Now suppose $(v)\mathfrak{a} \in S$ where $\mathfrak{a}$ is reduced. We have

$$v^2 N(\mathfrak{a}) = \prod_{i=1}^n N(\mathfrak{b}_i)^{\alpha_i}$$

and

$$\mathfrak{a} = \mathfrak{a}_i = Q_{i-1}\mathbb{Z} + (P_{i-1} + \omega)\mathbb{Z},$$

where from above

$$\frac{P_{i-1} + \omega}{Q_{i-1}} > \frac{v^2\sqrt{D}/2}{\prod_{i=1}^n N(\mathfrak{b}_i)^{\alpha_i}} \geq \frac{\sqrt{D}/2}{\prod_{i=1}^n N(\mathfrak{b}_i)^{\alpha_i}}.$$

Theorem 27. *Let $\mathcal{O}_1, \mathcal{O}_2, \dots, \mathcal{O}_k, \dots$ be a sequence of orders each of discriminant D_i where $D_i < D_{i+1}$. Suppose further that in each $\mathcal{O}_i$ there exists an independent set of principal ideals $\{\mathfrak{b}_{i,j} : (j = 1, \dots, n)\}$ such that $N(\mathfrak{b}_{i,j})$ is fixed for each value of i . Then*

$$R(D_i) \gg (\log D_i)^{n+1}.$$

Proof. The set

$$S = \left\{ \prod_{i=1}^n \mathfrak{b}_i^{\alpha_i} : \alpha_i \geq 0 \ i = 1, \dots, n, \ \prod_{i=1}^n N(\mathfrak{b}_i)^{\alpha_i} < \sqrt{D}/2 \right\}$$

contains only principal ideals. Let $\gamma_1, \dots, \gamma_s$ represent the quadratic irrationals associated with the primitive part of these ideals. Hence,

$$\varepsilon = \prod_{\gamma \in I} \gamma \geq \prod_{\gamma_i \in S} \gamma_i.$$

And,

$$\gamma_i = \frac{b_i + \omega_n}{N(\mathfrak{b}_i)^{\alpha_i}} > \frac{\sqrt{D}/2}{\prod N(\mathfrak{b}_i)^{\alpha_i}}.$$

Hence,

$$\varepsilon > \prod' \frac{\sqrt{D}/2}{\prod N(\mathfrak{b}_i)^{\alpha_i}} = \varepsilon_0.$$

One may now follow the same procedure as in Yamamoto [86] Theorem 3.1 or Halter-Koch [21] §3 to show that

$$R(D_i) \gg \left(\log \sqrt{D_i} \right)^{n+1}.$$

■

14. BASIC OBSERVATIONS ON KREEPERS

Our objective is to conclude that keepers must look, up to square factors, like

$$D_n = (qrx^n + (mz^2x^k - ly^2)/q)^2 + 4rly^2x^n. \quad (14.1)$$

As an aside, a slight issue arises if $2 \mid (Ax + C)$ and $4 \nmid (A^2, C^2, G)$. In this case one could consider $D_n/4$ and this should still be a keeper (provided D_n was a keeper to begin with). The problem now is that we are unable to express $D_n/4$ in the form above. As an illustration, consider

$$D_n = (30 \cdot 61^n + 10)^2 + 4 \cdot 5 \cdot 61^n$$

which corresponds to the above form with the selections:

$$r = 5, \quad m = l = 1, \quad q = 6, \quad k = 1.$$

The point is that $4 \mid D_n$ but 4 does not divide any of the terms q, r, l, m . In other words, it is a “fluke” square. While it is possible for other fluke squares to occur, that is $D_n \equiv 0 \pmod{9}$ might occur for n in some infinite subset of $\mathbb{N}$, it is only for 4 that it is possible to divide by the square and still be able to represent $D_n = A^2x^{2n} + Bx^n + C^2$ with $A, B, C \in \mathbb{Z}$. Dividing the above example by 4, one finds

$$D_n = (15 \cdot 61^n + 5)^2 + 5 \cdot 61^n$$

and it is clear that there is no way to represent this in the form (14.1), even though this is a keeper¹. This explains some curious 2's which float about later on. Anyway, we can always write our discriminants as

$$D_n = \frac{1}{d^2} [(Ax^n + C)^2 + 4Gx^n] \quad \text{where } G = (B - 2AC)/4, \quad (14.2)$$

where $A, B, C, G \in \mathbb{Z}$ (not to be mistaken with the earlier A, B, C). Recall, that a discriminant D_n of an order must satisfy $D_n \equiv 0, 1 \pmod{4}$. Note that we can also write

$$D_n = \frac{1}{d^2} [(Ax^n - C)^2 + 4Hx^n] \quad \text{where } H = (B + 2AC)/4.$$

Next, any common factors of C and x can be moved into the square divisors. By considering,

$$W := \max_{i \in \mathbb{N}}(x^i, C) \quad m := \min\{i \in \mathbb{N} : (x^i, C) = W\} \quad n := v + 2m$$

so that $(x, C/W) = 1$, we can write

$$\begin{aligned} D_n &= \frac{1}{d^2} \left[\left(AW^2 \frac{x^{2m}}{W^2} x^v + W \frac{C}{W} \right)^2 + 4GW^2 \frac{x^{2m}}{W^2} x^v \right] \\ &= \frac{W^2}{d^2} \left[\left(AW \frac{x^{2m}}{W^2} x^v + \frac{C}{W} \right)^2 + 4G \frac{x^{2m}}{W^2} x^v \right] \\ &= \frac{W^2}{d^2} \left[(\overline{A}x^v + \overline{C})^2 + 4\overline{G}x^v \right] \end{aligned}$$

where $(x, \overline{C}) = 1$ and $\overline{A}, \overline{C}, \overline{G} \in \mathbb{Z}$. Furthermore, any square factors of (A^2, C^2, G) can also be removed, so that without loss of generality we may suppose that

$$D_n = \left(\frac{c}{d} \right)^2 [(Ax^n + C)^2 + 4Gx^n]$$

with $A, C, G \in \mathbb{Z}$, $(x, C) = 1$ and (A^2, C^2, G) squarefree.

The next few results don't use any of the properties of keepers, we are merely interested in determining an explicit formula for $(Ax^n + C)^2 + 4Gx^n$. Consequently, we define

$$E_n := (Ax^n + C)^2 + 4Gx^n \quad (14.3)$$

where $(x, C) = 1$, and (A^2, C^2, G) is squarefree.

The first result is a representation of A, C, G .

Theorem 28. *Given E_n as in (14.3) and the conditions on A, C, G, x stated above, we have that*

$$E_n = (qrNx^n + P(M - L)/q)^2 + 4rLNPx^n \quad (14.4)$$

where $r, q, P, N \in \mathbb{Z}^+$, $M, L \in \mathbb{Z}$, and the following conditions are satisfied:

- 1) r is squarefree,
- 2) $(P, rqN) = 1$,
- 3) $rq \mid M - L$,
- 4) $(M, L) = 1$,
- 5) $(rq, ML) = 1$,
- 6) $(r, LNP) = 1$.

¹It can be written in the form (14.1) if we allow $l, m \in \mathbb{Q}$ but this would not be wise

Proof. The selections we make are:

$$\begin{aligned} r &:= (A, C, G), & N &:= \left(\frac{A}{r}, \frac{G}{r}\right), & q &:= \frac{A}{rN}, \\ P &:= \left(\frac{C}{r}, \frac{G}{Nr}\right), & L &:= \frac{G}{rNP}, & M &:= \frac{AC + G}{rNP}. \end{aligned}$$

These selections make (14.3) and (14.4) equivalent, so it only remains to show that the conditions indicated hold.

Clearly, r, q, P, N are positive and r is squarefree since (A^2, C^2, G) is squarefree by assumption. Next,

$$\left(\frac{A}{rN}, \frac{G}{rN}\right) = 1 \text{ implies that } (q, PL) = 1, \quad (14.5)$$

and

$$\left(\frac{A}{r}, \frac{C}{r}, \frac{G}{r}\right) = 1 \text{ implies } \left(Nq, \frac{C}{r}, \frac{G}{r}\right) = 1$$

which implies that $(Nq, P) = 1$ and $(N, C/r) = 1$. Next if $p|r$ and $p|P$ then $p^2|G$ would give $p^2|(A^2, C^2, G)$ which is not allowed. Hence, $(P, r) = 1$ so that we now have $(rqN, P) = 1$, yielding (2). Similarly, $(r, N)^2|G$ implies $(r, N) = 1$, otherwise (A^2, C^2, G) would not be squarefree. Next,

$$M - L = \frac{AC}{rNP} = \frac{Cq}{P} \text{ so } C = \frac{P(M - L)}{q} \text{ and thus } qC = P(M - L).$$

Also, $(qr, P) = 1$ and $r|C$ gives $qr|M - L$, which proves (3). Next,

$$(M, L) = \left(\frac{G}{rNP} + \frac{AC}{rNP}, \frac{G}{rNP}\right) = \left(\frac{G}{rNP}, \frac{AC}{rNP}\right) = \left(\frac{G}{rNP}, \frac{A}{N} \frac{C}{rP}\right).$$

From the definition of P we know $(C/rP, G/Nr) = 1$ which entails

$$(M, L) = (A/N, L) = (qr, L) = (r, L),$$

where we have used (14.5) to recall that $(q, L) = 1$.

Furthermore, $q|(M - L)$ entails $(q, M) = 1$. Returning to (M, L) , we have

$$(M, L) = \left(r, \frac{G}{rNP}\right).$$

But r is squarefree so $(r, G/r) = 1$ which implies that $(M, L) = 1$ and $(r, L) = 1$. This gives (4). Also, $(r, M) = 1$ since $qr|(M - L)$. Thus we have $(rq, ML) = 1$, giving (5) and $(r, LNP) = 1$, giving (6). $\blacksquare$

Our next objective is to determine the terms which divide E_n and those which are coprime to E_n .

Lemma 29. $(E_n, NP) = 1$

Proof. Let $p|(E_n, NP)$, then

$$E_n \equiv \left(qrNx^n + \frac{P(M - L)}{q}\right)^2 \pmod{NP}$$

which means

$$p|qrNx^n + \frac{P(M - L)}{q}.$$

Now, if $p|N$ then $p|P(M - L)/q$. Also,

$$\left(N, \frac{P(M - L)}{qr}\right) = \left(N, \frac{C}{r}\right) = 1 \text{ because } (N, P) = 1 \text{ by (2) in Theorem 28.}$$

From $(N, r) = 1$ we find

$$(N, P(M - L)/q) = 1,$$

and so it is impossible to have $p|N$.

That only leaves $p|P$ in which case $p|qrNx^n$ but from (2) in Theorem 28, $(P, qrN) = 1$ and we must have $p|x$. However, this would mean $p|x$ and $p|C$ which contradicts $(x, C) = 1$. Hence $p \nmid P$ and so $(E_n, NP) = 1$. ■

Theorem 30. *For the family of discriminants given by (14.4) with $n \in I$ (an infinite subset of $\mathbb{N}$) there exists an infinite set $I' \subseteq I$ such that for every $n \in I'$ we have $M = v'mz^2Z$, $L = vly^2Y$ with Z, Y, z, y positive and*

$$ml|E'_n, \quad vv'zy|F_n, \quad vv'|2, \quad (ZY, E'_n) = 1, \quad \left(\frac{F_n}{zy}, mlZY\right) = 1$$

and $E_n = F_n^2 E'_n$ where E'_n is squarefree. Here m, l, z, y, Z, Y are the same for all $n \in I'$.

Proof. Define $y_n^2 v_n := (F_n^2, L)$ where v_n is squarefree. Next, write

$$\tilde{F}_n = \frac{F_n}{y_n v_n} \quad \text{so that} \quad \left(v_n \tilde{F}_n^2, \frac{L}{v_n y_n^2}\right) = 1.$$

And,

$$E_n = y_n^2 v_n^2 \tilde{F}_n^2 E'_n = S_n^2 + 4rv_n y_n^2 \frac{L}{v_n y_n^2} NPx^n \quad (14.6)$$

where

$$S_n = qrNx^n + \frac{P(M - L)}{q}.$$

From (14.6) we have $y_n v_n | S_n$ which in turn implies that $v_n^2 | 4rv_n \frac{L}{v_n y_n^2} NPx^n$. Now

$$\left(\frac{L}{v_n y_n^2}, v_n\right) = 1.$$

Also, $v_n | E_n$ and $(NP, E_n) = 1$ implies $(v_n, NP) = 1$. And $(r, L) = 1$ entails $(r, v_n) = 1$; hence

$$\left(v_n, \frac{L}{v_n y_n^2} rNP\right) = 1.$$

It follows that we must have $v_n | 4x^n$.

However, if v_n divides both x and S_n then $v_n | P(M - L)/q$ and so $v_n | (x, C)$ which is impossible because $(x, C) = 1$. Since v_n is squarefree, we have that $v_n | 2$.

Therefore, (14.6) becomes

$$\left(\frac{F_n}{y_n}\right)^2 E'_n = \left(\frac{S_n}{y_n}\right)^2 + 4r \frac{L}{y_n^2} NPx^n \quad (14.7)$$

or if $v_n = 2$

$$\left(\frac{F_n}{2y_n}\right)^2 E'_n = \left(\frac{S_n}{2y_n}\right)^2 + 2r \frac{L}{2y_n^2} NPx^n \quad (14.8)$$

We now investigate the factors of $L/v_n y_n^2$. Since

$$\left(\frac{F_n}{v_n y_n}, \frac{L}{v_n y_n^2}\right) = 1$$

it follows that for any prime factor, p , of $L/v_n y_n^2$, (14.7) or (14.8) gives

$$p \mid \frac{S_n}{v_n y_n^2} \quad \text{if and only if} \quad p \mid E'_n.$$

Consequently, we define l_n to be the product of all prime powers, p^α , of $L/v_n y_n^2$ which satisfy

$$p^\alpha \parallel \frac{L}{v_n y_n^2} \quad \text{and} \quad p \mid E'_n,$$

and Y_n to be the product of all prime powers, p^β , which satisfy

$$p^\beta \parallel \frac{L}{v_n y_n^2} \quad \text{and} \quad (E'_n, p) = 1.$$

Clearly, since $(F_n, L/v_n y_n^2) = 1$ we have $L/v_n y_n^2 = l_n Y_n$. We note that, l_n is squarefree, since for any square factor u^2 of l_n we have

$$u \mid E'_n \implies u \mid S_n/y_n \implies u^2 \mid (S_n/y_n)^2 \implies u^2 \mid (F_n/y_n)E'_n \implies u^2 \mid E'_n,$$

which is impossible because E'_n is squarefree. Thus l_n is squarefree and we find

$$L = v_n l_n y_n^2 Y_n, \quad (Y_n, E'_n) = 1, \quad v_n y_n \mid F_n, \quad l_n \mid E'_n, \quad v_n \mid 2.$$

Also, we can write equation (14.6) as

$$E_n = z_n^2 (v'_n)^2 \tilde{F}_n^2 E'_n = (S'_n)^2 + 4r v'_n z_n^2 \frac{M}{v'_n z_n^2} N P x^n$$

where

$$S'_n = q r N x^n - P(M - L)/q \quad \text{and} \quad z_n^2 v'_n = (F_n^2, M).$$

By similar reasoning we get

$$M = m_n v'_n z_n^2 Z_n, \quad (Z_n, E'_n) = 1, \quad v'_n z_n \mid F_n, \quad m_n \mid E'_n, \quad v'_n \mid 2.$$

and $4 \nmid v_n v'_n$ because $(M, L) = 1$.

As there are only a finite number of choices for $m_n, l_n, z_n, y_n, v_n, v'_n, Z_n, Y_n$ for fixed values of L and M there must exist some infinite set $I' \subseteq I$ for which

$$M = v' m z^2 Z, \quad L = v l y^2 Y, \quad m l \mid E'_n, \quad z y v v' \mid F_n, \quad (ZY, E'_n) = 1, \quad v v' \mid 2.$$

Finally,

$$\left(\frac{F_n}{v y}, \frac{L}{v y^2} \right) = 1, \quad \left(\frac{F_n}{v' z}, \frac{M}{v' z^2} \right) = 1 \quad \text{and} \quad (L, M) = 1,$$

entails,

$$\left(\frac{F_n}{v v' z y}, \frac{L M}{v v' z^2 y^2} \right) = 1 \quad \text{which gives} \quad \left(\frac{F_n}{v v' z y}, m l Z Y \right) = 1.$$

Moreover, $(v, l Y) = (v', m Z) = 1$ and $(L, M) = 1$ implies $(v, m Z) = (v', l Y) = 1$. Thus $(v v', m l Z Y) = 1$, which gives $(F_n / z y, m l Z Y) = 1$. $\blacksquare$

This completes our investigation of E_n ; we have determined the factors coprime to E_n and those which divide each part of it.

15. INDEPENDENT IDEALS IN KREEPERS

Completing the square on $4A^2 E_n$ gives,

$$\begin{aligned} 4A^2 E_n &= 4A^4 x^{2n} + 4A^2 B x^n + 4A^2 C^2 \\ &= (2A^2 x^n + B)^2 + 16 \frac{(2AC - B)}{4} \frac{(2AC + B)}{4} \\ A^2 E_n &= (A^2 x^n + B/2)^2 - 4GH \end{aligned}$$

and $B/2$ is an integer because $4 \mid (B - 2AC)$. In terms of the constants found in the previous section, this equation becomes

$$\begin{aligned} q^2 r^2 N^2 F_n^2 E'_n &= ((qrN)^2 x^n + rNP(L + M))^2 - 4(rLNP)(rMNP), \\ q^2 F_n^2 E'_n &= (q^2 r N x^n + P(L + M))^2 - 4P^2 LM, \end{aligned}$$

where from the previous results we may suppose that

$$LM = vv'lmz^2y^2ZY, \quad vv'zy|F_n, \quad vv'|2, \quad lm|E'_n, \\ q^2 \left(\frac{F_n}{zy} \right)^2 E'_n = \left(\frac{q^2 r N x^n + P(L+M)}{zy} \right)^2 - 4vv'P^2lmZY. \quad (15.1)$$

We first define, $\omega'_n := (\sigma'_n - 1 + \sqrt{E'_n})/\sigma'_n$ where E'_n is squarefree and $\sigma'_n = 2$ if $E'_n \equiv 1 \pmod{4}$ and $\sigma'_n = 1$ otherwise. Then

$$d^2 D_n = c^2 F_n^2 E'_n$$

and we further define

$$\mathcal{O}_n = \mathbb{Z} + \omega_n \mathbb{Z}$$

where

$$\omega_n := \frac{t_n + \sqrt{D_n}}{2} \quad \text{and} \quad t_n = \begin{cases} 0 & \text{if } D_n \equiv 0 \pmod{4} \\ 1 & \text{if } D_n \equiv 1 \pmod{4} \end{cases}$$

Proposition 31. $\mathcal{O}_n = \mathbb{Z} + \frac{cF_n\sigma'_n}{2d}\omega'_n\mathbb{Z}$

Proof. Suppose $4|D_n$, then $2|cF_n/d$ and

$$\sqrt{D_n} = \frac{cF_n}{d}\sqrt{E'_n} = \frac{cF_n}{d}(\sigma'_n\omega'_n - \sigma'_n + 1)$$

Therefore,

$$\begin{aligned} \mathcal{O}_n &= \mathbb{Z} + \frac{t_n + cF_n(\sigma'_n\omega'_n - \sigma'_n + 1)/d}{2}\mathbb{Z} \\ &= \mathbb{Z} + \left(t_n + \frac{cF_n}{2d}(1 - \sigma'_n) \right) \mathbb{Z} + \frac{cF_n}{2d}\sigma'_n\omega'_n\mathbb{Z} \end{aligned}$$

Now suppose that $4 \nmid D_n$. Then cF_n/d is odd and $D_n \equiv E'_n \equiv 1 \pmod{4}$, so $\sigma'_n = 2$. In either case, $\mathcal{O}_n = \mathbb{Z} + cF_n\sigma'_n\omega'_n/(2d)\mathbb{Z}$. $\blacksquare$

Our objective now is to find an ideal arising from (15.1) which has a norm coprime to the conductor of some order. This will require a treatment of several separate cases.

15.1. Finding an ideal prime to the conductor

To begin with, we write (15.1) as

$$q^2 \left(\frac{F_n}{zy} \right)^2 E'_n = s_n^2 - 4vv'P^2lmZY.$$

Now suppose $2|q$, then $(q, LM) = 1$ implies $2 \nmid vv'lmzyYZ$ and so $vv' = 1$. Put

$$\alpha_n := s_n/2 + \frac{q}{2} \frac{F_n}{zy} \sqrt{E'_n} \quad \text{then} \quad \alpha_n \in \mathbb{Z} + \frac{F_n}{zy}\omega'_n\mathbb{Z}.$$

Further, $N(\alpha_n) = P^2lmZY$ and $(F_n/zy, P^2lmZY) = 1$. Consequently, from now on we suppose that $2 \nmid q$.

Case I: $2 \nmid F_n/zy$.

In this case, $vv' = 1$ and we put $\beta_n = s_n + qF_n/zy\sqrt{E'_n}$ so that $N(\beta_n) = 4P^2lmZY$. Next,

$$\begin{aligned} q^2(F_n/zy)^2 E'_n &\equiv s_n^2 \pmod{4} \\ E'_n &\equiv s_n^2 \pmod{4} \\ &\equiv 1 \pmod{4}. \end{aligned}$$

Hence $\sigma'_n = 2$ and $\sqrt{E'_n} = 2\omega'_n - 1$. Then

$$\beta_n = s_n + \frac{qF_n}{zy}(2\omega'_n - 1) = s_n - \frac{qF_n}{zy} + \frac{2qF_n}{zy}\omega'_n.$$

Put $\alpha_n := \beta_n/2$, so then $N(\alpha_n) = P^2lmZY$ and $\alpha_n \in \mathbb{Z} + \frac{F_n}{zy}\omega'_n\mathbb{Z}$. Also note that $(N(\alpha_n), F_n/zy) = 1$.

Case II: $2 \mid F_n/zy$. In this case, we must have $2 \nmid qP$.

subcase (a): $2 \mid F_n/2zy$. We have,

$$q^2 \left(\frac{F_n}{2zy} \right)^2 E'_n = (s_n/2)^2 - vv'P^2lmZY$$

then

$$\begin{aligned} vv'P^2lmZY &\equiv (s_n/2)^2 \pmod{4} \\ &\equiv \{0, 1\} \pmod{4}. \end{aligned}$$

If $vv' = 2$ then $2 \mid P^2lmZY$ which is impossible because vv' and P^2lmZY are coprime. Hence $vv' = 1$ and we take

$$\alpha_n = s_n/2 + \frac{qF_n}{2zy}\sqrt{E'_n},$$

so

$$N(\alpha_n) = P^2lmZY \quad \text{and} \quad \alpha_n \in \mathbb{Z} + \frac{F_n}{2zy}\omega'_n\mathbb{Z}.$$

subcase (b): $2 \nmid F_n/2zy$. Then $E'_n \equiv (s_n/2)^2 - vv'P^2lmZY \pmod{4}$. Take,

$$\alpha_n = s_n/2 + \frac{qF_n}{2zy}\sqrt{E'_n}.$$

Suppose that $\sigma'_n = 2$, then $E'_n \equiv 1 \pmod{4}$. If $2 \nmid s_n/2$ then $4 \mid vv'P^2lmZY$ which means that $vv' = 1$. Then from $(F_n/zy, lmZY) = 1$ we have that $2 \nmid lmZY$. Also, $2 \nmid P$ since $(E_n, P) = 1$. Hence, $2 \nmid (s_n/2)$ is impossible. Thus, $2 \mid s_n/2$ and $vv' = 1$, so

$$N(\alpha_n) = P^2lmZY \quad \text{and} \quad \alpha_n \in \mathbb{Z} + \frac{F_n}{zy}\omega'_n\mathbb{Z}.$$

Now suppose that $\sigma'_n = 1$. Take,

$$\alpha_n = s_n/2 + \frac{qF_n}{2zy}\sqrt{E'_n} \in \mathbb{Z} + \frac{F_n}{2zy}\omega'_n\mathbb{Z},$$

then

$$N(\alpha_n) = vv'lmZYP^2 \quad \text{and} \quad (F_n/2zy, vv') = 1 \quad \text{because} \quad 2 \nmid F_n/2zy.$$

In conclusion, we have established that in all cases:

$$\begin{aligned} \alpha_n &\in \mathbb{Z} + \frac{F_n}{\lambda zy}\omega'_n\mathbb{Z} \quad \text{with } \lambda = 1 \text{ or } 2, \\ \left(N(\alpha_n), \frac{F_n}{\lambda zy} \right) &= 1, \\ N(\alpha_n) &= \begin{cases} vv'P^2lmZY & \text{when } 2 \nmid F_n/2zy, 2 \nmid q, \sigma'_n = 1. \\ P^2lmZY & \text{otherwise.} \end{cases} \end{aligned}$$

In each case we find that $\alpha_n \in \overline{\mathcal{O}}_n := \mathbb{Z} + f_1\omega'_n\mathbb{Z}$ with $N(\alpha_n, f_1) = 1$ and $f_1 = F_n/(\lambda zy)$.

15.2. Bounded norms in keepers

By the definition of a keeper, the continued fraction expansion of ω_n has some $Q_i = x^g$ with g fixed independently of n . In other words, there exists some $\mu_n \in \mathcal{O}_n$ such that $N(\mu_n) = x^g$. Also recall Proposition 31 which states that

$$\mathcal{O}_n = \mathbb{Z} + f_2 \omega'_n \mathbb{Z} \quad \text{where } f_2 = \frac{cF_n \sigma'_n}{2d}.$$

Let $f_3 = (f_1, f_2)$, so that α_n, μ_n are both contained in $\mathcal{O}_n^* := \mathbb{Z} + f_3 \omega'_n \mathbb{Z}$. We know that $(x, C) = 1$, so then $(x, F_n/zy) = 1$. Hence,

$$(N(\alpha_n), f_3) = (N(\mu_n), f_3) = 1.$$

Thus in the order $\mathcal{O}_n^*$ we have the two principal ideals $\mathfrak{a}_n = (\alpha_n)$ and $\mathfrak{b}_n = (\mu_n)$ (whose norms are fixed for infinitely many $n \in I$). Furthermore, both ideals have norms which are coprime to the conductor. Hence, we may apply Theorems 26 and 27 from pages 61 and 62.

Proposition 32. *If $\mathfrak{a}_n$ and $\mathfrak{b}_n$ are independent ideals in $\mathcal{O}_n^*$ then*

$$R(\mathcal{O}_n) \gg (\log D_n)^3$$

Proof. By Theorem 27,

$$R(\mathcal{O}_n^*) \gg (\log \Delta(\mathcal{O}_n^*))^3 = (\log(f_3^2 \Delta(\mathcal{O}'_n)))^3$$

where $\mathcal{O}'_n := \mathbb{Z} + \omega'_n \mathbb{Z}$ and $\Delta(\mathcal{O})$ denotes the discriminant of the order $\mathcal{O}$. By Theorem 23,

$$R(\mathcal{O}_n) > \frac{(f_3, f_2)}{2f_3} R(\mathcal{O}_n^*) \text{ implies } R(\mathcal{O}_n) > \frac{1}{2} R(\mathcal{O}_n^*) \gg (\log(f_3^2 \Delta(\mathcal{O}'_n)))^3. \quad (15.2)$$

Now,

$$f_3 = (f_1, f_2) = \left(\frac{F_n}{\lambda zy}, \frac{cF_n \sigma'_n}{2d} \right).$$

Therefore,

$$2\lambda zy d f_3 = (2dF_n, cF_n \sigma'_n zy \lambda) = F_n(2d, c\sigma'_n zy \lambda)$$

so

$$f_3 = F_n \frac{(2d, c\sigma'_n zy \lambda)}{2\lambda zy d}.$$

This allows us to say,

$$(\log f_3^2 \Delta(\mathcal{O}'_n))^3 \gg (\log F_n^2 \Delta(\mathcal{O}'_n))^3 \gg (\log F_n^2 E'_n)^3 = (\log d^2 D_n / c^2)^3 \gg (\log D_n)^3$$

Hence, from (15.2), we have that $R(\mathcal{O}_n) \gg (\log D_n)^3$. ■

Since the conclusion of the Proposition contradicts the conditions on keepers, in order for the sequence of discriminants $\{D_n\}$ to be a keeper, $\mathfrak{a}_n$ and $\mathfrak{b}_n$ must be dependent ideals in $\mathcal{O}_n^*$. And,

$$\left(\frac{F_n}{\lambda zy}, N(\mathfrak{a}_n)N(\mathfrak{b}_n) \right) = 1,$$

so Theorem 26 says that there must exist nonnegative integers e and f , with e and f not both 0 such that

$$N(s(\mathfrak{a}_n))^e = N(s(\mathfrak{b}_n))^f. \quad (15.3)$$

15.3. *Splitting factors*

We now determine the splitting factors of $N(\mathfrak{a}_n)$. We know that

$$q^2 \left(\frac{F_n}{zy} \right)^2 E'_n \equiv s_n^2 \pmod{4P^2 vv' lmZY}$$

We also know that if $p|lm$ then $p|E'_n$ in which case p can not split. Now consider p an odd prime divisor of $vv'P^2ZY$, in which case $p|PZY$. Since $(PZY, qE_n) = 1$ we have that $p \nmid s_n$ and therefore,

$$\left(\frac{E'_n}{p} \right) = 1.$$

Now suppose that $p = 2$ so that $2 \nmid q$. Since $(F_n/zy, lmZYP) = 1$ we have that $2 \nmid F_n/zy$. Then $N(\alpha_n) = lmZYP^2$ and

$$q^2 \left(\frac{F_n}{zy} \right)^2 E'_n \equiv s_n^2 \pmod{4lmZYP^2}.$$

That is $E'_n \equiv 1 \pmod{8}$; in other words $\left(\frac{E'_n}{2} \right) = 1$. In all cases we have that $N(s(\mathfrak{a}_n)) = P^2ZY$.

Now consider the splitting primes of $N(\mathfrak{b}_n) = x^g$. If $p|x$ then $p \nmid C$ and

$$E_n \equiv C^2 \pmod{4p}.$$

Also, $(E_n, x) = 1$, implies $\left(\frac{F_n^2}{p} \right) = 1$, hence

$$\left(\frac{E'_n}{p} \right) = 1 \quad \text{for both } p \text{ odd and even.}$$

Hence, $N(s(\mathfrak{b}_n)) = x^g$, and (15.3) gives

$$x^{gf} = (YZP^2)^e. \tag{15.4}$$

Finally we show that neither e nor f is zero.

If $e = 0$ then $x^{gf} = 1$ and since in this case we must have $f > 0$ we find $x = 1$ which is impossible. On the other hand, if $f = 0$ then Y, Z and P are all ± 1 , in which case

$$\frac{q^2 E_n}{(zy)^2} = s_n^2 \pm 4mlvv'$$

But $mlvv'|E_n/(zy)^2$ which by Theorem 15 implies that the period length of $q^2 E_n/(zy)^2$ is bounded for all n . From Raney's automata (or just note that we still have an exceptional function field whose unit satisfies the Schinzel condition) it is clear that any rational multiple of a sleeper is again a sleeper. Hence, $c^2 E_n/d^2 = D_n$ must also be a sleeper; in other words, D_n is not a kreeper. It follows that if D_n is to be a kreeper we must have (15.4) with e, f each being positive. Since $(x, P) = 1$ we must have $P = 1$. Taking $d := (gf, e)$, $k := gf/d$, $h := e/d$ gives,

$$(ZY)^h = x^k, \quad (h, k) = 1,$$

which implies that $x = R^h$, $ZY = R^k$ and since we have supposed that x is not a power, $h = 1$ and $ZY = x^k$. From Theorem 28, $(Z, Y) = 1$ hence

$$Z = U^k, \quad Y = T^k, \quad \text{where } (U, T) = 1.$$

The objective of the next 2 sections will be to show that $T = 1$ and $U = x$.

16. PART OF THE CONTINUED FRACTION EXPANSION OF ω_n 16.1. Simplifying the formula for D_n

There is no longer any need to distinguish between factors of E'_n and F_n , which means that we may attempt to absorb the terms v and v' into l and m respectively.

First, $(v, lY) = (v', mZ) = 1$ implies that vl and $v'm$ are squarefree. Next, $(qr, ML) = 1$ implies $(qr, vv') = 1$, and $vv'|E_n$ implies $(vv', NPx) = 1$. We also have, $(qrN, yzml) = 1$, hence

$$(qrNUT, yzmlvv') = 1.$$

Furthermore, $(L, M) = 1$ gives $(v, mzU) = 1$ and $(v', lyT) = 1$. Combining these with $(Tyl, mzU) = 1$ gives $(Tylv, v'mzU) = 1$. It is clear that $qr| mz^2v'U^k - vly^2T^k$ still. Lastly, $yv|F_n$ and $l|E'_n$ and

$$q^2E_n = q^2F_n^2E'_n = (q^2rNx^n + v'mz^2U^k - vly^2T^k)^2 + 4rq^2Nly^2vT^{k+n}U^n.$$

Hence,

$$\begin{aligned} & y^2vl|(q^2rNx^n + v'mz^2U^k - vly^2T^k)^2, \\ & yvl|q^2rNx^n + v'mz^2U^k. \end{aligned}$$

By symmetry, $mzv'|q^2rNx^n + vly^2T^k$. Thus all the required conditions are still met. Consequently, up to this point we have determined that for some infinite subset I of $\mathbb{N}$, a creeper must be of the form

$$D_n = \left(\frac{c}{d}\right)^2 \left[(qrN(UT)^n + (z^2mU^k - y^2lT^k)/q)^2 + 4rNly^2T^{k+n}U^n \right] \quad (16.1)$$

where m, l, r are squarefree, and

$$(qr, UT) = 1, \quad (qrNUT, yzml) = 1, \quad (Tyl, mzU) = 1, \quad qr|z^2mU^k - y^2lT^k,$$

and for every $n \in I$,

$$yl|q^2rNx^n + z^2mU^k \quad \text{and} \quad mz|q^2rNx^n + y^2lT^k, \quad (16.2)$$

and $N > 0, x = UT$. Let μ be the least positive difference of any two integers in I . Then $\nu, \nu + \mu \in I$ for some ν , and

$$yl|q^2rNx^\nu + z^2mU^k \quad \text{and} \quad yl|q^2rNx^{\nu+\mu} + z^2mU^k$$

which means $yl|q^2rNx^\nu(x^\mu - 1)$, so $yl|x^\mu - 1$ because $(yl, qrNx) = 1$. By symmetry, $zm|x^\mu - 1$. Thus, $ylmz|x^\mu - 1$. Hence the conditions (16.2) become

$$yl|q^2rN(UT)^\nu + z^2mU^k, \quad mz|q^2rN(UT)^\nu + y^2lT^k, \quad (TU)^\mu \equiv 1 \pmod{ylmz},$$

for any $n \in I$, such that $n \equiv \nu \pmod{\mu}$. Since the signs of m and l are not specified there is no loss of generality in supposing that q, U, T are all positive.

Since (16.1) can also be represented as

$$D_n = \left(\frac{c}{d}\right)^2 \left[(qrN(UT)^n - (z^2mU^k - y^2lT^k)/q)^2 + 4rNmz^2T^nU^{k+n} \right] \quad (16.3)$$

we may also assume without loss of generality that $U > T$. Furthermore, it is clear that there is no loss in generality in supposing that $(c, d) = 1$.

16.2. Some notation

We now rewrite equations (16.1) and (16.3) as

$$\begin{aligned} D_n &= \left(\frac{s_1}{d}\right)^2 + 4\left(\frac{c}{d}\right)^2 rNly^2T^{k+n}U^n \\ &= \left(\frac{s_2}{d}\right)^2 + 4\left(\frac{c}{d}\right)^2 rNmz^2T^nU^{k+n}, \end{aligned}$$

where

$$\begin{aligned} s_1 &= cqrNx^n + c(z^2mU^k - y^2lT^k)/q, \\ s_2 &= cqrNx^n - c(z^2mU^k - y^2lT^k)/q. \end{aligned}$$

Then we take

$$S_1 := (s_1 - td)/2 \quad \text{and} \quad S_2 := (s_2 - td)/2$$

where an infinite subset of I has been chosen so that t_n is fixed and we set $t = t_n$.

We also take:

$$\alpha = \omega_n + S_1/d, \quad \bar{\alpha} = \bar{\omega}_n + S_1/d \quad \text{and} \quad \beta = \omega_n + S_2/d, \quad \bar{\beta} = \bar{\omega}_n + S_2/d.$$

Then,

$$\alpha\bar{\alpha} = -c^2rNly^2T^kx^n/d^2 \quad \text{and} \quad \beta\bar{\beta} = -c^2rNmz^2U^kx^n/d^2.$$

Further to these equations, we also have

$$q^2d^2D_n = (qs_3)^2 - 4c^2ml(zy)^2(UT)^k$$

where

$$qs_3 = cq^2rNx^n + mz^2U^k + ly^2T^k.$$

Also of relevance will be

$$S_1 + S_2 + td = Ax^n = cqrNx^n.$$

We now detail the initial segment of the continued fraction expansion of ω_n . From this we shall discover that in some circumstances this segment is seriously longer than linear in n so that the complete expansion could not possibly constitute a keeper.

Before commencing we need to determine the common factors between some of the terms. First, we define $g := (s_1, s_2, d)$. It is also worth noting that $(z, s_1/c) \mid 2$. Since $z \mid E_n$, any common factor of z and s_1/c must divide $4rNly^2T^kx^n$. We already know that $(z, rNlyx) = 1$, hence the greatest common divisor divides 4. If $4 \mid z$ then $4 \mid s_2/c$ and so

$$E_n = d^2D_n/c^2 = (s_2/c)^2 + 4rNmz^2U^kx^n \equiv 0 \pmod{16}.$$

But, if $4 \mid s_1/c$ we have

$$E_n = d^2D_n/c^2 = (s_1/c)^2 + 4rNly^2T^kx^n \equiv 0 \pmod{16}$$

and so $16 \mid 4rNly^2T^kx^n$, which is impossible. Similarly, $(y, s_2/c)$ must divide 2.

Lemma 33. *We always have $g \mid 2$. Moreover, $g = 2$ if and only if $2 \mid d$.*

Proof. By definition, g divides $s_1 + s_2 = 2cqrNx^n$ and $s_1 - s_2 = 2c(mz^2U^k - ly^2T^k)$. Since $d^2 \mid E_n$ we have $(g, xN) = 1$ which implies that $(g, TUN) = 1$. Hence, $g \mid 2qrc$ and since $(c, d) = 1$ we have $g \mid 2qr$. From,

$$d^2D_n/c^2 = E_n = (s_1/c)^2 + 4rNly^2T^kx^n$$

and $g \mid s_1$ we have $g^2 \mid 4rNly^2T^kx^n$ which implies $g^2 \mid 4rly^2$. Thus $g \mid (2rq, 2y)$, and $(rq, y) = 1$ means that $g \mid 2$.

For the second part, if $g = 2$ then clearly $2 \mid d$. Next, if $2 \nmid d$ then $2 \mid s_1$ and $2 \mid s_2$ hence $2 \mid g$ so then $g = 2$. ■

16.3. Certain factors of d

We now define $d_y := (S_1 + td, d)$ and $d_z := (S_2 + td, d)$. Ultimately, we intend to show that this notation is justified by demonstrating that $d_y \mid y$. Before doing that, we let $d'_y := (s_1/g, d/g)$ and $\tau_y := d_y/d'_y$. Note that since $S_1 + td = (s_1 + td)/2$, when $g = 2$ we have $d'_y \mid d_y$. And when $g \neq 2$, $d'_y \mid (s_1 + td, d) = (2(S_1 + td), d) = d_y$ since in this case, $2 \nmid d$. Hence $\tau_y \in \mathbb{Z}$. Similarly, we define $d'_z := (s_2/g, d/g)$ and $\tau_z := d_z/d'_z$.

Lemma 34. $d'_y \mid y$ and $d'_z \mid z$.

Proof. We have

$$\left(\frac{d}{d'_z}\right)^2 D_n = \left(\frac{s_2}{d'_z}\right)^2 + \frac{4Hc^2x^n}{(d'_z)^2}. \quad (16.4)$$

Since $d^2 D_n = c^2 E_n$ and $(c, d) = 1$ we must have $d'_z \mid E_n$, whence $(d'_z, x) = 1$. It follows that $(d'_z)^2 \mid 4rmz^2$ because $(N, E_n) = 1$. If d'_z is odd then $d'_z \mid z$ since r and m are squarefree and relatively prime. Now suppose that d'_z is even. Also, we must have $2 \mid d/d'_z$ so then $2 \mid d$ and $g = 2$. Hence, $2d'_z = (s_2, d)$, therefore $2 \mid s_2/d'_z$. From (16.4) we have $4 \mid 4H/(d'_z)^2$ so $(d'_z)^2 \mid H$, therefore $d'_z \mid z$. By symmetry, $d'_y \mid y$ and $(d'_y, d'_z) = 1$. ■

By definition, $\tau_z = ((S_2 + td)/d'_z, d/d'_z)$. We also have the following result.

Lemma 35. $\tau_z \mid g$.

Proof. By definition,

$$g = \left(\frac{s_2}{d'_z}, \frac{d}{d'_z}\right) = \left(\frac{s_2}{d'_z} + \frac{td}{d'_z}, \frac{d}{d'_z}\right) = \left(\frac{2(S_2 + td)}{d'_z}, \frac{d}{d'_z}\right).$$

Hence $\tau_z \mid g$. ■

Moreover, $\tau_z = 2$ implies that $2 \parallel d/d'_z$, hence $2 \nmid d/d'_z$. Next, we write $d = \bar{d}d_z d_y$ (so when $\tau_z = 2$ we have $(2, \bar{d}d_y) = 1$). That $\bar{d} \in \mathbb{Z}$ follows from the next result.

Lemma 36. $\tau_y d'_y \mid y$ and $\tau_z d'_z \mid z$, in other words $d_y \mid y$ and $d_z \mid z$.

Proof. We treat the 2 cases of t separately. From Lemma 34 we have $d'_z \mid z$, thus we are reduced to treating the case of $\tau_z = 2$. First suppose that $t = 0$, then $S_2 = s_2/2$ and

$$d^2 D_n/4 = S_2^2 + c^2 r N m z^2 U^k x^n.$$

Furthermore, $\tau_z = 2$ implies $2 \parallel d/d'_z$, hence,

$$\frac{d^2 D_n}{4(d'_z)^2} - \left(\frac{S_2}{d'_z}\right)^2 = \frac{c^2 r N m z^2 U^k x^n}{(d'_z)^2}. \quad (16.5)$$

$\tau_z \mid d/d'_z$ by definition and $4 \mid D_n$ since $t = 0$ which imply that $4\tau_z^2(d'_z)^2 \mid d^2 D_n$. Moreover, $\tau_z \mid S_2/d'_z$, implies that τ_z^2 divides the left side of (16.5). Hence,

$$\tau_z^2 \mid c^2 r N m (z/d'_z)^2 U^k x^n.$$

From $\tau_z \mid d$ and $(d, cxN) = 1$ we must have $\tau_z^2 \mid rm(z/d'_z)^2$. But since r and m are coprime and squarefree, rm is squarefree. Hence $\tau_z \mid z/d'_z$, that is $\tau_z d'_z \mid z$.

Now suppose that $t = 1$. Then we have

$$d^2 D_n/4 = S_2^2 + tdS_2 + td^2/4 + c^2 r N m z^2 U^k x^n$$

because $s_2 = 2S_2 + td$. This is,

$$\frac{d^2 D_n}{4(d'_z)^2} - \frac{td^2}{4(d'_z)^2} - \frac{S_2}{d'_z} \left(\frac{S_2 + td}{d'_z} \right) = \frac{c^2 r N m z^2 U^k x^n}{(d'_z)^2}. \quad (16.6)$$

Now, $\tau_z = 2$ implies $2 \parallel d/d'_z$, so that $d^2 D_n / (4(d'_z)^2) \equiv 1 \pmod{4}$ and $td^2 / (4(d'_z)^2) \equiv 1 \pmod{4}$. Hence

$$4 \mid \frac{d^2 D_n - td^2}{4(d'_z)^2} \quad \text{which entails} \quad \tau_z^2 \mid \frac{d^2 D_n - td^2}{4(d'_z)^2}.$$

Also, $\tau_z \mid (S_2 + td)/d'_z$ and $\tau_z \mid d/d'_z$ by definition; hence $\tau_z \mid S_2/d'_z$. Thus,

$$\tau_z^2 \mid \frac{S_2}{d'_z} \left(\frac{S_2 + td}{d'_z} \right).$$

Therefore, τ_z^2 divides the left side of (16.6). Hence, $\tau_z^2 \mid c^2 r N m (z/(d'_z)^2) U^k x^n$ and, following the same reasoning as in the $t = 0$ case, we have that $d'_z \tau_z \mid z$, that is $d_z \mid z$. By symmetry, $d_y \mid y$. $\blacksquare$

16.4. Expansion of ω_n

We now commence to expand ω_n . Its continued fraction begins as

$$\begin{aligned} \omega_n &= (S_1 + td)/d - (\bar{\omega}_n + S_1/d) \\ &= \frac{(S_1 + td)/d_y}{d/d_y} - (\bar{\omega}_n + S_1/d) \end{aligned}$$

and $((S_1 + td)/d_y, d/d_y) = 1$ by the definition of d_y (in fact this is what motivates the definition of d_y). Hence we may apply Lemma 2 on page 8, and find that after the expansion of $S_1 + td/d$, of length h_0 , a new complete quotient in the continued fraction expansion is

$$\frac{-(-1)^{h_0+1}}{(\bar{\omega}_n + S_1/d)(d/d_y)^2} - \frac{c_0}{d/d_y}$$

where

$$c_0 \equiv \frac{(-1)^{h_0}}{(S_1 + td)/d_y} \pmod{d/d_y}.$$

By choosing $(-1)^{h_0+1} = \text{sign}(l)$ this element becomes

$$\frac{\omega_n + S_1/d}{c^2 r N |l| y^2 T^k x^n / (d_y)^2} - \frac{c_0}{d/d_y} = \frac{\omega_n + S_1/d - c_0 c^2 r N |l| d_y (y/d_y)^2 T^k x^n / d}{c^2 r N |l| (y/d_y)^2 T^k x^n}. \quad (16.7)$$

Now define:

$$\begin{aligned} s &:= \max_{i \in \mathbb{N}} \{(x^i, c)\}, \\ u &:= \frac{c}{s}. \end{aligned}$$

Hence $(u, x) = 1$. Recall, $(x, qrz yml) = 1$ so then $(s, qrz yml) = 1$.

We will denote the element in (16.7) as θ_{h_0} . From it, we can write

$$\theta_{h_0} = \frac{\omega_n + P_{h_0}}{Q_{h_0}} = \frac{A_0}{B_0} - \frac{\bar{\beta}}{c^2 r N |l| (y/d_y)^2 T^k x^n} \quad (16.8)$$

where

$$\begin{aligned} A_0 &= c q r N x^n - c_0 c^2 r N |l| d_y (y/d_y)^2 T^k x^n, \\ B_0 &= c^2 d r N |l| (y/d_y)^2 T^k x^n. \end{aligned}$$

Next, we define $\Delta_0 := (A_0, B_0)$. We need to determine Δ_0 before we can apply Lemma 2 again. Then,

$$\Delta_0 = crNx^n(q - c_0c|l|d_y(y/d_y)^2T^k, cd|l|(y/d_y)^2T^k),$$

and define, $\delta := (c, q)$, $G_0 := A_0/(\delta crNx^n)$. Then

$$\Delta_0 = crNx^n\delta(G_0, d).$$

Next, set $d_1 := (G_0, d)$. So $\Delta_0 = crNx^n\delta d_1$. Since $A_0 = dP_{h_0} + (s_2 + td)/2$, we have

$$(d, A_0) = (d, (s_2 + td)/2) = (d, S_2 + td) = d_z.$$

Since $G_0|A_0$ we can conclude that $d_1|d_z$. Moreover,

$$q - c_0c|l|d_y(y/d_y)^2T^kx^n \equiv q + \frac{cd_y(y/d_y)^2T^k}{S_1 + td} \pmod{d_z} \quad (16.9)$$

by the value of c_0 .

First note that $(d_z, S_1) = 1$. For let $p|(d_z, S_1)$ with $p > 1$ and recall $d_z|S_2$ and $d_z|d$. Hence $p|S_1 + S_2 + td = cqrNx^n$. Since $(z, qrNx) = 1$ we must have $p|c$. But then $p|(c, d)$ which is a contradiction. Hence, we have that $S_1 + td$ is invertible modulo d_z , so (16.9) becomes,

$$\begin{aligned} q - c_0c|l|d_y(y/d_y)^2T^kx^n &\equiv \frac{q(S_1 + td) + cly^2T^k}{S_1 + td} \pmod{d_z} \\ &\equiv \frac{cq^2rNx^n + cmz^2U^k + cly^2T^k + tdq}{2(S_1 + td)} \pmod{d_z} \\ &\equiv \frac{qs_3}{S_1 + td} \pmod{d_z} \\ &\equiv 0 \pmod{d_z} \end{aligned}$$

Hence $d_1 = d_z$. In summary,

$$\Delta_0 = crNx^n\delta d_z \quad \text{and} \quad \frac{B_0}{\Delta_0} = \frac{cd|l|(y/d_y)^2T^k}{\delta d_z}.$$

From (16.8), by applying Lemma 2, we find the next partial quotients are the expansion of A_0/B_0 of length p_0 . By choosing $(-1)^{p_0+1} = \text{sign}(m)$, the next element in the continued fraction expansion is θ_{h_1} where $h_1 := h_0 + p_0$ and

$$\theta_{h_1} = \frac{c^2rN|l|(y/d_y)^2T^kx^n}{-\beta \text{sign}(m)(B_0/\Delta_0)^2} - \frac{c_1}{B_0/\Delta_0}$$

where $c_1 \equiv -\text{sign}(m)\Delta_0/A_0 \pmod{B_0/\Delta_0}$,

$$= \frac{\omega_n + S_2/d - c_1(c/\delta)|m|z(z/d_z)U^k/d}{(c/\delta)^2|ml|(y/d_y)^2(z/d_z)^2(UT)^k}. \quad (16.10)$$

16.5. General complete quotients

For induction purposes, we suppose that we have a complete quotient satisfying

$$P_{h_{2i-1}} = S_2/d - sm_{2i-1}r_{2i-1}u_{2i-1}u'_{2i-1}z_{2i-1}z'_{2i-1}c_{2i-1}U^{ki}/d \quad (16.11a)$$

$$Q_{h_{2i-1}} = r_{2i-1}l_{2i-1}m_{2i-1}(su_{2i-1}y_{2i-1}z_{2i-1})^2(UT)^{ki} \quad (16.11b)$$

where:

$$\begin{aligned} m_{2i-1} &\in \{1, |m|\}, \quad l_{2i-1} \in \{1, |l|\}, \quad r_{2i-1} \in \{1, r\}, \\ y_{2i-1} &|y/d_y, \quad z_{2i-1}|z/d_z, \quad z'_{2i-1}|z, \quad u_{2i-1}|u, \quad u'_{2i-1}|u, \\ (z_{2i-1}y_{2i-1}, u/u_{2i-1}) &= 1, \quad (c_{2i-1}, sT) = 1. \end{aligned}$$

The complete quotient, θ_{h_1} , in (16.10) is the case of $i = 1$ with the selections:

$$\begin{aligned} r_1 &= 1, \quad l_1 = |l|, \quad m_1 = |m|, \quad z_1 = z/d_z, \quad z'_1 = z, \\ y_1 &= y/d_y, \quad u_1 = u/\delta, \quad u'_1 = 1. \end{aligned}$$

Note that $u/u_1 = \delta$ and $\delta|q$, hence $(z_1 y_1, u/u_1) = 1$. Also observe that since $sT|B_0/\Delta_0$ and $(B_0/\Delta_0, A_0/\Delta_0) = 1$ we have $(c_1, sT) = 1$.

From (16.11), we find that the current line in the continued fraction expansion is

$$\frac{\omega_n + P_{h_{2i-1}}}{Q_{h_{2i-1}}} = \frac{cqrNx^n - sr_{2i-1}m_{2i-1}u_{2i-1}u'_{2i-1}z_{2i-1}z'_{2i-1}c_{2i-1}U^{ki}}{dr_{2i-1}m_{2i-1}l_{2i-1}(su_{2i-1}z_{2i-1}y_{2i-1})^2(UT)^{ki}} - \frac{(\bar{\omega}_n + S_1/d)}{r_{2i-1}m_{2i-1}l_{2i-1}(su_{2i-1}z_{2i-1}y_{2i-1})^2(UT)^{ki}}.$$

Now, define:

$$\begin{aligned} A_{2i-1} &:= cqrNx^n - sr_{2i-1}m_{2i-1}u_{2i-1}u'_{2i-1}z_{2i-1}z'_{2i-1}c_{2i-1}U^{ki}, \\ B_{2i-1} &:= dr_{2i-1}m_{2i-1}l_{2i-1}(su_{2i-1}z_{2i-1}y_{2i-1})^2(UT)^{ki}, \\ \Delta_{2i-1} &:= (A_{2i-1}, B_{2i-1}). \end{aligned}$$

16.6. Some helpful results

Theorem 37. *If $A_{2i-1} = dP_{h_{2i-1}} + (s_1 + td)/2$ then $d_y u_{2i-1} l_{2i-1} y_{2i-1} | A_{2i-1}$.*

Proof. From the element,

$$\theta_{h_{2i-1}} = \frac{\omega_n + P_{h_{2i-1}}}{Q_{h_{2i-1}}} = \frac{\sqrt{D_n} + 2P_{h_{2i-1}} + t}{2Q_{h_{2i-1}}}$$

we have

$$-D_n + (2P_{h_{2i-1}} + t)^2 = -4Q_{h_{2i-1}-1}Q_{h_{2i-1}}.$$

Moreover, $d_y^2 u_{2i-1}^2 l_{2i-1} y_{2i-1}^2$ divides $d^2 D_n$ and $d^2 Q_{h_{2i-1}}$, hence

$$d_y u_{2i-1} l_{2i-1} y_{2i-1} | d(2P_{h_{2i-1}} + t). \quad (16.12)$$

Also,

$$A_{2i-1} = dP_{h_{2i-1}} + (s_1 + td)/2.$$

Since $y_{2i-1} | y/d_y$ we have $d_y u_{2i-1} l_{2i-1} y_{2i-1} | s_1$, hence

$$d_y u_{2i-1} l_{2i-1} y_{2i-1} | 2A_{2i-1} - s_1 \text{ which implies } d_y u_{2i-1} l_{2i-1} y_{2i-1} | 2A_{2i-1}.$$

Thus we are only left to show that if $2^a | d_y u_{2i-1} l_{2i-1} y_{2i-1}$ then $2^a | A_{2i-1}$, where $a > 0$.

Lemma 38. *If $a \geq 1$ and $2^a | d_y u_{2i-1} l_{2i-1} y_{2i-1}$ then $2^a | A_{2i-1}$.*

Proof. First observe that $2^a | d_y u_{2i-1} l_{2i-1} y_{2i-1}$ implies $2^a | cly$ thus $2^{2a-1} | c^2 l y^2$ and $2^a | s_1$. To begin with, if $2 | u_{2i-1}$ then $2 \nmid d$ and $4 | D_n$ so $t = 0$. We split the proof into 2 cases according to the value of t .

First consider the case of $t = 1$, so $2 \nmid u_{2i-1}$. From above,

$$d_y l_{2i-1} y_{2i-1} | d(2P_{h_{2i-1}} + 1).$$

Without loss of generality, let us suppose that $2^a | d_y l_{2i-1} y_{2i-1}$. Then $2^a | s_1$ and since D_n is odd, $2^a | d$ as well. Hence d'_y which is equal to $(s_1/2, d/2)$ must be divisible by 2^{a-1} (recall Lemma 36 on page 73, $2 | d$ if and only if $g = 2$). Thus, either $2 | l_{2i-1} y_{2i-1}$ or $2^a | d'_y$. First, suppose that $2^a \nmid d'_y$, that is $2 | l_{2i-1} y_{2i-1}$. Then

$2^a \mid ly$ which means $2^{2a+1} \mid 4ly^2$. Furthermore, $d'_y = (s_1/2, d/2)$ is exactly divisible by 2^{a-1} implies that at least one of d and s_1 is exactly divisible by 2^a . Then,

$$d^2 D_n = s_1^2 + 4c^2 r N ly^2 T^k x^n$$

means that $2^{a+1} \mid s_1$ if and only if $2^{a+1} \mid d$, because D_n is odd. Hence $2^a \parallel s_1$ and $2^a \parallel d$. Then

$$A_{2i-1} = dP_{h_{2i-1}} + (s_1 + td)/2 \equiv 0 \pmod{2^a}$$

as we required.

Second, suppose that $2^a \mid d'_y$. Since $d'_y \mid y$ we have $2^a \mid y$. Then from

$$s_1^2 = d^2 D_n - 4c^2 r N ly^2 T^k x^n$$

we again have $2^{a+1} \mid s_1$ if and only if $2^{a+1} \mid d$. Hence,

$$(s_1 + td)/2 \equiv 0 \pmod{2^a}$$

so $2^a \mid A_{2i-1}$.

Now consider the case of $t = 0$, from (16.12),

$$d_y u_{2i-1} l_{2i-1} y_{2i-1} \mid 2dP_{h_{2i-1}}$$

and

$$2A_{2i-1} = 2dP_{h_{2i-1}} + s_1.$$

Suppose that $2^a \mid d_y u_{2i-1} l_{2i-1} y_{2i-1}$ and $2^a \nmid A_{2i-1}$. Then from above, precisely one of $2dP_{h_{2i-1}}$ and s_1 is exactly divisible by 2^a . Recall that $\tau_y d'_y l_{2i-1} y_{2i-1} u_{2i-1} \mid lyc$, so then $\tau_y d'_y l_{2i-1} y_{2i-1} u_{2i-1} \mid s_1$.

Let us first suppose that $2^a \parallel s_1$ and $2 \nmid d_y$. If $g = 2$ then $2 \nmid u_{2i-1}$ and recall that $2 \nmid d_y$ implies that we have $\tau_y = 1$. That is,

$$2 = g = \left(\frac{2(S_1 + td)}{d'_y}, \frac{d}{d'_y} \right) \quad \text{and} \quad 1 = \tau_y = \left(\frac{S_1 + td}{d'_y}, \frac{d}{d'_y} \right).$$

So, $2 \mid d/d'_y$ and $2 \nmid (S_1 + td)/d'_y = s_1/(2d'_y)$. Since $2 \nmid d'_y$ we have that $2 \parallel s_1$. However,

$$s_1^2 = d^2 D_n - 4c^2 r N ly^2 T^k x^n \equiv 0 \pmod{8} \quad (16.13)$$

implies that $4 \mid s_1$ which is a contradiction.

If $g = 1$ then $2 \nmid d$ and from,

$$d^2 D_n = s_1^2 + 4c^2 r N ly^2 T^k x^n$$

we have $2^{2a} \parallel D_n$. From the earlier assumption, $2^{a+1} \mid 2dP_{h_{2i-1}}$, that is $2^a \mid P_{h_{2i-1}}$. Since $t = 0$, we have

$$D_n - 4P_{h_{2i-1}}^2 = 4Q_{h_{2i-1}-1}Q_{h_{2i-1}},$$

which gives

$$D_n = 4Q_{h_{2i-1}-1}Q_{h_{2i-1}} + 4P_{h_{2i-1}}^2 \equiv 0 \pmod{2^{2a+1}}$$

which is a contradiction.

Now suppose that $2^a \parallel s_1$, $2^{a+1} \mid 2dP_{h_{2i-1}}$, $2 \mid d_y$, $2 \nmid u_{2i-1}$ and $2^a \mid d_y l_{2i-1} y_{2i-1}$. Hence, $2^a \mid ly$ so $2^a \parallel ly$ since $ly \mid s_1$, so $2^a \parallel d_y l_{2i-1} y_{2i-1}$, because this term divides ly . From,

$$d^2 D_n = s_1^2 + 4c^2 r N ly^2 T^k x^n$$

we have $2^{2a} \parallel d^2 D_n$. Now let $2^b \parallel d_y$ with $a \geq b$, so $2^{a-b} \mid l_{2i-1} y_{2i-1}$. It follows that $2^{2a-2b} \parallel (\bar{d}d_z)^2 D_n$ and since $4 \mid D_n$ we must have $a - b > 0$. Hence, $2^{2a-2b-1} \mid Q_{h_{2i-1}}$. Now,

$$-d^2 D_n + d^2 (2P_{h_{2i-1}} + t)^2 = -4d^2 Q_{h_{2i-1}-1}Q_{h_{2i-1}}$$

so $2^{2a} \parallel -4d^2 Q_{h_{2i-1}-1}Q_{h_{2i-1}}$ which means $2^{2a-2b-2} \parallel (\bar{d}d_z)^2 Q_{h_{2i-1}-1}Q_{h_{2i-1}}$ which is a contradiction. In conclusion, it is not possible to have $2^a \parallel s_1$ and $2^{a+1} \mid 2dP_{h_{2i-1}}$.

Now consider, $2^{a+1} \mid s_1$ and $2^a \parallel 2dP_{h_{2i-1}}$. Then from

$$d^2 D_n = s_1^2 + 4c^2 r N l y^2 T^k x^n \equiv 0 \pmod{2^{2a+1}}$$

we have $2^{2a+1} \mid d^2 D_n$. Let $2^b \parallel d_y$ with $b \geq 0$, so $2^{a-b} \mid u_{2i-1} l_{2i-1} y_{2i-1}$. Then,

$$2^{2a-2b+1} \mid (\bar{d} d_z)^2 D_n \text{ and } 4 \mid D_n \text{ implies } 2a - 2b + 1 \geq 2.$$

Thus $a > b$. Hence, $2^{2a-2b-1} \mid Q_{h_{2i-1}}$. We also have,

$$-d^2 D_n + 4d^2 P_{h_{2i-1}} = -4d^2 Q_{h_{2i-1}-1} Q_{h_{2i-1}}.$$

Since $2^{2a} \parallel 4d^2 P_{h_{2i-1}}$ we must have $2^{2a} \parallel 4d^2 Q_{h_{2i-1}-1} Q_{h_{2i-1}}$, that is

$$2^{2a-2b-2} \parallel (d_z \bar{d})^2 Q_{h_{2i-1}-1} Q_{h_{2i-1}}$$

which is again a contradiction. Thus the original assumption that $2^a \nmid A_{2i-1}$ must be wrong. $\blacksquare$

By the above Lemma we have $d_y u_{2i-1} l_{2i-1} y_{2i-1} \mid A_{2i-1}$. $\blacksquare$

In the procedure for expanding ω_n , the next lemma will prove useful,

Lemma 39. *For any rational integers a, b, d, f such that $d \mid ab$ and $(f, d) = 1$ there exist rational integers x, y such that*

$$dxy = ab$$

and $x \mid a, y \mid b, (fx, b/y) = 1$

Proof. Just take $x = a/(a, d)$ and $y = b(a, d)/d$. $\blacksquare$

16.7. Determining Δ_{2i-1}

Returning to the expansion of ω_n , we write:

$$\begin{aligned} w_{2i-1} &:= (m_{2i-1}, u/u_{2i-1}), \\ e_{2i-1} &:= s d_y u_{2i-1} l_{2i-1} y_{2i-1} r_{2i-1} w_{2i-1}. \end{aligned}$$

We note, by the formula for A_{2i-1} , that if $n \geq ki$ then

$$w_{2i-1} u_{2i-1} s U^{ki} r_{2i-1} \mid A_{2i-1}$$

and by Theorem 37,

$$d_y u_{2i-1} l_{2i-1} y_{2i-1} \mid A_{2i-1}.$$

Now, $(ly, smrU) = 1$ implies $(d_y l_{2i-1} y_{2i-1}, s w_{2i-1} U^{ki} r_{2i-1}) = 1$. Hence,

$$\begin{aligned} &\frac{w_{2i-1} u_{2i-1} s U^{ki} r_{2i-1} d_y u_{2i-1} l_{2i-1} y_{2i-1}}{(w_{2i-1} u_{2i-1} s U^{ki} r_{2i-1}, d_y u_{2i-1} l_{2i-1} y_{2i-1})} \mid A_{2i-1}, \\ &w_{2i-1} u_{2i-1} s U^{ki} r_{2i-1} d_y l_{2i-1} y_{2i-1} \mid A_{2i-1}, \\ &U^{ki} e_{2i-1} \mid A_{2i-1}. \end{aligned}$$

We have determined

$$\begin{aligned} \frac{B_{2i-1}}{U^{ki} e_{2i-1}} &= \frac{dr_{2i-1} l_{2i-1} m_{2i-1} (s u_{2i-1} y_{2i-1} z_{2i-1})^2 (UT)^{ki}}{s d_y u_{2i-1} l_{2i-1} y_{2i-1} r_{2i-1} w_{2i-1} U^{ki}} \\ &= \frac{d s m_{2i-1} u_{2i-1} y_{2i-1} z_{2i-1}^2 T^{ki}}{d_y w_{2i-1}} \end{aligned}$$

and

$$\begin{aligned} A_{2i-1} &= s u_{2i-1} r_{2i-1} w_{2i-1} U^{ki} \times \\ &\left(\frac{ur}{u_{2i-1} r_{2i-1} w_{2i-1}} q N T^n U^{n-ki} - \frac{m_{2i-1}}{w_{2i-1}} u'_{2i-1} z_{2i-1} z'_{2i-1} c_{2i-1} \right). \end{aligned}$$

Hence,

$$\frac{A_{2i-1}}{e_{2i-1}U^{ki}} = \frac{urqNT^nU^{n-ki} - u_{2i-1}r_{2i-1}m_{2i-1}u'_{2i-1}z_{2i-1}z'_{2i-1}c_{2i-1}}{u_{2i-1}r_{2i-1}w_{2i-1}l_{2i-1}y_{2i-1}}.$$

Now we define,

$$G_{2i-1} := \frac{A_{2i-1}}{U^{ki}e_{2i-1}}.$$

Any common factor of z_{2i-1} and G_{2i-1} must divide u/u_{2i-1} , but we have the condition $(z_{2i-1}, u/u_{2i-1}) = 1$ so $(G_{2i-1}, z_{2i-1}) = 1$.

Also, any common factor of G_{2i-1} and m_{2i-1}/w_{2i-1} must divide $u/u_{2i-1}w_{2i-1}$. But

$$\left(\frac{m_{2i-1}}{w_{2i-1}}, \frac{u}{u_{2i-1}w_{2i-1}} \right) = 1$$

by the definition of w_{2i-1} . Hence, $(G_{2i-1}, m_{2i-1}/w_{2i-1}) = 1$.

Also note that if $p|(sT, G_{2i-1})$ then, when $n > ki$,

$$p|m_{2i-1}u'_{2i-1}z_{2i-1}z'_{2i-1}c_{2i-1}.$$

Since $(c_{2i-1}, sT) = 1$, we must have $(s, G_{2i-1}) = (T, G_{2i-1}) = 1$. In summary,

$$\begin{aligned} \Delta_{2i-1} &= U^{ki}e_{2i-1} \left(\frac{A_{2i-1}}{e_{2i-1}U^{ki}}, \frac{B_{2i-1}}{e_{2i-1}U^{ki}} \right) \\ &= U^{ki}e_{2i-1}(G_{2i-1}, \bar{d}d_z y_{2i-1}u_{2i-1}). \end{aligned}$$

Since,

$$A_{2i-1} = dP_{h_{2i-1}} + S_1 + td$$

we have, $(d, A_{2i-1}) = d_y$. Thus $(\bar{d}d_z, G_{2i-1}) = 1$. Hence,

$$\Delta_{2i-1} = U^{ki}e_{2i-1}\bar{\Delta}_{2i-1}$$

where $\bar{\Delta}_{2i-1} := (y_{2i-1}u_{2i-1}, G_{2i-1})$.

This completes the investigation of Δ_{2i-1} . Note that we have determined,

$$\frac{B_{2i-1}}{\Delta_{2i-1}} = \frac{dsm_{2i-1}u_{2i-1}y_{2i-1}z_{2i-1}^2T^{ki}}{w_{2i-1}d_y\bar{\Delta}_{2i-1}}.$$

16.8. Continuing the expansion of ω_n

From the complete quotient

$$\theta_{h_{2i-1}} = \frac{A_{2i-1}}{B_{2i-1}} - \frac{\bar{\alpha}}{r_{2i-1}l_{2i-1}m_{2i-1}(su_{2i-1}y_{2i-1}z_{2i-1})^2(UT)^{ki}}$$

we apply Lemma 2, so the next partial quotients are the expansion of A_{2i-1}/B_{2i-1} of length p_{2i-1} . The parity of p_{2i-1} is determined by $(-1)^{p_{2i-1}+1} = \text{sign}(l)$. Following this, the next complete quotient is $\theta_{h_{2i}}$ where

$$h_{2i} := h_{2i-1} + p_{2i-1}.$$

By Lemma 2, $\theta_{h_{2i}}$ is equal to

$$\begin{aligned} &\frac{r_{2i-1}l_{2i-1}m_{2i-1}(su_{2i-1}y_{2i-1}z_{2i-1})^2(UT)^{ki}}{-\bar{\alpha}(B_{2i-1}/\Delta_{2i-1})^2} - \frac{c_{2i}}{B_{2i-1}/\Delta_{2i-1}} = \\ &\frac{r_{2i-1}l_{2i-1}(\omega_n + S_1/d)(w_{2i-1}d_y\bar{\Delta}_{2i-1})^2}{s^2u^2rN|l|y^2m_{2i-1}z_{2i-1}^2U^{n-ki}T^{n+k(i+1)}} - \frac{c_{2i}}{B_{2i-1}/\Delta_{2i-1}} \quad (16.14) \end{aligned}$$

where

$$c_{2i} \equiv -\text{sign}(l) \Delta_{2i-1}/A_{2i-1} \pmod{B_{2i-1}/\Delta_{2i-1}}.$$

Also note that $sT|B_{2i-1}/\Delta_{2i-1}$ and $(A_{2i-1}/\Delta_{2i-1}, B_{2i-1}/\Delta_{2i-1}) = 1$ imply that $(c_{2i}, sT) = 1$.

The point of Lemma 39 is to replace the product $d_y w_{2i-1} \bar{\Delta}_{2i-1}$ with one dependent on terms involving y and u . More specifically, by replacing the terms in the Lemma known as f, a, b, d by $z_{2i-1}, y/d_y, u/w_{2i-1}, \bar{\Delta}_{2i-1}$ respectively, then the Lemma says (note that f and d are coprime because $(z_{2i-1}, G_{2i-1}) = 1$ implies $(z_{2i-1}, \bar{\Delta}_{2i-1}) = 1$) that there exists two numbers, say y_{2i} and u_{2i} , such that

$$y_{2i}|y/d_y, \quad u_{2i}|u/w_{2i-1}, \quad y_{2i}u_{2i}\bar{\Delta}_{2i-1} = \frac{y}{d_y} \frac{u}{w_{2i-1}}$$

and

$$\left(z_{2i-1}y_{2i}, \frac{u}{u_{2i}w_{2i-1}} \right) = 1.$$

In other words, $yu = d_y w_{2i-1} y_{2i} u_{2i} \bar{\Delta}_{2i-1}$. Consequently, the first term in (16.14) can be written as

$$\frac{l_{2i-1}}{|l|} \frac{r_{2i-1}}{r} \frac{(\omega_n + S_1/d)}{s^2 y_{2i}^2 u_{2i}^2 m_{2i-1} z_{2i-1}^2 N T^{n+k(i+1)} U^{n-ki}},$$

which by taking:

$$l_{2i} := \frac{|l|}{l_{2i-1}}, \quad r_{2i} := \frac{r}{r_{2i-1}}, \quad m_{2i} := m_{2i-1}, \quad z_{2i} := z_{2i-1},$$

becomes

$$\frac{\omega_n + S_1/d}{m_{2i} r_{2i} l_{2i} (s u_{2i} y_{2i} z_{2i})^2 N T^{n+k(i+1)} U^{n-ki}}.$$

Also observe that

$$\left(z_{2i-1}y_{2i}, \frac{u}{u_{2i}w_{2i-1}} \right) = 1 \text{ implies that } \left(z_{2i}y_{2i}, \frac{u}{u_{2i}w_{2i-1}} \right) = 1$$

and $(w_{2i-1}, y_{2i}) = 1$ implies that $(y_{2i}, u/u_{2i}) = 1$. Finally, $(z_{2i}, u/u_{2i-1}) = 1$ implies that $(z_{2i}, w_{2i-1}) = 1$ and so

$$(y_{2i} z_{2i}, u/u_{2i}) = 1.$$

Now the second term in (16.14) is

$$\begin{aligned} \frac{c_{2i} w_{2i-1} d_y \bar{\Delta}_{2i-1}}{d s m_{2i-1} u_{2i-1} y_{2i-1} z_{2i-1}^2 T^{ki}} &= \frac{c_{2i} y u}{d s y_{2i} u_{2i} m_{2i-1} u_{2i-1} y_{2i-1} z_{2i-1}^2 T^{ki}} \\ &= \frac{y}{y_{2i-1}} \frac{u}{u_{2i-1}} \frac{c_{2i}}{d s m_{2i} u_{2i} y_{2i} z_{2i}^2 T^{ki}}. \end{aligned} \quad (16.15)$$

If we take:

$$y'_{2i} := \frac{y}{y_{2i-1}}, \quad u'_{2i} := \frac{u}{u_{2i-1}},$$

the term in (16.15) becomes

$$\frac{c_{2i} y'_{2i} u'_{2i}}{d s m_{2i} u_{2i} y_{2i} z_{2i}^2 T^{ki}} = \frac{N s u_{2i} u'_{2i} l_{2i} r_{2i} y_{2i} y'_{2i} c_{2i} U^{n-ki} T^{n+k}}{d r_{2i} l_{2i} m_{2i} (s u_{2i} y_{2i} z_{2i})^2 N U^{n-ki} T^{n+k(i+1)}}.$$

In conclusion, the entire element is

$$\theta_{h_{2i}} = \frac{\omega_n + S_1/d - N s r_{2i} l_{2i} u_{2i} u'_{2i} y_{2i} y'_{2i} c_{2i} U^{n-ki} T^{n+k}/d}{r_{2i} l_{2i} m_{2i} (s u_{2i} y_{2i} z_{2i})^2 N T^{n+k(i+1)} U^{n-ki}} \quad (16.16)$$

with

$$\begin{aligned} m_{2i} &\in \{1, |m|\}, \quad l_{2i} \in \{1, |l|\}, \quad r_{2i} \in \{1, r\}, \quad u_{2i}|u, \quad u'_{2i}|u, \\ z_{2i}|z/d_z, \quad y_{2i}|y/d_y, \quad y'_{2i}|y, \quad (c_{2i}, sT) &= 1, \quad (z_{2i} y_{2i}, u/u_{2i}) = 1. \end{aligned}$$

16.9. Repeating the procedure on $\theta_{h_{2i}}$

The above procedure can now be repeated on $\theta_{h_{2i}}$. From (16.16), we find a new line in the continued fraction expansion,

$$\frac{\omega_n + P_{h_{2i}}}{Q_{h_{2i}}} = \frac{cqrNx^n - Nsr_{2i}l_{2i}u_{2i}u'_{2i}y_{2i}y'_{2i}c_{2i}U^{n-ki}T^{n+k}}{dr_{2i}l_{2i}m_{2i}(su_{2i}y_{2i}z_{2i})^2NT^{n+k(i+1)}U^{n-ki}} - \frac{(\bar{\omega}_n + S_2/d)}{r_{2i}l_{2i}m_{2i}(su_{2i}y_{2i}z_{2i})^2NT^{n+k(i+1)}U^{n-ki}}$$

Now, define:

$$\begin{aligned} A_{2i} &:= cqrNx^n - Nsr_{2i}l_{2i}u_{2i}u'_{2i}y_{2i}y'_{2i}c_{2i}U^{n-ki}T^{n+k}, \\ B_{2i} &:= dr_{2i}l_{2i}m_{2i}(su_{2i}y_{2i}z_{2i})^2NT^{n+k(i+1)}U^{n-ki}, \\ \Delta_{2i} &:= (A_{2i}, B_{2i}). \end{aligned}$$

Before the next element in the continued fraction expansion can be determined, we need to evaluate Δ_{2i} .

We set,

$$\begin{aligned} w_{2i} &:= (l_{2i}, u/u_{2i}), \\ e_{2i} &:= sd_z u_{2i} m_{2i} z_{2i} r_{2i} w_{2i}. \end{aligned}$$

By using the same method of proof as that used in Theorem 37, we find that $d_z u_{2i} m_{2i} z_{2i} | A_{2i}$ and $A_{2i} = dP_{h_{2i}} + (s_2 + td)/2$. Since $su_{2i}w_{2i} | c$, $su_{2i}w_{2i} | sl_{2i}u_2$ and $r_{2i} | r$ we have that $sr_{2i}u_{2i}w_{2i} | A_{2i}$. So,

$$\frac{sd_z u_{2i} m_{2i} z_{2i} r_{2i} u_{2i} w_{2i}}{(d_z u_{2i} m_{2i} z_{2i}, sr_{2i} u_{2i} w_{2i})} \Big| A_{2i} \text{ which implies } \frac{sd_z u_{2i} m_{2i} z_{2i} r_{2i} w_{2i}}{(d_z m_{2i} z_{2i}, sr_{2i} w_{2i})} \Big| A_{2i}.$$

Thus $e_{2i} | A_{2i}$ because $(d_z m_{2i} z_{2i}, sr_{2i} w_{2i}) = 1$.

We also obviously have that $NT^n U^{n-ki} | \Delta_{2i}$. Then,

$$\begin{aligned} \frac{B_{2i}}{e_{2i} T^n U^{n-ki} N} &= \frac{dr_{2i}l_{2i}m_{2i}(su_{2i}y_{2i}z_{2i})^2 NT^{n+k(i+1)} U^{n-ki}}{sd_z u_{2i} m_{2i} z_{2i} r_{2i} w_{2i} NT^n U^{n-ki}} \\ &= \frac{dl_{2i}su_{2i}z_{2i}y_{2i}^2 T^{k(i+1)}}{d_z w_{2i}} \end{aligned}$$

and

$$A_{2i} = su_{2i}r_{2i}w_{2i}NT^n U^{n-ki} \left(\frac{ur}{u_{2i}r_{2i}w_{2i}} qU^{ki} - \frac{l_{2i}}{w_{2i}} u'_{2i}y_{2i}y'_{2i}c_{2i}T^k \right).$$

Hence,

$$\frac{A_{2i}}{e_{2i} T^n U^{n-ki} N} = \frac{urqU^{ki} - u_{2i}r_{2i}l_{2i}u'_{2i}y_{2i}y'_{2i}c_{2i}T^k}{d_z u_{2i}r_{2i}m_{2i}z_{2i}w_{2i}}.$$

Now define,

$$G_{2i} := \frac{A_{2i}}{e_{2i} T^n U^{n-ki} N}.$$

Since $(y, qrx) = 1$ any common factor of y_{2i} and G_{2i} must divide u/u_{2i} , but this is impossible by the conditions given for $\theta_{h_{2i}}$ in (16.16). Thus, $(G_{2i}, y_{2i}) = 1$. Moreover, any common factor of G_{2i} and l_{2i}/w_{2i} would need to divide $u/(u_{2i}w_{2i})$. But

$$\left(\frac{l_{2i}}{w_{2i}}, \frac{u}{u_{2i}w_{2i}} \right) = 1$$

by the definition of w_{2i} . Hence, $(G_{2i}, l_{2i}/w_{2i}) = 1$.

Next if $p|(T, G_{2i})$ then $p|urqU^{ki}$ which is impossible. Any common factor of U and G_{2i} must divide $u_{2i}r_{2i}l_{2i}u'_{2i}y_{2i}y'_{2i}c_{2i}T^k$ which is impossible because $(c_{2i}, U) = 1$. Hence $(G_{2i}, UT) = 1$, and so we have $(G_{2i}, s) = 1$. Lastly,

$$(d, A_{2i}) = (d, (s_2 + td)/2) = (d, S_2 + td) = d_z.$$

And since $d_z|e_{2i}$ we have that $(G_{2i}, \bar{d}d_y) = 1$. So,

$$\begin{aligned} \Delta_{2i} &= NT^n U^{n-ki} e_{2i}(G_{2i}, \bar{d}d_y s l_{2i} u_{2i} z_{2i} y_{2i} T^{k(i+1)}/w_{2i}) \\ &= NT^n U^{n-ki} e_{2i}(G_{2i}, z_{2i} u_{2i}) \\ &= NT^n U^{n-ki} e_{2i} \bar{\Delta}_{2i} \end{aligned}$$

where $\bar{\Delta}_{2i} := (z_{2i} u_{2i}, G_{2i})$.

We have determined,

$$\frac{B_{2i}}{\Delta_{2i}} = \frac{d s l_{2i} u_{2i} z_{2i} y_{2i}^2 T^{k(i+1)}}{w_{2i} d_z \bar{\Delta}_{2i}}.$$

Replacing the symbols, f, a, b, d , in Lemma 39 by $y_{2i}, z/d_z, u/w_{2i}$ and $\bar{\Delta}_{2i}$ respectively we see that there exists two numbers, say z_{2i+1} and u_{2i+1} , such that

$$z_{2i+1}|z/d_z, \quad u_{2i+1}|u/w_{2i}, \quad z_{2i+1} u_{2i+1} \bar{\Delta}_{2i} = \frac{zu}{d_z w_{2i}}$$

and

$$\left(y_{2i} z_{2i+1}, \frac{u}{u_{2i+1} w_{2i}} \right) = 1.$$

Now define $y_{2i+1} := y_{2i}$, which means that $(y_{2i+1} z_{2i+1}, u/(u_{2i+1} w_{2i})) = 1$. Then, $(w_{2i}, z) = 1$ because $(z, l) = 1$ and $(w_{2i}, y_{2i+1}) = 1$ since $(y_{2i}, u/u_{2i}) = 1$. In conclusion, we have $(y_{2i+1} z_{2i+1}, u/u_{2i+1}) = 1$. This completes the investigation of $\bar{\Delta}_{2i}$.

16.10. Finding another complete quotient

From,

$$\theta_{h_{2i}} = \frac{A_{2i}}{B_{2i}} - \frac{\bar{\beta}}{r_{2i} l_{2i} m_{2i} (s u_{2i} y_{2i} z_{2i})^2 T^{n+k(i+1)} U^{n-ki}}$$

we apply Lemma 2 and find that the next partial quotients are the expansion of A_{2i}/B_{2i} of length p_{2i} where the parity of p_{2i} is determined by $(-1)^{p_{2i}+1} = \text{sign}(m)$, and a new complete quotient in the expansion is $\theta_{h_{2i+1}}$ where

$$h_{2i+1} := h_{2i} + p_{2i}$$

and

$$\theta_{h_{2i+1}} = \frac{r_{2i} l_{2i} m_{2i} (s u_{2i} y_{2i} z_{2i})^2 N U^{n-ki} T^{n+k(i+1)}}{-\bar{\beta} \text{sign}(m) (B_{2i}/\Delta_{2i})^2} - \frac{c_{2i+1}}{B_{2i}/\Delta_{2i}} \quad (16.17)$$

where $c_{2i+1} \equiv -\text{sign}(m) \Delta_{2i}/A_{2i} \pmod{B_{2i}/\Delta_{2i}}$. Also note that

$$sT|B_{2i}/\Delta_{2i} \quad \text{and} \quad \left(\frac{B_{2i}}{\Delta_{2i}}, \frac{A_{2i}}{\Delta_{2i}} \right) = 1 \quad \text{implies} \quad (sT, c_{2i+1}) = 1.$$

Focusing just on the first term of (16.17),

$$\frac{m_{2i} r_{2i} (\omega_n + S_2/d) (d_z w_{2i} \bar{\Delta}_{2i})^2}{s^2 u^2 r |m| z^2 l_{2i} y_{2i}^2 T^{k(i+1)} U^{k(i+1)}} = \frac{r_{2i}}{r} \frac{m_{2i}}{|m|} \frac{(\omega_n + S_2/d)}{s^2 z_{2i+1}^2 u_{2i+1}^2 y_{2i+1}^2 l_{2i} (UT)^{k(i+1)}} \quad (16.18)$$

because $zu = \bar{\Delta}_{2i} z_{2i+1} u_{2i+1} d_z w_{2i}$. Furthermore, by taking:

$$r_{2i+1} := \frac{r}{r_{2i}}, \quad m_{2i+1} := \frac{|m|}{m_{2i}}, \quad l_{2i+1} := l_{2i},$$

(16.18) becomes,

$$\frac{(\omega_n + S_2/d)}{r_{2i+1}l_{2i+1}m_{2i+1}(su_{2i+1}y_{2i+1}z_{2i+1})^2(UT)^{k(i+1)}}.$$

Likewise, the second term is

$$\frac{c_{2i+1}w_{2i}d_z\bar{\Delta}_{2i}}{dsl_{2i}u_{2i}z_{2i}y_{2i}^2T^{k(i+1)}} = \frac{u}{u_{2i}} \frac{z}{z_{2i}} \frac{c_{2i+1}}{dsz_{2i+1}u_{2i+1}l_{2i}y_{2i}^2T^{k(i+1)}}$$

which if we take:

$$z'_{2i+1} := \frac{z}{z_{2i}}, \quad u'_{2i+1} := \frac{u}{u_{2i}},$$

becomes

$$\frac{u'_{2i+1}z'_{2i+1}c_{2i+1}}{dsl_{2i+1}u_{2i+1}y_{2i+1}^2z_{2i+1}T^{k(i+1)}}.$$

In summary, the entire element, $\theta_{h_{2i+1}}$ is

$$\theta_{h_{2i+1}} = \frac{\omega_n + S_2/d - sm_{2i+1}r_{2i+1}u_{2i+1}u'_{2i+1}z_{2i+1}z'_{2i+1}c_{2i+1}U^{k(i+1)}/d}{r_{2i+1}l_{2i+1}m_{2i+1}(su_{2i+1}z_{2i+1}y_{2i+1})^2(UT)^{k(i+1)}} \quad (16.19)$$

with:

$$\begin{aligned} m_{2i+1} &\in \{1, |m|\}, \quad l_{2i+1} \in \{1, |l|\}, \quad r_{2i+1} \in \{1, r\}, \\ u_{2i+1}|u, \quad u'_{2i+1}|u, \quad y_{2i+1}|y/d_y, \quad z_{2i+1}|z/d_z, \quad z'_{2i+1}|z, \\ (z_{2i+1}y_{2i+1}, u/u_{2i+1}) &= 1, \quad (c_{2i+1}, sT) = 1. \end{aligned}$$

Thus, from the complete quotient $\theta_{h_{2i-1}}$ we find another complete quotient $\theta_{h_{2i+1}}$ which satisfies exactly the same requirements as $\theta_{h_{2i-1}}$. Moreover, only a bounded number of the these complete quotients are not reduced. More precisely,

$$\begin{aligned} \frac{A_{2i}}{B_{2i}} &= \frac{k_1x^n - k_2U^{n-ki}T^{n+k}}{k_3T^{n+k(i+1)}U^{n-ki}} = \frac{k_1U^{ki}}{k_3T^{k(i+1)}} - \frac{k_2}{k_3T^{ki}} \\ &> k_4(U/T)^{ki} > 1 \end{aligned}$$

for some $i \geq W$, where W only depends on the parameters $m, l, y, z, r, d, c, N, U, T$.

Similarly,

$$\begin{aligned} \frac{A_{2i+1}}{B_{2i+1}} &= \frac{k_1x^n - k_2U^{k(i+1)}}{k_3(UT)^{k(i+1)}} = \frac{k_1x^n - k'_2U^{ki}}{k'_3(UT)^{ki}} = \frac{k_1x^{n-ki}}{k'_3} - \frac{k'_2}{k'_3T^{ki}} \\ &> k_4x^{n-ki} > 1 \end{aligned}$$

for some V such that $n - ki \geq V$, that is $n - V \geq ki$. And again, V depends only on the parameters $m, l, y, z, r, d, c, N, U, T$.

Because the values of the pairs (P_{h_i}, Q_{h_i}) are all distinct for $i = 1, 2, \dots, 2(n - V)/k + 1$ we see that

$$lp(\omega_n) > \sum_{j=W}^{(n-V)/k} p_{2j}.$$

Our interest now falls on the length of the expansion of A_{2i}/B_{2i} . Basically, since we have $\Theta(n)$ of these if the lengths are unbounded then, from above, the period length can not possibly be linear in n . In the next section we show that in order to have the length of A_{2i}/B_{2i} bounded for all i we must have $T = 1$.

17. THE LENGTH OF THE CONTINUED FRACTION EXPANSION OF A_{2i}/B_{2i}

In this section we recall that

$$d^2 D_n = (Ax^n + C)^2 + 4Gx^n = (2S_1 + td)^2 + 4Gx^n.$$

Let i be fixed with $W \leq i \leq (n - V)/k$ so that $\theta_{h_{2i}}$ is reduced. We have

$$\begin{aligned} P_{h_{2i}} &= S_1/d - Nsl_{2i}r_{2i}u_{2i}u'_{2i}y_{2i}y'_{2i}c_{2i}U^{n-ki}T^{n+k}/d \\ &= S_1/d - J/d. \end{aligned}$$

As usual, we have

$$d^2(t - D_n)/4 + d^2(tP_{h_{2i}} + P_{h_{2i}}^2) \equiv 0 \pmod{d^2Q_{h_{2i}}},$$

where in the above case

$$Q_{h_{2i}} = r_{2i}m_{2i}l_{2i}N(su_{2i}z_{2i}y_{2i})^2U^{n-ki}T^{n+k(i+1)}.$$

In other words,

$$-Gx^n - tdJ - 2S_1J + J^2 \equiv 0 \pmod{d^2Q_{h_{2i}}}.$$

Now $U^{n-ki}T^n$ divides every term on the left side so

$$\begin{aligned} GU^{ki} &\equiv -(td + 2S_1 - J)(Nsl_{2i}r_{2i}u_{2i}u'_{2i}y_{2i}y'_{2i}c_{2i}T^k) \\ &\quad \pmod{d^2Nl_{2i}r_{2i}m_{2i}(su_{2i}z_{2i}y_{2i})^2T^{k(i+1)}} \\ &\equiv -(Ax^n + C - J)(Nsl_{2i}r_{2i}u_{2i}u'_{2i}y_{2i}y'_{2i}c_{2i}T^k) \\ &\quad \pmod{d^2Nl_{2i}r_{2i}m_{2i}(su_{2i}z_{2i}y_{2i})^2T^{k(i+1)}}. \end{aligned}$$

Consequently, modulo $T^{k(i+1)}$ we have

$$GU^{ki} \equiv -CNsl_{2i}r_{2i}u_{2i}u'_{2i}y_{2i}y'_{2i}c_{2i}T^k \pmod{T^{k(i+1)}}. \quad (17.1)$$

Returning now to A_{2i}/B_{2i} , we previously found that

$$\frac{A_{2i}}{B_{2i}} = \frac{AU^{ki} - Nsr_{2i}l_{2i}u_{2i}u'_{2i}y_{2i}y'_{2i}c_{2i}T^k}{dNl_{2i}r_{2i}m_{2i}(su_{2i}y_{2i}z_{2i})^2T^{k(i+1)}}.$$

Writing (17.1) as

$$CNsl_{2i}r_{2i}u_{2i}u'_{2i}y_{2i}y'_{2i}c_{2i}T^k = -GU^{ki} - f_{2i}T^{k(i+1)}$$

where $f_{2i} \in \mathbb{Z}$, we find,

$$\begin{aligned} \frac{CA_{2i}}{CB_{2i}} &= \frac{CAU^{ki} - CNsr_{2i}l_{2i}u_{2i}u'_{2i}y_{2i}y'_{2i}c_{2i}T^k}{CNdl_{2i}r_{2i}m_{2i}(su_{2i}y_{2i}z_{2i})^2T^{k(i+1)}} \\ &= \frac{ACU^{ki} + GU^{ki} + f_{2i}T^{k(i+1)}}{CNdl_{2i}r_{2i}m_{2i}(su_{2i}y_{2i}z_{2i})^2T^{k(i+1)}}. \end{aligned} \quad (17.2)$$

From, $A = cqrN$ and $C = c(mz^2U^k - ly^2T^k)/q$ we have,

$$AC + G = c^2rNmz^2U^k$$

and (17.2) becomes

$$\begin{aligned} \frac{A_{2i}}{B_{2i}} &= \frac{c^2rNmz^2U^{k(i+1)} + f_{2i}T^{k(i+1)}}{CNdl_{2i}r_{2i}m_{2i}(su_{2i}y_{2i}z_{2i})^2T^{k(i+1)}} \\ &= \frac{1}{CNdl_{2i}r_{2i}m_{2i}(su_{2i}y_{2i}z_{2i})^2} \left[c^2rNmz^2 \left(\frac{U^k}{T^k} \right)^{i+1} + f_{2i} \right] \\ &= \frac{1}{F_{2i}} (E\xi^{i+1} + f_{2i}) \end{aligned}$$

where F_{2i} , E are bounded integers and $\xi = U^k/T^k$.

17.1. Depth of a sequence of rationals

We now provide an aside regarding the depth of $(a/b)^h$ as $h \rightarrow \infty$ for coprime a and b . Recall that $\delta(a)$ for $a \in \mathbb{Q}$ represents the depth of the regular continued fraction expansion of a .

Theorem 40. *If a and b are two coprime integers where $1 < b < a$ then*

$$\lim_{i \rightarrow \infty} \delta((a/b)^i) = \infty$$

Proof. Whether this is so was asked by Mendès France and proved by Pourchet in a letter to him. A summary of Pourchet's response is given by van der Poorten in [70].¹ ■

It is clear that $\delta(1/a) \geq \delta(a) - 2$, $\delta(-a) \geq \delta(a) - 2$ and $\delta(a+n) = \delta(a)$ for any $n \in \mathbb{Z}$. Furthermore, Mendès France [44] has shown that for any $a \in \mathbb{Q}$, $n \in \mathbb{Z}^+$,

$$\delta(na) \geq \frac{\delta(a) - 1}{\kappa(n) + 2} - 1$$

where $\kappa(n)$ is a positive valued function whose values depend only on n . Consequently, for any sequence $a_i \in \mathbb{Q}$ satisfying

$$\lim_{i \rightarrow \infty} \delta(a_i) = \infty$$

and any $n \in \mathbb{Z}^+$,

$$\lim_{i \rightarrow \infty} \delta\left(\frac{1}{a_i}\right) = \infty \implies \lim_{i \rightarrow \infty} \delta\left(\frac{n}{a_i}\right) = \infty \implies \lim_{i \rightarrow \infty} \delta\left(\frac{a_i}{n}\right) = \infty.$$

From the result above, if $(a, b) = 1$ and $1 < b < a$ then

$$\lim_{i \rightarrow \infty} \delta\left(\left(\frac{a}{b}\right)^i\right) = \infty.$$

Consequently, we find

$$\lim_{i \rightarrow \infty} \delta\left(\frac{A_{2i}}{B_{2i}}\right) = \lim_{i \rightarrow \infty} \delta\left(\frac{1}{F_{2i}}(E\xi^{i+1} + f_{2i})\right) = \infty \quad (17.3)$$

for $T > 1$.

17.2. Period length of ω_n

By our criteria for a kreeper we must have for some $a, b \in \mathbb{Q}$,

$$an + b = lp(\omega_n) > \sum_{j=W}^{(n-V)/k} p_{2j}$$

for all $n \in I$. Hence we must have

$$\sum_{i=W}^{(n-V)/k} \delta\left(\frac{A_{2i}}{B_{2i}}\right) < an + b. \quad (17.4)$$

By (17.3) there exists a $\gamma > W$ such that

$$\delta\left(\frac{A_{2i}}{B_{2i}}\right) > k(a+1)$$

¹Van der Poorten provides the following correction to the given argument: Consider $p_{n+1} < p_n a^{h\varepsilon_n}$ so that $a^h = p_{\psi(h)} < a^{h(\varepsilon_1 + \dots + \varepsilon_{\psi(h)})}$ and then consider $\varepsilon_1 + \dots + \varepsilon_{\psi(h)} = \psi(h)\varepsilon$.

for all $i \geq \gamma$. So,

$$\begin{aligned} \sum_{i=W}^{(n-V)/k} \delta\left(\frac{A_{2i}}{B_{2i}}\right) &\geq \sum_{j=\gamma}^{(n-V)/k} \delta\left(\frac{A_{2i}}{B_{2i}}\right) \\ &> k(a+1)[(n-V)/k - \gamma]. \end{aligned}$$

We now wish to establish an n so that this is larger than $an + b$, that is

$$k(a+1)\left(\frac{n-V}{k} - \gamma\right) > an + b.$$

This will certainly be the case when

$$n > b + (a+1)(V + k\gamma)$$

and since all the terms on the right side are bounded, there must exist an infinitude of values of $n \in I$ such that $lp(\omega_n) > an + b$ for any fixed a, b . In conclusion D_n can not be a keeper if $T > 1$. In other words, we must have $T = 1$ and $U = x$, so that a keeper must satisfy

$$D_n = \frac{c^2}{d^2} \left[(qrNx^n + (mz^2x^k - ly^2)/q)^2 + 4rNly^2x^n \right].$$

17.3. Showing that $N = 1$

Our only remaining objective is to show that necessarily $N = 1$. Clearly, there is no loss in generality in supposing that N is not a power of x . In §16 we established the existence of the following complete quotient in all keepers,

$$\theta_{h_{2i}} = \frac{\omega_n + S_1/d - Nsr_{2i}l_{2i}u_{2i}u'_{2i}y_{2i}y'_{2i}c_{2i}x^{n-ki}/d}{r_{2i}m_{2i}l_{2i}(su_{2i}y_{2i}z_{2i})^2Nx^{n-ki}}.$$

By taking $i = \lfloor n/k \rfloor$ we find an element, $\eta \in \mathcal{O}_n$, whose norm can be written as RNx^ν where $R \mid D_n$, $(N, E_n) = 1$ and $0 \leq \nu < k$. Hence the norm of η is bounded independently of n , and coprime to the conductor of the order $\mathcal{O}_n^*$. First, recall the ideal $\mathfrak{b}_n \in \mathcal{O}_n^*$ from page 69 which has norm x^g , with g fixed independently of n . Suppose that $N > 1$ and that the ideals (η) , $\mathfrak{b}_n$ are dependent in $\mathcal{O}_n^*$. Then by Theorem 26 we must have

$$N^e x^{\nu e} = x^{gf}$$

for some nonnegative integers e, f , not both zero. Since N is not a power of x , we must have $e = 0, f > 0$. But this would imply that $x = 1$ which is impossible. Hence (η) and $\mathfrak{b}_n$ are independent ideals in $\mathcal{O}_n^*$. Then by Proposition 32, $R(\mathcal{O}_n) \gg (\log D_n)^3$; that is the sequence of discriminants $\{D_n\}$ is not a keeper. Thus we must have $N = 1$.

In conclusion we have shown that a keeper must look like

$$D_n = \frac{c^2}{d^2} \left[(qrx^n + (mz^2x^k - ly^2)/q)^2 + 4rly^2x^n \right]$$

where m, l, r are squarefree and

$$rly^2mz^2 \mid d^2D_n \quad (rq, lymzx) = 1 \quad (ly, mz) = 1.$$

It remains to show that any sequence of discriminants which satisfy such a form are indeed keepers; we do this in Chapter VII. But before we do that, the next Chapter examines some other aspects of interest regarding keepers.

Chapter VI. Additional remarks about creepers

This chapter contains some extra thoughts about creepers. Since it is only intended to highlight various features, we will suppose throughout that c and d are equal to 1. These terms merely complicate the expansion without adding anything of interest. The full expansion in Chapter VII details the impact they have.

18. DON'T BE $\square$

This section examines the effect the terms y^2 and z^2 have upon creepers (in other words, D_n possesses square factors). The discriminants are no longer fundamental but this certainly doesn't prevent us from investigating their continued fraction expansions. (As is well known, multiplying a continued fraction expansion by an integer may furnish a shorter period length, which provides evidence for sometimes considering non-maximal orders).

The detailed expansion covered in Chapter VII allows the presence of rational squares in order to complete all the cases in one swell foop. But in practice one may multiply in any rationals they like and this raises the question as to which order is the simplest to consider. As we shall soon see, the correct approach is to always divide out the obvious square factors (y and z) but not any other squares. To demonstrate the hows and whys we consider the following example which highlights all the important details.

Example 9.

$$D_n = (2 \cdot 509 \cdot x^n + (3^2 11 \cdot x - 5^2 7)/2)^2 + 4 \cdot 509 \cdot (5^2 7) \cdot x^n \quad (x = 1319011)$$

The expansion commences as

$$\omega = S_1 + 1 - (\bar{\omega} + S_1)$$

$$\frac{\omega + S_1}{509(5^2 7)x^n} = \frac{2}{5^2 7} - \frac{\bar{\beta}}{509(5^2 7)x^n}$$

After the partial quotients arising from the expansion of $2/(5^2 7)$, we have the complete quotient,

$$\theta_{h_1} = \frac{\omega + S_2}{(5^2 7)(3^2 11)x} - \frac{c_1}{5^2 7} \quad \text{where} \quad c_1 \equiv -1/2 \pmod{5^2 7}.$$

The next line in the expansion is

$$\frac{2 \cdot 509 \cdot x^{n-1} - 87(3^2 11)}{(5^2 7)(3^2 11)} - \frac{\bar{\alpha}}{(5^2 7)(3^2 11)x}.$$

In parallel to the non-square case we would like to be able to say that $2 \cdot 509 \cdot x^{n-1} - 87(3^2 11) \equiv 0 \pmod{5^2 7}$, but unfortunately this may not be the case. For the above x and $n \equiv 4 \pmod{15}$ ¹, then,

$$(2 \cdot 509 \cdot x^{n-1} - 87(3^2 11), 5^2 7) = 5 \cdot 7$$

rather than $5^2 7$ as desired. The consequence is that a factor 5 from y will divide some Q_{h_i} 's where l does not. Specifically, the next complete quotient of interest is

$$\theta_{h_2} = \frac{(\omega + S_1)}{509 \cdot 5^2(3^2 11)x^{n-1}} - \frac{c_2}{5(3^2 11)},$$

$$\text{where} \quad c_2 \equiv -\frac{7}{(2 \cdot 509x^{n-1} - 87 \cdot 99)/5} \pmod{5 \cdot 3^2 11}.$$

¹since $x^{15} \equiv 1 \pmod{(5^2 7)(3^2 11)}$ we should consider n according to its residue modulo 15

The following line in the expansion is

$$\frac{2x - 172 \cdot 5}{5^2(3^211)} - \frac{\bar{\beta}}{509 \cdot 5^2(3^211)x^{n-1}}.$$

Fortunately, $2x - 172 \cdot 5 \equiv 0 \pmod{3^211}$ (all that we can guarantee is that it is 0 modulo $3 \cdot 11$), and after the expansion of $(2x - 172 \cdot 5)/(5^23^211)$ we have the complete quotient

$$\theta_{h_3} = \frac{(\omega + S_2)}{5^2x^2} - \frac{c_3}{5^2} \quad \text{where} \quad c_3 \equiv -\frac{3^211}{2x - 172 \cdot 5} \pmod{5^2}.$$

From this, we find that the next line is

$$\frac{2 \cdot 509 \cdot x^{n-2} - 23}{5^2} - \frac{\bar{\alpha}}{5^2x^2}.$$

Now $5 \nmid 2 \cdot 509 \cdot x^{n-2} - 23$, so we find

$$\theta_{h_4} = \frac{\omega + S_1}{509(5^27)x^{n-2}} - \frac{c_4}{5} \quad \text{where} \quad c_4 \equiv \frac{-1}{(2 \cdot 509x^{n-2} - 23)/5} \pmod{5}.$$

The point here is that the square factors y^2 and z^2 manifest themselves in unusual places. More importantly, unless one sits down and physically computes the particular case in question there is seemingly no way in which the occurrence of these square factors can be easily predicted. To illustrate this the following table lists some factorizations of Q_{h_i} in the case of $n = 19$.

h	Factors other than x	Exponent of x
2	$(3^211)(5^27)$	1
9	$5^2(3^211)509$	18
14	5^2	2
17	$(5^27)509$	17
166	$(3^211)(5^27)$	17
$\vdots$		
173	$(3^211)509$	2
174	1	18
175	$(5^27)509$	1
185	$5^2(3^211)509$	0
193	$(5^27)509$	1
196	5^2	18
203	$5^2(3^211)509$	2
208	$(3^211)(5^27)$	17
$\vdots$		
321	$(5^27)509$	17
326	5^2	2
331	$5^2(3^211)509$	18
336	$(3^211)(5^27)$	1
340	5^2	0
342	$(3^211)(5^27)$	1
347	$(3^211)509$	18
348	1	2
349	$(5^27)509$	17
$\vdots$		
508	$(3^211)(5^27)$	17
519	$5^2(3^211)509$	2

522	5^2	18
527	$(5^2 7)509$	1
539	$(3^2 11)509$	0
551	$(5^2 7)509$	1
556	5^2	18
559	$5^2(3^2 11)509$	2
570	$(3^2 11)(5^2 7)$	17
$\vdots$		
729	$(5^2 7)509$	17
730	1	2
731	$(3^2 11)509$	18
736	$(3^2 11)(5^2 7)$	1
738	5^2	0
742	$(3^2 11)(5^2 7)$	1
747	$5^2(3^2 11)509$	18
752	5^2	2
757	$(5^2 7)509$	17
$\vdots$		
870	$(3^2 11)(5^2 7)$	17
875	$5^2(3^2 11)509$	2
882	5^2	18
885	$(5^2 7)509$	1
893	$5^2(3^2 11)509$	0
903	$(5^2 7)509$	1
904	1	18
905	$(3^2 11)509$	2
912	$(3^2 11)(5^2 7)$	17
$\vdots$		
1061	$(5^2 7)509$	17
1064	5^2	2
1069	$5^2(3^2 11)509$	18
1076	$(3^2 11)(5^2 7)$	1
1078	1	0

Notice that because of the square factors, the points which would normally be halfway, that is exponent of x is zero, are no longer guaranteed of being halfway. For this congruence class, the period length of ω_n is given by

$$lp(\omega_n) = \begin{cases} \frac{862}{15}n - \frac{478}{15} & \text{if } n \equiv 4 \pmod{30} \\ \frac{862}{15}n - \frac{208}{15} & \text{if } n \equiv 19 \pmod{30} \end{cases}$$

But it is clear that the above table is merely a demonstration of my ramblings in §3 regarding units in different orders. In other words, if $D' = D/(3^2 5^2)$ then the fundamental unit of the ring of integers of $\mathbb{Q}(\sqrt{D})$ is some power of the fundamental unit of the ring of integers of $\mathbb{Q}(\sqrt{D'})$ ¹. Moreover, one might notice that, in the instance above the extra factor floating about is always 5^2 and never 3^2 and that the expansion appears to be three segments of a shorter expansion.

¹here, unlike other sections, we are not supposing that D' is squarefree

Some other interesting cases include:

$$n \equiv 9 \pmod{30} \quad lp(\omega) = \frac{392}{3}n - 56,$$

$$n \equiv 21 \pmod{30} \quad lp(\omega) = \frac{392}{3}n - 48.$$

Both of these cases are examples needing six segments. Furthermore, since six is fairly big, we would expect that for $D' = D/(3^2 5^2)$ we should have

$$\left(\frac{D'}{3}\right) = \left(\frac{D'}{5}\right) = -1$$

and indeed this is the case. On the short side of things,

$$n \equiv 7 \pmod{30} \quad lp(\omega) = \frac{1}{3}(53n + 19)$$

which doesn't require any extra segments; as expected,

$$\left(\frac{D'}{3}\right) = \left(\frac{D'}{5}\right) = 1$$

One might also be inclined to look at

$$D_n = (31 \cdot 1306^n + 9 \cdot 1306^k - 5)^2 + 4 \cdot 31 \cdot 9 \cdot 1306^n$$

where it is necessary to take $k \equiv 1 \pmod{5}$. This also demonstrates that one might consider a keeper as a function of three variables, n , k and x . It seems this viewpoint is not particularly helpful.

19. THE PROPER WAY TO HANDLE SQUARE FACTORS

The previous section attempts to provide a method for handling square factors. Indeed by using such an approach it is not difficult at all to obtain the continued fraction expansion of such a keeper. However, the splitting into residue classes of n is unfortunate and unavoidable, under such an approach. Instead we consider an alternative where this is not necessary. Among the advantages of this approach are: only one "formula" needs to be given, period lengths match up, and the period lengths are shorter (all evidence for suggesting this is *the* way to handle square factors). The suggestion is that decency requires first removing obvious square factors from D_n .

The main difficulty is that we no longer possess an integer representation of our discriminant. This is not a serious obstacle, it just means that we can never "reciprocate" easily; Lemma 2 is required at *every* step of the continued fraction expansion.

Our keeper looks like

$$D_n = (grx^n + (mz^2x^k - ly^2)/q)^2 + 4ly^2rx^n$$

where l , m are squarefree. Instead of taking $S_1 = (s_1 - t)/2$ we are now going to consider

$$S_1 = \frac{s_1 - tzy}{2} \quad \text{and} \quad S_2 = \frac{s_2 - tzy}{2}$$

where $D_n/(zy)^2 \equiv t \pmod{4}$. However, t might no longer be 0 or 1; if so, we replace D_n by $4D_n$. The point of this is that we are interested in

$$\alpha = \omega + S_1/zy = \frac{z\omega + S_y}{z}$$

and

$$\beta = \omega + S_2/zy = \frac{y\omega + S_z}{y}$$

where S_y and S_z denote S_1/y and S_2/z respectively (both of which are integers by the division requirements).

As before, we need to know the quantities,

$$\begin{aligned} \alpha\bar{\alpha} &= \frac{(zy\bar{\omega} + S_1)(zy\omega + S_1)}{z^2y^2} = \frac{z^2y^2\omega\bar{\omega} + zyS_1(\omega + \bar{\omega}) + S_1^2}{z^2y^2} \\ &= -\frac{lr x^n}{z^2}, \end{aligned}$$

and

$$\begin{aligned} \beta\bar{\beta} &= \frac{(zy\bar{\omega} + S_2)(zy\omega + S_2)}{z^2y^2} = \frac{z^2y^2\omega\bar{\omega} + S_2zy(\omega + \bar{\omega}) + S_2^2}{z^2y^2} \\ &= \frac{-rmx^{n+k}}{y^2}, \end{aligned}$$

neither of which is integral. For the sake of brevity we are not going to concern ourselves about the signs of Q_{h_i} which we know can easily be handled. Anyway, the expansion of $\omega = (t + \sqrt{D_n/zy})/2$ commences as

$$\omega_n = \frac{S_y}{z} + t - \bar{\alpha}.$$

Using Lemma 2 we discover that the next partial quotients are the expansion of $(S_y + tz)/z$ followed by the complete quotient

$$\frac{\omega_n + S_y/z}{lr x^n} - \frac{c_0}{z}, \quad c_0 \equiv -1/S_y \pmod{z}.$$

The next line in the expansion is

$$\frac{q - c_0ly}{lzy} - \frac{\bar{\beta}}{lr x^n}.$$

Obviously $q - c_0ly \not\equiv 0 \pmod{ly}$. However it is the case that $q - c_0ly \equiv 0 \pmod{z}$ (this is similar to the cases for squarefree l, m where certain divisions automatically hold).

$$\begin{aligned} q - c_0ly &\equiv q + lyS_y^{-1} \equiv S_y^{-1}(qS_1 + ly^2)y^{-1} \pmod{z} \\ &\equiv C(q^2rx^n - ly^2 + 2ly^2) \pmod{z} \quad \text{for some } C \neq 0 \\ &\equiv Cqs_2 \pmod{z}. \end{aligned}$$

The next partial quotients arise from the expansion of $\frac{(q - c_0ly)/z}{ly}$ followed by

$$\frac{\omega_n + S_z/y}{lmx^k} - \frac{c_1}{ly}, \quad c_1 \equiv -z/q \pmod{ly}.$$

And from this we find that the next line is

$$\frac{qrx^{n-k} - c_1zm}{yzlm} - \frac{\bar{\alpha}}{lmx^k}.$$

We now note that

$$\begin{aligned} qrx^{n-k} - c_1zm &\equiv (q^2rx^{n-k} + z^2m)/z \pmod{ly} \\ &\equiv S_1/z \pmod{ly} \end{aligned}$$

Thus, the next partial quotients are the expansion of $\frac{(qrx^{n-k} - c_1zm)/ly}{mz}$ followed by the complete quotient

$$\frac{\omega + S_y/z}{rmx^{n-k}} - \frac{c_2}{mz}, \quad c_2 \equiv \frac{-ly}{qrx^{n-k}} \pmod{mz}.$$

The next line is

$$\frac{qx^k - c_2y}{mzy} - \frac{\bar{\beta}}{rmx^{n-k}}.$$

And so on ...

The point is that the square factors of D_n , namely z^2 and y^2 , no longer pollute any of the Q_{h_i} 's. The catch is that while it is easy to show

$$lp\left(\sqrt{D_n}/\kappa\right) = an + b \text{ implies that } lp\left(\sqrt{D_n}\right) = O(n)$$

it is not immediately obvious that the stronger

$$lp\left(\sqrt{D_n}/\kappa\right) = an + b \text{ implies that } lp\left(\sqrt{D_n}\right) = a'n + b' \quad (19.1)$$

holds. In the case of keepers we physically demonstrate this in Chapter VII. However, it is an interesting question as to whether or not one can prove that (19.1) holds for any sequence of discriminants without prior knowledge of the actual expansion. The R - L sequences of Raney and the phrase “finite-state transducer” suggest that this is conceivable, but an immediate proof eludes us. Anyway, this also demonstrates that while the period length being in an arithmetic progression is exciting, the real focus should be on regulators in which case the problem of multiplying by rationals becomes trivial (recall §3, page 13, regarding powers of fundamental units).

Returning to our earlier example of

$$D_n = (2 \cdot 509 \cdot 1319011^n + (3^2 \cdot 11 \cdot 1319011 - 5^2 \cdot 7)/2)^2 + 4 \cdot 509 \cdot 5^2 \cdot 7 \cdot 1319011^n,$$

enough has been said to make it plain that the expansion of $D_n/(3 \cdot 5)^2$ is

$$\omega_n = \left[\frac{S_y - 2}{3} + 1, 1, 2, -1, 1, 2, 5, 2, \frac{(2 \cdot 509x^{n-1} - 16 \cdot 3 \cdot 11)/5 \cdot 7 - 14}{3 \cdot 11}, \right. \\ \left. 2, 2, 1, 4, \frac{(2x - 5 \cdot 7)/3 \cdot 11}{5}, 1, 4, \frac{(2 \cdot 509x^{n-2} - 3)/5 - 2}{3}, 1, 2, \right. \\ \left. \frac{2x^2 - 5 \cdot 7}{3 \cdot 5 \cdot 7}, 1, 2, 5, 2, \frac{(2 \cdot 509x^{n-3} - 16 \cdot 3 \cdot 11)/5 \cdot 7 - 14}{3 \cdot 11}, 2, 2, \dots \right]$$

where the pattern is now evident. When $x \equiv 1 \pmod{ml}$ then by considering $D_n/(zy)^2$ instead of D_n , the period lengths of ω won't split into separate congruence classes. For the above example we have that $lp(\omega) = 16n - 6$, a far cleaner formula than the ones for D_n on page 90.

An explicit expansion of this example can be found in the Tables, at page 145.

20. A COMPOSITION APPROACH

In this section we demonstrate that it is possible to find a unit purely from ideal compositions. As stated earlier, one detects a keeper by finding a suitable pair of ideals and composing them appropriately until a unit is formed. A minor problem is that one is not automatically guaranteed that the unit found is indeed fundamental (but with a little thought one can recover from this difficulty). Since this section has nothing to do with period lengths, for ease of exposition, we will suppose that y and z are equal to 1.

The starting point is the simple, yet essential, observation that D_n can be represented as

$$D_n = s_1^2 + 4lr^n \quad (20.1)$$

as well as by

$$D_n = s_2^2 + 4rmx^{n+k}. \quad (20.2)$$

From (20.1) we have the ideal,

$$\mathfrak{a} = \langle rlx^n, \omega + S_1 \rangle.$$

Equally, (20.2), produces an ideal

$$\mathfrak{b} = \langle rmx^{n+k}, \omega + S_2 \rangle.$$

The observant reader will realize that $\mathfrak{b}$ is unreduced and consequently does not exist in the primitive, reduced cycle (in fact $\mathfrak{a}$ might also be unreduced). But this is not a problem because all we want to do is construct the unit by composition.

Crudely composing with no regard for the P_h 's yields,

$$\begin{aligned} \mathfrak{a}\mathfrak{b} &= \langle mlx^k, \omega + P_1 \rangle && \text{provided} && \text{(i) } r|S_1 + S_2 + t \\ \mathfrak{a}^2\mathfrak{b} &= \langle rmx^{n-k}, \omega + P_2 \rangle && \text{provided} && \text{(ii) } l|S_1 + P_1 + t \\ \mathfrak{a}^3\mathfrak{b}^2 &= \langle rlx^{n-2k}, \omega + P_3 \rangle && \text{provided} && \text{(iii) } m|P_1 + P_2 + t \end{aligned}$$

The ideal $\mathfrak{a}^2\mathfrak{b}^2 = \langle x^{2k}, \omega + p \rangle$ can also be created but the unit can be found without doing that.

The requirements stated on the right are the situations which need to be fulfilled so that constants which are common to both norms involved in the composition are removed from the composite. For example, in $\mathfrak{a}$ and $\mathfrak{a}\mathfrak{b}$ the only common constant factor (excluding x of course) between the norms rlx^n and mlx^k is l . If it were the case that $l \nmid S_1 + P_1 + t$ then the norm of $\mathfrak{a}^2\mathfrak{b}$ would contain l^2 . Evidently, it would not be possible to form any new composites without some power of l dividing its norm, and hence we could not easily obtain a unit.

It is immediate that if r, l and m divide D_n then because of (20.1) and (20.2) each constant will divide its corresponding s_i (that is $l|s_1, m|s_2$ and consequently any future ideals will behave equally nicely. Conveniently, the requirements (ii)-(iii) are equivalent to l and m dividing D_n .

Condition (ii): $l|S_1 + P_1 + t$. What is P_1 ? The rules of composition determine that P_1 has to satisfy (amongst other things)

$$P_1 \equiv S_1 \pmod{l} \quad (20.3)$$

$$P_1 \equiv S_2 \pmod{mx^k} \quad (20.4)$$

which says that modulo l , P_1 is S_1 . That is, $S_1 + P_1 + t$ modulo l is equal to $2S_1 + t$ modulo l which of course is equal to s_1 modulo l . Hence, (ii) is satisfied if and only if $l|s_1$.

Condition (iii): $m|P_1 + P_2 + t$. This time we have that P_2 must satisfy,

$$P_2 \equiv P_1 \pmod{m} \quad (20.5)$$

From (20.4) we have $P_1 \equiv S_2 \pmod{m}$. In conjunction with (20.5) this implies that $P_2 \equiv S_2 \pmod{m}$. The condition $P_1 + P_2 + t \equiv 0 \pmod{m}$ then becomes $2S_2 + t \equiv 0 \pmod{m}$, which in other words is $m|s_2$.

20.1. Remarks about the term called r

Note that condition (i) is trivial. Indeed, r acts very differently from l and m . Whereas the powers of l and m dividing each norm would increase if the above divisibility conditions failed to hold, this will not happen for r . Regardless of the value of r , we would find either $\langle rl, \omega + p \rangle$ or $\langle rm, \omega + p' \rangle$. The point is that if $r \nmid D_n$ then the square of this ideal is not $\langle 1, \omega + P \rangle$. One could take $\Theta(n)$ powers of this ideal; in which case $R(D_n) \gg n^3$, going against creeper regulations. In fact this is precisely what Yamamoto does in order to find $R(D_n) \gg n^3$ infinitely often for a parametrized family D_n . (Actually, this is also exactly the situation described in Chapter V when we considered $N > 1$ on page 86). The problem then is that one has no idea about the unit (or the period length). For example,

$$D_n = (3 \cdot 2^n + 2 - 1)^2 + 4 \cdot 3 \cdot 2^n \quad \text{and} \quad E_n = (5 \cdot 2^n + 2 - 1)^2 + 4 \cdot 5 \cdot 2^n$$

have the period lengths,

n	$lp(\omega_n)$		n	$lp(\omega_n)$
5	128		5	61
6	188		6	238
7	564		7	458
8	838		8	702
9	1874		9	1168
10	559		10	714
11	8972	and	11	726
12	1210		12	4474
13	21674		13	27524
14	48574		14	55583
15	3660		15	33236
16	148955		16	252814
17	2926		17	34250
18	237644		18	248296

The expectation is that these discriminants should behave like a “random” discriminant. Indeed, $h(D_{16}) = 1$ showing that these period lengths are as long as can be expected. A question is whether D_{17} is a natural aberration or part of something more sinister, noting that $h(D_{17}) = 92$ is mildly large¹.

Beyond the element, $(\omega + P_h)/r^v$ where $2^n/r < r^v < 2^n$, our current methods cease to function; that is we can't write down a reduced principal ideal for each n , hence the difficulty in predicting the period length or finding a unit.

20.2. Sequences of ideals

Returning now to the sequence of ideals formed from composing $\mathfrak{a}$ and $\mathfrak{b}$. Since $\mathfrak{a}^3\mathfrak{b}^2 = \langle rlx^{n-2k}, \omega + S_1 \rangle$ we can inductively apply the preceding procedure to obtain

$$\begin{aligned}
\mathfrak{a}^3\mathfrak{b}^3 &= \langle mlx^{3k}, \omega + P_4 \rangle \\
\mathfrak{a}^4\mathfrak{b}^3 &= \langle rmx^{n-3k}, \omega + P_5 \rangle \\
\mathfrak{a}^5\mathfrak{b}^4 &= \langle rlx^{n-4k}, \omega + S_1 \rangle \\
&\vdots \\
\mathfrak{a}^{j+1}\mathfrak{b}^j &= \langle rlx^{n-jk}, \omega + S_1 \rangle
\end{aligned}$$

¹Even more perversely, $(43 \cdot 2^{18} + 1)^2 + 4 \cdot 43 \cdot 2^{18}$ has a class number of 204

where $j = \lfloor n/k \rfloor$ (in the above case it is supposed that j is even, when j is odd the final ideal produced would be $\langle rm x^{n-jk}, \omega + P \rangle$)

Remark. In the situation where $k \nmid n$ this process terminates with an ideal $\langle rm, \omega + P \rangle$ or $\langle rl, \omega + P \rangle$ when n/k is odd or even respectively. Furthermore, since $rl \mid D_n$ or $rm \mid D_n$ the square of this ideal has norm 1.

As is clearly apparent, if $k \nmid n$ then we are not going to end happily with the above process. The two different cases which can arise are:

1. $mlx^{jk} > x^n$ and $rmx^{n-jk} < 1$,
2. $x^{(j+1)k} > x^n$ and $rlx^{n-(j+1)k} < 1$.

Naturally, the same technique applies to both of them, hence we shall only consider the first case. In other words, let us suppose that we arrive at the ideal $\mathfrak{r} = \langle rm x^\nu, \omega + P \rangle$ where $0 < \nu < k$ and $P \equiv S_2 \pmod{mx^\nu}$. We now compose $\mathfrak{r}$ with $\mathfrak{ab}$. Since $\nu < k$ we have $G = \gcd(rmx^\nu, mlx^k, P_1 + P + t) = mx^\nu$. Hence the composite is given by $\mathfrak{c} = \langle rlx^{k-\nu}, \omega + p \rangle$ where $p \equiv P_1 \pmod{lx^{k-\nu}}$ and $p \equiv S_2 \pmod{m}$.

And we can continue composing as before. The sequence of ideals now runs in a reverse pattern to before. That is, we now find the following sequence of ideals,

$$\langle x^{n-\nu}, \omega + S_1 \rangle, \langle rm x^{\nu+k}, \omega + S_2 \rangle, \langle mlx^{n-\nu-k}, \omega + P' \rangle, \langle rlx^{\nu+2k}, \omega + P'' \rangle.$$

This procedure continues to repeat until we find an ideal whose norm has an exponent of 0 for x , and the square of this ideal has norm 1. Moreover, by backtracking and counting the compositions, we find that this ideal will be $\mathfrak{a}^{2n+2k}\mathfrak{b}^{2n}$, unless rl or rm is 1, in which case we will not have needed to take the square.

Of course, this composition is nothing other than an application of the continued fraction method in disguise. When expanding ω_n we “jumped” between complete quotients $(\omega + P_{h_i})/Q_{h_i}$ and $(\omega + P_{h_{i+1}})/Q_{h_{i+1}}$ and this composition is exactly the same procedure. One could also detail the expansion using R - L sequences, much as was done in [72].

In Chapter VII we show that the period length of (20.1) is given by $lp(\omega_n) = an + b$. We suggest to the reader that by appropriately using the notion of distance and infrastructure, it is possible to reach the same conclusion from this composition approach.

21. THE FUNDAMENTAL UNIT OF KREEPERS

Any creeper can be represented, up to square factors, as

$$D_n = (qrx^n + (mz^2x^k - ly^2)/q)^2 + 4lrx^n$$

where the usual conditions apply for q, r, l, m, z, y . But on the other hand, such an equation itself represents infinitely many keepers by appropriately Chinese Remaindering the division requirements in order to find other suitable x 's. For example, suppose that we have decided that we want $r = 3, m = 7, l = 11$ and $q = k = 1$. Then it only remains to determine some x and n . The required division conditions are:

$$x^n \equiv 1 \pmod{7}, \quad 3x^n + 7x \equiv 0 \pmod{11}, \quad x \equiv 2 \pmod{3}.$$

There are several possible solutions. For example, we could consider either

$$x = 71 \text{ with } n \equiv 7 \pmod{10} \quad \text{or} \quad x = 2 \text{ with } n \equiv 15 \pmod{30}.$$

Whatever, it is immediate that once we have a creeper then another creeper can be formed by considering $x' \equiv x \pmod{qrlmzy}$. Obviously, all the division requirements will then hold for x' . However, the original choices of r, ly and mz can not be completely arbitrary since there are selections for which no keepers exist.

In other words, the use of x rather than some other letter is appropriate in that this is a parametrization. But once this is realized, it cries out to us consider such an equation as a function of x and examine the corresponding function fields. In order to avoid any possible confusion we replace x by X to indicate our desire now to consider X as an indeterminate (recall our convention that an x means that we are interested in the numerical not the functional case).

Since we are now in the function field setting the issue of nontrivial square factors needs to be considered. In our cases such occurrences are uncommon, but they do exist, for example when $r = z = y = 1$, $l = -1$, $m = 4$, $D_n(X)$ has the nontrivial square factor of $(X + 1)^2$ when $n = 2$. However, since periodicity is never assumed but rather derived, it is perfectly allowable to ignore any such factors.

In this section, again it will be convenient to have $(n, k) = 1$, thus we replace X by $X^{(n,k)}$ if necessary. Since rational are units in $\mathbb{Q}[X]$, multiplying our discriminant by a rational is harmless, that is the period length is preserved. Multiplying through by appropriate rationals, our preferred form is

$$Y_n^2 = ((q^2 r X^n + m z^2 X^k + l y^2)/2)^2 - m z^2 l y^2 X^k$$

where as before r, l, m, z, y are integers. The following notations will prove useful,

$$\begin{aligned} S_1 &= (q^2 r X^n + m z^2 X^k - l y^2)/2, & \alpha &= Y_n + S_1, \\ S_2 &= (q^2 r X^n - m z^2 X^k + l y^2)/2, & \beta &= Y_n + S_2, \\ S_3 &= (q^2 r X^n + m z^2 X^k + l y^2)/2, & \gamma &= Y_n + S_3. \end{aligned}$$

So that

$$\alpha \bar{\alpha} = -r l y^2 X^n, \quad \beta \bar{\beta} = -r m z^2 X^{n+k}, \quad \gamma \bar{\gamma} = m z^2 l y^2 X^k.$$

The continued fraction expansion of Y commences as

$$\begin{aligned} Y_n &= S_3 - \bar{\gamma} \\ \frac{\gamma}{-m z^2 l y^2 X^k} &= \frac{-(q^2 r X^{n-k} + m z^2)}{m z^2 l y^2} - \frac{\bar{\alpha}}{-m z^2 l y^2 X^k} \\ \frac{\alpha}{-r X^{n-k}/m z^2} &= -m z^2 q^2 X^k - \frac{\bar{\beta}}{-r X^{n-k}/m z^2} \\ \frac{\beta}{-(m z^2)^2 X^{2k}} &= \frac{-q^2 r X^{n-2k}}{(m z^2)^2} - \frac{\bar{\alpha}}{-(m z^2)^2 X^{2k}} \\ \frac{\alpha}{-r l y^2 X^{n-2k}/(m z^2)^2} &= \dots \end{aligned}$$

And at this point we have

$$\theta_2 = \frac{\alpha}{-r X^{n-k}/m z^2} \quad \text{and} \quad \theta_4 = \frac{\alpha}{-r l y^2 X^{n-2k}/(m z^2)^2} = \theta_2 \left(\frac{m z^2 X^k}{l y^2} \right)$$

Hence, inductively we find that

$$\theta_{2(i+1)} = \theta_2 \left(\frac{m z^2 X^k}{l y^2} \right)^i$$

provided $n - k(i + 1) \geq k$.

Recall the earlier notation,

$$\begin{aligned} \lambda_1 &:= k \\ \lambda_{i+2} &:= \begin{cases} \lambda_i + k - n \varepsilon_i & \text{if } i \equiv 1 \pmod{2} \\ 0 & \text{if } i \equiv 0 \pmod{2} \end{cases} \end{aligned}$$

and

$$\varepsilon_i = \begin{cases} 1 & \text{if } \lambda_i \geq n - k \\ 0 & \text{if } \lambda_i < n - k \text{ or } 2|i. \end{cases}$$

Also note that, $\lambda_{2i-1} \equiv ki \pmod{n}$ so that $\lambda_{2n-1} = 0$.

Then the elements so far appearing in the continued fraction expansion are

$$\frac{\beta}{-(mz^2)^{(i+1)/2} X^{\lambda_i} / (ly^2)^{(i-3)/2}} \quad \text{and} \quad \frac{\alpha}{-r(ly^2)^{i/2-1} X^{n-\lambda_i} / (mz^2)^{i/2}}$$

for i odd and even respectively.

The points $\varepsilon_j = 1$ correspond to a place in the expansion where the above induction is not valid. Let us now suppose that we are in such a situation, that is $\lambda_i \geq n - k$, then we find

$$\begin{aligned} \frac{\alpha}{-r(ly^2)^{i/2-1} X^{n-\lambda_i} / (mz^2)^{i/2}} &= \frac{-(mz^2)^{i/2} (q^2 r X^{\lambda_i} + mz^2 X^{k-n+\lambda_i})}{qr(ly^2)^{i/2-1}} - \frac{\bar{\gamma}}{-r(ly^2)^{i/2-1} X^{n-\lambda_i} / (mz^2)^{i/2}} \\ \frac{\gamma}{(mz^2)^{i/2+1} X^{k+\lambda_i-n} / r(ly^2)^{i/2-2}} &= \frac{(ly^2)^{i/2-2} q^2 X^{2n-k-\lambda_i} + mz^2 X^{n-\lambda_i}}{q(mz^2)^{i/2+1}} - \frac{\bar{\alpha}}{(mz^2)^{i/2+1} X^{k+\lambda_i-n} / r(ly^2)^{i/2-2}} \end{aligned}$$

and since $\lambda_{i+2} = \lambda_i + k - n$ in this case, the next complete quotient and line in the expansion are:

$$\begin{aligned} \theta_{i+2} &= \frac{\alpha}{r^2(ly^2)^{i/2-1} X^{n-\lambda_{i+2}} / (mz^2)^{i/2+1}} = \\ &= \frac{(mz^2)^{i/2+1} q X^{\lambda_{i+2}}}{r(ly^2)^{i/2-1}} - \frac{\bar{\beta}}{r^2(ly^2)^{i/2-1} X^{n-\lambda_{i+2}} / (mz^2)^{i/2+1}}. \end{aligned}$$

And from here, we may apply the inductive process again.

Regarding the ε_j 's,

$$\varepsilon_{2j-1} = \left\lfloor \frac{(j+1)k}{n} \right\rfloor - \left\lfloor \frac{jk}{n} \right\rfloor$$

and

$$\sum_{j=0}^{2ns-1} \varepsilon_j = \sum_{j=0}^{ns} \varepsilon_{2j-1} = \left\lfloor \frac{(ns+1)k}{n} \right\rfloor = sk$$

hence there are precisely sk values of $i \in \{1, \dots, 2ns-1\}$ where $\varepsilon_j = 1$.

Clearly, in the primitive period there can't be more than k instances of $\varepsilon_j = 1$, so that we may consider $s = 1$. We know that $\lambda_{2n-3} = n - k$ which implies that $n - \lambda_{2n-3} = k$, and we must have

$$\begin{aligned} \theta_{2n-2} &= \frac{\alpha}{(-1)^k r^k (ly^2)^{n-k-1} X^k / (mz^2)^{n-1}} = \\ &= \frac{(-1)^k (mz^2)^{n-1} (q^2 r X^{n-k} + mz^2)}{r^k (ly^2)^{n-k-1}} - \frac{\bar{\gamma}}{(-1)^k r^k (ly^2)^{n-k-1} X^k / (mz^2)^{n-1}} \\ \theta_{2n-1} &= \frac{\gamma}{(mz^2)^n / r^k (ly^2)^{n-k-2}} = \dots \end{aligned}$$

which is the end of the quasi-period. We want to find a numerical unit so instead of finding the function field unit by multiplying all the elements in a quasi-period we will multiply all the elements in the full period. Recalling the expansion, we know that since every even line was of the shape α/Q , we have that $Q_i Q_{i+1} = ry^2 l X^n$

for every odd i . Furthermore, in the quasi-period there are $n - 1$ occurrences of α as a numerator, $n - k - 1$ times of β and $k + 1$ times of γ . Altogether we find,

$$\varepsilon_n = \frac{\alpha^{2(n-1)} \beta^{2(n-k-1)} \gamma^{2(k+1)}}{(r(l y^2) X^n)^{2(n-1)} \frac{(m z^2)^n}{r^k (l y^2)^{n-k-2}}} = \frac{\alpha^{2(n-1)} \beta^{2(n-k-1)} \gamma^{2(k+1)}}{r^{2n-k-2} (l y^2)^{n+k} (m z^2)^n X^{2n(n-1)}}$$

As for the integrality of this unit, clearly $X^{2n(n-1)}$ divides the numerator since by construction $\varepsilon = T(X) + U(X)Y_n$ where $T(X)$ and $U(X)$ are polynomials. Now, $\alpha^2 = Y_n^2 + S_1^2 + 2S_1 Y_n$, of which each term is divisible by rly^2 . Similarly, for β , γ and rmz^2 , meaning that this unit is an element of $\mathcal{O}_K$. Alternatively, one might note that any creeper with n fixed yields a sleeper and thus the unit must be integral by Schinzel's condition.

It may be noted that in this situation, since r , ly^2 and mz^2 are units then we are essentially retreating the earlier known cases of keepers but in the function field setting rather than the numerical.

22. FUNCTION FIELD KREEPERS

The previous section considered the case of taking a numerical creeper and considering the corresponding function fields. But this is a simplistic view of the matter. As was remarked at the conclusion of the section, we basically redid the work of earlier authors but now in the function field setting. This immediately provokes the question as to whether the extensions to keepers as demonstrated throughout here (that of allowing r , l and m to be non-units) can also be dealt with in the function field case. It can be.

Our principle interest is unashamedly numerical and so our description of function field keepers is rather scant. But it is evident that if we work over the field $\mathbb{F}_q[X]$ then all the remarks made previously which determine keepers over the rationals can be translated into the case $\mathbb{F}_q[X]$. The situation where the function is in $\mathbb{Q}[X]$ is only slightly trickier. In this case there are infinitely many functions of any given degree which prohibits the bounding of the sets which contain the partial quotients of bounded degree. But since this function must be arithmetic in period length over $\mathbb{F}_p[X]$ for each prime, p , it is straightforward that over $\mathbb{Q}[X]$ it must again be in an arithmetic progression (just apply the reduction principle of Yu and Van der Poorten as described in §5).

Finding keepers over $\mathbb{Q}[X]$ is considerably harder than finding keepers over $\mathbb{Z}$ (for brevity we suppose that $z = y = 1$). Consider the condition,

$$q^2 r X^n + l \equiv 0 \pmod{m} \quad q, r, l, m \in \mathbb{Z}[X]$$

If $m_i, \dots, m_g$ represent all the complex zeros of m then

$$q^2(m_i) r(m_i) m_i^n + l(m_i) = 0$$

for every $i = 1, \dots, g$. But this implies,

$$m_i^n = \frac{-l(m_i)}{q^2(m_i) r(m_i)}$$

which can only be true as $n \rightarrow \infty$ if $|m_i| = 1$ or 0 . But $(m, X) = 1$ implies $|m_i| \neq 0$. Thus $m(X)$ must consist of powers of cyclotomic polynomials. Similarly, for $l(X)$.

Any roots of $q(X)r(X)$ must come from $m(X)X^k - l(X)$. Unfortunately, we have not been able to likewise restrict the possibilities for $q(X)$ and $r(X)$. Some

simple cases which we have found are:

$$\begin{aligned} Y^2 &= ((X+1)X^n + X^k + 1)^2 - 4(X+1)X^n, \quad k \text{ odd}, \\ Y^2 &= (X^n + (X-1)X^k + 1)^2 - 4X^n, \\ Y^2 &= (X^n + X^3 + X^2 + 1)^2 + 4(X^3 + X^2 + X + 1)X, \end{aligned}$$

where in the last example $l(X) = (X^4 - 1)/(X - 1)$ and $k = 1$, whence it is necessary to have $n \equiv 1 \pmod{4}$. The expansion with $n = 9$ is included in the Tables at page 149.

We have been unable to find an example where more than one of the functions q, r, l, m is not a constant. Whether this is our fault or that of the problem is unknown.

Consequently, it seems wise to allow the functions q, r, m and l to also depend on n . If such a set of discriminants can still be written as a kreeper and satisfy the usual division requirements we say that it is an *extended-kreeper*. There will be more to say about such things later.

Each of the above examples corresponds to a case of a torsion divisor on the Jacobian of a hyperelliptic curve; its order being $\Theta(n^2)$. But this is exactly what Flynn was able to report after his investigations into torsion on hyperelliptic curves. In fact he found that the curve

$$Y_g^2 = (A^{(g)}(X))^2 - tX^g(X-1)^{g+1} \quad (22.1)$$

where

$$A^{(g)}(X) = (X^{g+1} - t(X-1)^g - X^g(X-1)) / 2$$

has torsion of order $2g^2 + 2g + 1$ at infinity. Since we are talking about the same thing shouldn't Flynn's curve be a kreeper?

The answer is yes. After a few simple transformations, (22.1) is transformed into

$$Y_g^2 = ((X-1)X^g + (X-1))^2 + 4(X-1)X^g$$

which is an obvious kreeper with $r = X - 1$. This is more than just a coincidence. Our results say that this must be a kreeper, but more importantly without them one would realise this from a deeper investigation into Flynn's curve. It arises by considering two divisors which are composed in such a way as to yield a unit. But this is precisely the manner in which we derived our keepers and so the two approaches are identical. Following this realisation one is naturally led to ask whether further results regarding torsion on elliptic and hyperelliptic curves might yield new information about keepers, and this idea is the basis of Chapter VIII.

Example 10. Here is an example of a function field kreeper over a finite field.

$$Y^2 = (4(X^2 + 1)X^n + X^3 + 4X^2 + X + 1)^2 + (X^2 + 4)X^3.$$

In this example, where the base field needs to be $\mathbb{F}_5[X]$ in order to provide a kreeper, we have taken

$$r = 4(X^2 + 1), \quad l = X + 1, \quad m = X - 1, \quad k = 3, \quad n \equiv 0 \pmod{2}.$$

Here is the expansion for $n = 8$,

h	$a_h(X)$	$Q_h(X)$
0	$4X^{10} + 4X^8 + X^4 + 4X^3 + X + 1$	1
1	$3X^5 + X^3 + X$	$X^3(X+1)(X+4)$
2	$2X + 3$	$Q_2(X)$
3	$3X^2 + 3X + 3$	$X^5(X+2)(X+3)(X+4)$
4	$3X^4 + 3X^2$	X^6
5	$3X^5 + 2X^4 + 3X^3 + 2X^2 + 3X + 2$	$X^2(X+1)(X+2)(X+3)$

6	$2X^2 + 2$	$4X^8 + X^2 + 1$
7	$3X^6 + 3X^5 + 3X^4 + 3X^3 + 3X^2 + 3X$	$X(X+2)(X+3)(X+4)$
8	$2X + 3$	$Q_8(X)$
9	$3X$	$X^7(X+1)(X+4)$
10	$2X + 2$	$Q_{10}(X)$
11	$3X^3 + 2X^2 + 3X + 2$	$X^4(X+1)(X+2)(X+3)$
12	$3X^6 + 3X^4 + 2$	X^4
13	$2X$	$4X^9 + 4X^7 + X^3 + X + 1$
14	$3X^7 + X^5 + X^3 + 3X + 3$	$X(X+1)(X+4)$
15	$3X^8 + 3X^6 + 2X^2 + 3X$	X^2
16	$3X + 2$	$X^6(X+1)(X+2)(X+3)$
17	$2X + 2$	$Q_{17}(X)$
18	$3X^3 + X$	$X^5(X+1)(X+4)$
19	$2X + 3$	$Q_{19}(X)$
20	$3X^4 + 3X^3 + 3X^2 + 3X + 3$	$X^3(X+2)(X+3)(X+4)$
21	$3X^2 + 3$	X^8
22	$3X^7 + 2X^6 + 3X^5 + 2X^4 + 3X^3 + 2X^2 + 3$	$(X+1)(X+2)(X+3)$

where:

$$Q_2(X) = 4X^9 + 4X^8 + 3X^7 + 4X^6 + 4X^5 + 2X^3 + 2X^2 + 2X + 1 ,$$

$$Q_8(X) = 4X^9 + 4X^8 + 4X^7 + X^3 + X^2 + X + 1 ,$$

$$Q_{10}(X) = 4X^9 + X^8 + 4X^7 + X^5 + 4X^4 + X^3 + 1 ,$$

$$Q_{17}(X) = 4X^9 + X^8 + 4X^7 + X^6 + X^5 + 3X^4 + X^3 + 1 ,$$

$$Q_{19}(X) = 4X^9 + 4X^8 + 3X^7 + 3X^6 + 4X^5 + X^4 + 2X^3 + 2X^2 + 2X + 1 .$$

This line is halfway in the expansion because $(X+1)(X+2)(X+3) \mid Y^2$ over $\mathbb{F}_5[X]$.

The period lengths and regulators satisfy:

$n \pmod{6}$	$lp(Y)$	$R(Y)$
0	$2n - 2$	$\frac{1}{3} (2n^2 + 2n + 3)$
2, 4	$6n - 4$	$2n^2 + 2n + 3$

Chapter VII. The general expansion of a kreeper

In its full generality, a kreeper can be expressed as

$$D_n = \frac{c^2}{d^2} \left[(qrx^n + (z^2mx^k - y^2l)/q)^2 + 4ry^2lx^n \right]$$

with the conditions:

$$\begin{aligned} m, l, r \text{ squarefree}, \quad c^2 r l y^2 m z^2 \mid d^2 D_n, \\ (qr, x) = 1, \quad (qrx, mzy l) = 1, \quad (yl, mz) = 1, \end{aligned}$$

where of course $x > 1$. In this section we show that if these conditions hold then there exists $a, b \in \mathbb{Q}$ such that

$$lp(\omega_n) = an + b$$

for infinitely many n . (As we know, keeping the square factors in D_n causes more difficulty in expanding ω_n . While it is not difficult to show that $lp(\omega'_n) = a'n + b'$ implies that $lp(\omega_n) = O(n)$, it doesn't appear obvious that we have the stronger assertion that $lp(\omega_n) = an + b$.)

To begin with, we write

$$G = c^2 y^2 l r, \quad H = c^2 z^2 m r x^k, \quad A = c q r,$$

so then

$$D_n = (s_1/d)^2 + 4G/d^2 x^n = (s_2/d)^2 + 4H/d^2 x^n.$$

As before, we put:

$$\alpha = \omega_n + S_1/d, \quad \beta = \omega_n + S_2/d,$$

with this notation we have

$$S_1 + S_2 + td = Ax^n.$$

As was noticed in the examples earlier, there is a congruence class of n in which the constant partial quotients remain the same. Indeed it will greatly facilitate our working if we restrict our attention to such a class of n 's.

Define, as before,

$$\begin{aligned} s &:= \max_{i \in \mathbb{N}} \{(x^i, c)\}, \\ u &:= \frac{c}{s}, \end{aligned}$$

so that $(u, x) = 1$. In order to make things easier for ourselves when we come to the expansion of ω_n , we would like to have:

$$\begin{aligned} x^\mu &\equiv 1 \pmod{u^2 m z^2 l y^2}, \\ x^\mu &\equiv 0 \pmod{s^2}. \end{aligned}$$

By our conditions, such a μ must exist. We shall want to consider the congruence class, $I_\nu = \{n \in \mathbb{N} : n \equiv \nu \pmod{\mu}, n > \mu\}$. Our proof will show that D_n is a kreeper for $n \in I_\nu$, and since every n lies in some I_ν , we shall not be losing any generality in this restriction.

More precisely, if

$$\begin{aligned} K &\equiv 1 \pmod{u^2 m z^2 l y^2}, \\ K &\equiv 0 \pmod{s^2}, \end{aligned}$$

then

$$K^i \equiv K \pmod{u^2 s^2 m z^2 l y^2}.$$

So,

$$x^{i\mu+\nu} \equiv x^{\mu+\nu} \pmod{u^2 s^2 m z^2 l y^2},$$

that is, the value of $x^n \pmod{u^2 s^2 m z^2 l y^2}$ is the same for all $n \in I_\nu$.

As we have seen, the main points of interest in the continued fraction expansion of ω_n occur when $x|Q_h$. Consequently, we shall “jump” between these elements when expanding ω_n . Naturally, the object must then be to find the first occurrence of $x|Q_h$.

As before we take:

$$g = (s_1, s_2, d), \quad d_y = (S_1 + td, d), \quad d_z = (S_2 + td).$$

Lemma 34 on page 73 determined that $d_y|y$ and $d_z|z$.

23. DETERMINATION OF THE SPECIFIC ELEMENTS IN THE EXPANSION

In Chapter V we determined the existence of the following complete quotient in the expansion of ω_n ,

$$\theta_{h_0} = \frac{\omega_n + S_1/d - c_0 c^2 r |l| y(y/d_y) x^n / d}{c^2 r |l| (y/d_y)^2 x^n},$$

with:

$$A_0 = c q r x^n - c_0 c^2 r |l| y(y/d_y) x^n,$$

$$B_0 = d c^2 r |l| (y/d_y)^2 x^n,$$

$$\Delta_0 = (A_0, B_0),$$

$$c_0 \equiv \frac{-\text{sign}(l)}{(S_1 + td)/d_y} \pmod{\bar{d} d_z}.$$

Furthermore, $\Delta_0 = c r x^n \delta d_z$ where $\delta = (c, q)$. We also found,

$$\theta_{h_1} = \frac{\omega_n + S_2/d - c_1 (c/\delta) z(z/d_z) x^k / d}{(c/\delta)^2 |m l| (y/d_y)^2 (z/d_z)^2 x^k}, \quad (23.1)$$

where

$$c_1 \equiv -\text{sign}(m) \Delta_0 / A_0 \pmod{B_0 / \Delta_0}.$$

Moreover, for sufficiently large n , θ_{h_1} is reduced since $\theta_{h_1} > 1$ and $-1 \leq \bar{\theta}_{h_1} < 0$. The θ_{h_i} 's are going to represent the complete quotients in the expansion of ω_n where $x|Q_{h_i}$. As we already know, these points are the critical ones in the expansion and the partial quotients in between are not of great importance.

Since we wish to determine the fundamental unit, we will be multiplying the complete quotients together, as in Proposition 5. Consequently, if θ_{i+1} represents the $(i+1)$ th complete quotient, that is $\omega = [a_0, a_1, \dots, a_i, \theta_{i+1}]$, we define,

$$\Psi_{i+1} = \theta_1 \dots \theta_{i-1} \theta_i Q_i.$$

Then, we have $N(\Psi_{i+1}) = (-1)^i Q_i$. Also, if we apply Lemma 2 to

$$\theta_{h_i} = \frac{A_i}{B_i} - \gamma \quad (23.2)$$

then

$$\theta_{h_{i+1}} = \frac{(-1)^{p_i} \bar{\gamma}}{\gamma \bar{\gamma} (B_i / \Delta_i)^2} - \frac{c_i}{B_i / \Delta_i}$$

where $c_i \equiv (-1)^{p_i+1} \Delta_i / A_i \pmod{B_i / \Delta_i}$. Thus,

$$Q_{h_{i+1}} = (-1)^{p_i} \gamma \bar{\gamma} (B_i / \Delta_i)^2 Q_{h_i}.$$

Hence,

$$\frac{(-1)^{p_i} Q_{h_{i+1}}}{Q_{h_i}} = \frac{(A_i - B_i \theta_{h_i})(A_i - B_i \bar{\theta}_{h_i})}{\Delta_i^2}.$$

By Proposition 5,

$$\begin{aligned} \theta_{h_i+1} \theta_{h_i+2} \dots \theta_{h_{i+1}} &= \frac{(-1)^{p_i}}{A_i / \Delta_i - B_i / \Delta_i \theta_{h_i}} \\ &= \frac{(A_i - \bar{\theta}_{h_i} B_i) Q_{h_i}}{\Delta_i Q_{h_{i+1}}}. \end{aligned}$$

Hence,

$$\begin{aligned} \Psi_{h_{i+1}+1} &= \Psi_{h_i+1} \theta_{h_i+1} \dots \theta_{h_{i+1}} Q_{h_{i+1}} \\ &= \left(\frac{A_i - B_i \bar{\theta}_{h_i}}{\Delta_i} \right) \Psi_{h_i+1}. \end{aligned} \tag{23.3}$$

Also note that from (23.2), $(A_i - B_i \bar{\theta}_{h_i}) / \Delta_i = \bar{\gamma}$.

Combining the above results with those in Chapter V, we have

$$\Psi_{h_1+1} = \left(\frac{A_0 - B_0 \bar{\theta}_1}{\Delta_0} \right) \Psi_1$$

where $\Psi_1 = \alpha$,

$$\bar{\theta}_1 = \frac{\bar{\omega}_n + S_1/d - c_0 c^2 r |l| y(y/d_y) x^n / d}{c^2 r |l| (y/d_y)^2 x^n}$$

and

$$\frac{A_0 - B_0 \bar{\theta}_1}{\Delta_0} = \frac{\beta}{d \Delta_0}.$$

Thus, $\Psi_{h_1+1} = \frac{\alpha \beta}{d \Delta_0}$.

23.1. Recursions from Chapter V

As we have seen in the earlier examples, the development of the expansion of ω_n depends upon whether or not a power of x^k can be factored out from Q_{h_i} or not. In order to accommodate this we define:

$$\varepsilon_i = \begin{cases} 1 & \text{if } \lambda_i \geq n - k, \\ 0 & \text{if } \lambda_i < n - k, \end{cases}$$

where λ_i is defined recursively as

$$\begin{aligned} \lambda_1 &:= k, \\ \lambda_{i+2} &:= \begin{cases} \lambda_i + k - n \varepsilon_i & \text{if } i \equiv 1 \pmod{2} \\ 0 & \text{if } i \equiv 0 \pmod{2} \end{cases} \end{aligned}$$

Note that $\lambda_{2i-1} \equiv ki \pmod{n}$. The reason for keeping λ_{2i} equal to 0 is that in the expansion, only every second complete quotient has an increase/decrease in the exponent of x . And $\varepsilon_i = 1$ represent those points in the expansion when the progression of powers of x changes (remember that this issue does not arise in the

case of $k|n$). With these notations, the recursions in Chapter V may be written in the following manner,

$$\begin{aligned} P_{h_i} &= S_2/d - sm_i r_i u_i u'_i z_i z'_i c_i x^{\lambda_i} / d \\ Q_{h_i} &= r_i l_i m_i (su_i y_i z_i)^2 x^{\lambda_i} \end{aligned}$$

where:

$$\begin{aligned} l_i &\in \{1, |l|\}, \quad m_i \in \{1, |m|\}, \quad r_i \in \{1, r\}, \quad u_i | u, \quad u'_i | u, \\ y_i | y/d_y, \quad z_i | z/d_z, \quad z'_i | z, \quad \lambda_i < n, \\ (c_i, s) &= 1, \quad (z_i y_i, u/u_i) = 1. \end{aligned}$$

The complete quotient θ_{h_1} in (23.1) is the case of $i = 1$ with the selections:

$$\begin{aligned} l_1 &= |l|, \quad m_1 = |m|, \quad r_1 = 1, \quad u_1 = u/\delta, \quad u'_1 = 1, \\ y_1 &= y/d_y, \quad z_1 = z/d_z, \quad z'_1 = z. \end{aligned}$$

Note that other than for λ_i , all the above values depend only on i and $\lambda_i \pmod{\mu}$. In Chapter V we determined that from the complete quotient,

$$\theta_{h_i} = \frac{\omega_n + P_{h_i}}{Q_{h_i}} \quad (23.4)$$

another complete quotient $\theta_{h_{i+1}}$ is given by,

$$P_{h_{i+1}} = S_1/d - sr_{i+1} l_{i+1} u_{i+1} u'_{i+1} y_{i+1} y'_{i+1} c_{i+1} x^{n-\lambda_i} / d \quad (23.5a)$$

$$Q_{h_{i+1}} = r_{i+1} l_{i+1} m_{i+1} (su_{i+1} y_{i+1} z_{i+1})^2 x^{n-\lambda_i} \quad (23.5b)$$

where:

$$\begin{aligned} r_{i+1} &\in \{1, r\}, \quad l_{i+1} \in \{1, |l|\}, \quad m_{i+1} \in \{1, |m|\}, \quad u_{i+1} | u, \quad u'_{i+1} | u, \\ y_{i+1} &| y/d_y, \quad y'_{i+1} | y, \quad z_{i+1} | z/d_z, \quad (z_{i+1} y_{i+1}, u/u_{i+1}) = 1, \quad (c_{i+1}, s) = 1. \end{aligned}$$

Furthermore,

$$\Psi_{h_{i+1}+1} = \left(\frac{A_i - B_i \bar{\theta}_{h_i}}{\Delta_i} \right) \Psi_{h_i+1}$$

where:

$$\begin{aligned} A_i &= cqr x^n - sr_i m_i u_i u'_i z_i z'_i c_i x^{\lambda_i}, \\ B_i &= dr_i l_i m_i (su_i y_i z_i)^2 x^{\lambda_i}, \\ \Delta_i &= (A_i, B_i), \\ &= sd_y u_i l_i y_i r_i w_i \bar{\Delta}_i x^{\lambda_i}. \end{aligned}$$

In this case,

$$\frac{A_i - B_i \bar{\theta}_{h_i}}{\Delta_i} = \frac{\alpha}{d \Delta_i} \quad \text{which implies that} \quad \Psi_{h_{i+1}+1} = \left(\frac{\alpha}{d \Delta_i} \right) \Psi_{h_i+1}.$$

If $\lambda_i \leq n - k$ then $\varepsilon_i = 0$ and $\lambda_{i+2} = \lambda_i + k$. In this case, we established, in Chapter V, the existence of the complete quotient $\theta_{h_{i+2}}$ with

$$\begin{aligned} P_{h_{i+2}} &= S_2/d - sr_{i+2} m_{i+2} u_{i+2} u'_{i+2} z_{i+2} z'_{i+2} c_{i+2} x^{\lambda_{i+2}} / d \\ Q_{h_{i+2}} &= r_{i+2} l_{i+2} m_{i+2} (su_{i+2} y_{i+2} z_{i+2})^2 x^{\lambda_{i+2}} \end{aligned}$$

where

$$\begin{aligned} r_{i+2} &\in \{1, r\}, \quad l_{i+2} \in \{1, |l|\}, \quad m_{i+2} \in \{1, |m|\}, \quad u_{i+2} | u, \quad u'_{i+2} | u, \\ y_{i+2} &| y/d_y, \quad z_{i+2} | z/d_z, \quad z'_{i+2} | z, \quad (z_{i+2} y_{i+2}, u/u_{i+2}) = 1, \quad (c_{i+2}, s) = 1. \end{aligned}$$

Furthermore,

$$\begin{aligned} A_{i+1} &= cqr x^n - sr_{i+1}l_{i+1}u_{i+1}u'_{i+1}y_{i+1}y'_{i+1}c_{i+1}x^{n-\lambda_i}, \\ B_{i+1} &= dr_{i+1}l_{i+1}m_{i+1}(su_{i+1}y_{i+1}z_{i+1})^2x^{n-\lambda_i}, \\ \Delta_{i+1} &= (A_{i+1}, B_{i+1}), \\ &= sd_z u_{i+1}m_{i+1}z_{i+1}r_{i+1}w_{i+1}\overline{\Delta}_{i+1}x^{n-\lambda_i}. \end{aligned}$$

And,

$$\frac{A_{i+1} - B_{i+1}\overline{\theta}_{h_{i+1}}}{\Delta_{i+1}} = \frac{\beta}{d\Delta_{i+1}}.$$

From,

$$\Psi_{h_{i+2}+1} = \left(\frac{A_{i+1} - B_{i+1}\overline{\theta}_{h_{i+1}}}{\Delta_{i+1}} \right) \Psi_{h_{i+1}+1}$$

we have

$$\Psi_{h_{i+2}+1} = \left(\frac{\beta}{d\Delta_{i+1}} \right) \Psi_{h_{i+1}+1} = \left(\frac{\alpha\beta}{d^2\Delta_i\Delta_{i+1}} \right) \Psi_{h_i+1}.$$

23.2. When $\varepsilon_i = 1$

However, when $\lambda_i \geq n - k$ we have $\varepsilon_i = 1$ and this case was not considered in Chapter V. In this situation, we need to consider

$$\theta_{h_{i+1}} = \frac{s_1/d - sr_{i+1}l_{i+1}u_{i+1}u'_{i+1}y_{i+1}y'_{i+1}c_{i+1}x^{n-\lambda_i}/d}{r_{i+1}l_{i+1}m_{i+1}(su_{i+1}y_{i+1}z_{i+1})^2x^{n-\lambda_i}} - \frac{\overline{\alpha}}{r_{i+1}l_{i+1}m_{i+1}(su_{i+1}y_{i+1}z_{i+1})^2x^{n-\lambda_i}}.$$

There is a slight problem in notation now because there is going to be an extra intermediate complete quotient. Consequently, we will use overlines to represent the terms involved. This time we take:

$$\begin{aligned} \overline{A}_{i+1} &:= s_1 - sr_{i+1}l_{i+1}u_{i+1}u'_{i+1}y_{i+1}y'_{i+1}c_{i+1}x^{n-\lambda_i}, \\ \overline{B}_{i+1} &:= dr_{i+1}l_{i+1}m_{i+1}(su_{i+1}y_{i+1}z_{i+1})^2x^{n-\lambda_i}, \\ \overline{\Delta}_{i+1} &:= (\overline{A}_{i+1}, \overline{B}_{i+1}). \end{aligned}$$

Then,

$$\overline{A}_{i+1} = dP_{h_{i+1}} + (s_1 + td)/2. \quad (23.6)$$

In Theorem 37 on page 76, the only prominent fact needed was (23.6). Hence the result found there will equally apply in this case, that is $d_y u_{i+1}l_{i+1}y_{i+1} | \overline{A}_{i+1}$. We also define:

$$\begin{aligned} \overline{w}_{i+1} &:= (m_{i+1}, u/u_{i+1}), \\ \overline{e}_{i+1} &:= sd_y u_{i+1}l_{i+1}y_{i+1}r_{i+1}\overline{w}_{i+1}. \end{aligned}$$

Unlike the case of $\varepsilon_i = 0$, it is not immediately obvious that $u_{i+1}\overline{w}_{i+1} | \overline{A}_{i+1}$, but it does.

Lemma 41. $u_{i+1}\overline{w}_{i+1}$ divides $\overline{A}_{i+1}$.

Proof. First, $u_{i+1}\overline{w}_{i+1} | u$ and $u_{i+1}\overline{w}_{i+1} | u_{i+1}m_{i+1}$. Also note that $u^2 | D_n$ and $m_{i+1}u_{i+1}^2 | Q_{h_{i+1}}$. From

$$-D_n + (2P_{h_{i+1}} + t)^2 = -4Q_{h_{i+1}-1}Q_{h_{i+1}},$$

we have $u_{i+1}\bar{w}_{i+1} \mid (2P_{h_{i+1}} + t)$. And,

$$2\bar{A}_{i+1} = 2dP_{h_{i+1}} + td + s_1.$$

Since $u_{i+1}\bar{w}_{i+1} \mid s_1$ we have that $u_{i+1}\bar{w}_{i+1} \mid 2\bar{A}_{i+1}$. If $2 \nmid u$ then $u_{i+1}\bar{w}_{i+1} \mid \bar{A}_{i+1}$. Now suppose that $2 \mid u$, so then $2 \nmid d$ and $t = 0$. It is also clear that $u_{i+1} \mid \bar{A}_{i+1}$, so that we need only consider the case of $2 \mid \bar{w}_{i+1}$. Since m_{i+1} is squarefree we must have $2 \parallel m_{i+1}$ and

$$E_n = \frac{d^2 D_n}{c^2} = (s_1/c)^2 + 4rmz^2x^k \equiv 0 \pmod{4},$$

so $2 \mid s_1/c$. Then recalling that $u_{i+1}\bar{w}_{i+1} \mid c$ we must have $2u_{i+1}\bar{w}_{i+1} \mid s_1$. Moreover, $c^2m \mid d^2D_n$ and $2 \nmid d$ imply that $2u^2 \mid D_n$ which implies that $2u_{i+1}^2\bar{w}_{i+1}^2 \mid D_n$, and so $4u_{i+1}\bar{w}_{i+1} \mid D_n$. From

$$-D_n/4 + P_{h_{i+1}}^2 = -Q_{h_{i+1}-1}Q_{h_{i+1}},$$

we have $u_{i+1}\bar{w}_{i+1} \mid P_{h_{i+1}}$. Hence,

$$\bar{A}_{i+1} = dP_{h_{i+1}} + s_1/2 \equiv 0 \pmod{u_{i+1}\bar{w}_{i+1}}.$$

■

Since $s\bar{w}_{i+1}u_{i+1}r_{i+1} \mid \bar{A}_{i+1}$ and

$$(d_y l_{i+1} y_{i+1}, s\bar{w}_{i+1} r_{i+1}) = 1$$

we find $\bar{e}_{i+1} \mid \bar{A}_{i+1}$. We have,

$$\frac{\bar{B}_{i+1}}{\bar{e}_{i+1}} = \frac{sdm_{i+1}u_{i+1}y_{i+1}z_{i+1}^2x^{n-\lambda_i}}{d_y\bar{w}_{i+1}}$$

and

$$\frac{\bar{A}_{i+1}}{\bar{e}_{i+1}} = \frac{1}{d_y l_{i+1} y_{i+1} \bar{w}_{i+1}} \times \left[\frac{ur}{u_{i+1}r_{i+1}}x^n - \frac{u}{u_{i+1}} \frac{(mz^2x^k - ly^2)}{qr_{i+1}} - l_{i+1}u'_{i+1}y_{i+1}y'_{i+1}c_{i+1}x^{n-\lambda_i} \right]$$

The next two results establish that $\bar{A}_{i+1}/\bar{e}_{i+1}$ and $z_{i+1}m_{i+1}/\bar{w}_{i+1}$ are coprime.

Lemma 42. z_{i+1} and $\bar{A}_{i+1}/\bar{e}_{i+1}$ are coprime.

Proof. From,

$$-d^2D_n + (d(2P_{h_{i+1}} + t))^2 = -4d^2Q_{h_{i+1}-1}Q_{h_{i+1}} \quad (23.7)$$

we get

$$u_{i+1}m_{i+1}z_{i+1} \mid d(2P_{h_{i+1}} + t).$$

We have $(z_{i+1}, u/u_{i+1}) = 1$. It also holds that $z^2 \mid E_n$ which means that $z \mid s_2/c$, hence $z \mid s_2/u$ because $(s, u) = 1$. Let $\delta = (z, s_1/u)$. Since $\delta \mid s_2/u$ we get

$$\delta \mid (s_1 + s_2)/u = 2sqr x^n.$$

Since $(z, sqr x) = 1$ we get $\delta \mid 2$. Thus $(z_{i+1}, s_1/u_{i+1}) \mid 2$. Then from,

$$\frac{2\bar{A}_{i+1}}{u_{i+1}} = \frac{2dP_{h_{i+1}} + td}{u_{i+1}} + \frac{s_1}{u_{i+1}} \quad (23.8)$$

we have that $(z_{i+1}, \bar{A}_{i+1}/u_{i+1}) \mid 2$. Now suppose that $(z_{i+1}, \bar{A}_{i+1}/\bar{w}_{i+1}) = 2$.

We first consider the case of $4 \mid s_1/u_{i+1}$. Then $2 \parallel z_{i+1}$ and

$$E_n = (s_1/c)^2 + 4rly^2x^n \not\equiv 0 \pmod{8} \quad \text{since } (2, rlyx) = 1.$$

From (23.8), $4 \mid d(2P_{h_{i+1}} + t)/u_{i+1}$. Then

$$d^2 D_n = (d(2P_{h_{i+1}} + t))^2 + 4d^2 Q_{h_{i+1}-1} Q_{h_{i+1}} \equiv 0 \pmod{8}.$$

Thus, $2 \mid c$ since $d^2 D_n/c^2 \not\equiv 0 \pmod{8}$, hence $2 \nmid d$ and $t = 0$. Similarly,

$$\frac{D_n}{u_{i+1}^2} = \frac{4P_{h_{i+1}}^2}{u_{i+1}^2} + \frac{4Q_{h_{i+1}-1}Q_{h_{i+1}}}{u_{i+1}^2} \equiv 0 \pmod{8}$$

Thus,

$$8 \mid \frac{D_n}{u_{i+1}^2} \text{ which implies that } 8 \mid \frac{D_n}{u^2} \text{ because } (u/u_{i+1}, z_{i+1}) = 1.$$

It follows that $8 \mid D_n/c^2$ and since $2 \nmid d$, we have $8 \mid E_n$. But this is a contradiction, which means that $2 \parallel s_1/u_{i+1}$. Then from (23.8), we have $2 \parallel d(2P_{h_{i+1}} + t)$.

If $t = 0$ then $2 \nmid d$ and $2 \nmid P_{h_{i+1}}/u_{i+1}$. Moreover,

$$\frac{P_{h_{i+1}}^2}{u_{i+1}^2} \equiv -Q_{h_{i+1}-1} \frac{Q_{h_{i+1}}}{u_{i+1}^2} \pmod{2}.$$

Since $2 \mid z_{i+1}$ and $z_{i+1} \mid Q_{h_{i+1}}/u_{i+1}^2$, we get $2 \mid P_{h_{i+1}}/u_{i+1}$ which is a contradiction.

If $t = 1$ then D_n is odd and $2 \parallel d$ which means that $2 \parallel s_2$ and $2 \parallel s_1$. But then

$$q(s_1 + s_2)/2c = q^2 r x^n \equiv 0 \pmod{2}$$

which means $2 \mid (qrx, z)$ which is also a contradiction. In conclusion we must have $(z_{i+1}, \bar{A}_{i+1}/u_{i+1}) = 1$ and therefore, $(z_{i+1}, \bar{A}_{i+1}/\bar{e}_{i+1}) = 1$. ■

Lemma 43. $(m_{i+1}/\bar{w}_{i+1}, \bar{A}_{i+1}/\bar{w}_{i+1}u_{i+1}) = 1$.

Proof. Let $p \mid (m_{i+1}/\bar{w}_{i+1}, \bar{A}_{i+1}/\bar{w}_{i+1}u_{i+1})$ with $p > 1$. Then $p \nmid u/(u_{i+1}\bar{w}_{i+1})$ by the definition of $\bar{w}_{i+1}$. From (23.7), we have $m_{i+1}u_{i+1} \mid d(2P_{h_{i+1}} + t)$ and

$$\frac{2\bar{A}_{i+1}}{u_{i+1}\bar{w}_{i+1}} = \frac{2dP_{h_{i+1}} + td}{u_{i+1}\bar{w}_{i+1}} + \frac{s_1}{u_{i+1}\bar{w}_{i+1}}. \quad (23.9)$$

Hence, $p \mid s_1/(u_{i+1}\bar{w}_{i+1})$. Since $(m, s_1/c) \mid 2$ we must have $p = 2$. Thus,

$$2 \nmid \frac{u}{u_{i+1}\bar{w}_{i+1}}.$$

Also, if $2 \mid m_{i+1}/\bar{w}_{i+1}$ then since m_{i+1} is squarefree, $2 \nmid \bar{w}_{i+1}$. Hence $2 \nmid u/u_{i+1}$. We are now exactly in the same scenario as in Lemma 42 (since $m_{i+1} \mid Q_{h_{i+1}}$) and therefore we reach the same conclusion. ■

23.3. Expanding $\theta_{h_{i+1}}$

Next we take, $\bar{G}_{i+1} := \bar{A}_{i+1}/\bar{e}_{i+1}$. From Lemmas 42 and 43, we have

$$(\bar{G}_{i+1}, z_{i+1}m_{i+1}/\bar{w}_{i+1}) = 1.$$

Also note that if $p \mid (x, \bar{A}_{i+1})$ with $p > 1$ then p must divide s_1 , whence $p \mid ly^2$. However, this is impossible because $(x, ly^2) = 1$. Thus, $(\bar{G}_{i+1}, x) = 1$. Then,

$$\begin{aligned} \bar{\Delta}_{i+1} &= \bar{e}_{i+1} \left(\frac{\bar{A}_{i+1}}{\bar{e}_{i+1}}, \frac{\bar{B}_{i+1}}{\bar{e}_{i+1}} \right) \\ &= sdy_{i+1}l_{i+1}y_{i+1}r_{i+1}\bar{w}_{i+1} \left(\bar{G}_{i+1}, \frac{sdm_{i+1}u_{i+1}y_{i+1}z_{i+1}^2x^{n-\lambda_i}}{dy\bar{w}_{i+1}} \right) \\ &= sdy_{i+1}l_{i+1}y_{i+1}r_{i+1}\bar{w}_{i+1} (\bar{G}_{i+1}, \bar{d}d_z u_{i+1}y_{i+1}). \end{aligned}$$

Since,

$$\overline{A}_{i+1} = dP_{h_{i+1}} + S_1 + td$$

we have $(d, \overline{A}_{i+1}) = d_y$, and since $d_y | \overline{c}_{i+1}$ we have $(\overline{d}d_z, \overline{G}_{i+1}) = 1$. By writing, $\Delta'_{i+1} = (\overline{G}_{i+1}, u_{i+1}y_{i+1})$, we get

$$\overline{\Delta}_{i+1} = sd_y u_{i+1} l_{i+1} y_{i+1} r_{i+1} \overline{w}_{i+1} \Delta'_{i+1}.$$

Note that

$$\frac{\overline{B}_{i+1}}{\overline{\Delta}_{i+1}} = \frac{sdm_{i+1}u_{i+1}y_{i+1}z_{i+1}^2x^{n-\lambda_i}}{d_y \overline{w}_{i+1} \Delta'_{i+1}}.$$

Replacing the terms denoted by f , a , b , d in Lemma 39 by z_{i+1} , y/d_y , $u/\overline{w}_{i+1}$ and Δ'_{i+1} respectively, the Lemma says that there exist two numbers $\overline{y}_{i+2}$ and $\overline{u}_{i+2}$, such that

$$\overline{y}_{i+2} | y/d_y, \quad \overline{u}_{i+2} | u/\overline{w}_{i+1}, \quad \overline{y}_{i+2} \overline{u}_{i+2} \Delta'_{i+1} = \frac{y}{d_y} \frac{u}{\overline{w}_{i+1}},$$

and

$$\left(z_{i+1} \overline{y}_{i+2}, \frac{u}{\overline{u}_{i+2} \overline{w}_{i+1}} \right) = 1.$$

So $yu = \overline{y}_{i+2} \overline{u}_{i+2} d_y \overline{w}_{i+1} \Delta'_{i+1}$.

From,

$$\theta_{h_{i+1}} = \frac{\overline{A}_{i+1}}{\overline{B}_{i+1}} - \frac{\overline{\alpha}}{r_{i+1} l_{i+1} m_{i+1} (su_{i+1} y_{i+1} z_{i+1})^2 x^{n-\lambda_i}}$$

we apply Lemma 2 and find that the next partial quotients are the expansion of $\overline{A}_{i+1}/\overline{B}_{i+1}$ of length $\overline{p}_{i+1}$, where the parity of $\overline{p}_{i+1}$ is determined by $(-1)^{\overline{p}_{i+1}+1} = \text{sign}(l)$. The next complete quotient is then $\theta_{j_{i+2}}$ where

$$j_{i+2} = h_{i+1} + \overline{p}_{i+1}$$

and

$$\theta_{j_{i+2}} = \frac{r_{i+1} l_{i+1} m_{i+1} (su_{i+1} y_{i+1} z_{i+1})^2 x^{n-\lambda_i}}{-\text{sign}(l) \overline{\alpha} \left(\frac{sdm_{i+1}u_{i+1}y_{i+1}z_{i+1}^2x^{n-\lambda_i}}{d_y \overline{w}_{i+1} \Delta'_{i+1}} \right)^2} - \frac{\overline{c}_{i+2}}{\overline{B}_{i+1}/\overline{\Delta}_{i+1}} \quad (23.10)$$

where $\overline{c}_{i+2} \equiv -\text{sign}(l) \overline{\Delta}_{i+1}/\overline{A}_{i+1} \pmod{\overline{B}_{i+1}/\overline{\Delta}_{i+1}}$. Looking just at the first term of (23.10),

$$\frac{r_{i+1}}{r} \frac{l_{i+1}}{|l|} \frac{(\omega_n + S_1/d)(yu)^2}{m_{i+1}(suy_{i+1}\overline{y}_{i+2}\overline{u}_{i+2})^2 x^{2n-\lambda_i}}$$

which if we take:

$$\overline{r}_{i+2} := \frac{r}{r_{i+1}}, \quad \overline{l}_{i+2} := \frac{|l|}{l_{i+1}}, \quad \overline{m}_{i+2} := m_{i+1}, \quad \overline{z}_{i+2} := z_{i+1},$$

this becomes

$$\frac{\omega_n + S_1/d}{\overline{r}_{i+2} \overline{l}_{i+2} \overline{m}_{i+2} (s\overline{u}_{i+2} \overline{y}_{i+2} \overline{z}_{i+2})^2 x^{2n-\lambda_i}}.$$

The second term in (23.10) is

$$\frac{\overline{c}_{i+2} d_y \overline{w}_{i+1} \overline{\Delta}_{i+1}}{sdm_{i+1}u_{i+1}y_{i+1}z_{i+1}^2x^{n-\lambda_i}} = \frac{y}{y_{i+1}} \frac{u}{u_{i+1}} \frac{\overline{c}_{i+2}}{dsm_{i+1}z_{i+1}^2\overline{u}_{i+2}\overline{y}_{i+2}x^{n-\lambda_i}}.$$

If we take

$$\overline{y}'_{i+2} := \frac{y}{y_{i+1}} \quad \text{and} \quad \overline{u}'_{i+2} := \frac{u}{u_{i+1}}$$

this becomes

$$\frac{\overline{c}_{i+2} \overline{y}'_{i+2} \overline{u}'_{i+2}}{ds\overline{m}_{i+2} \overline{u}_{i+2} \overline{y}_{i+2} \overline{z}_{i+2}^2 x^{n-\lambda_i}}.$$

Also note that since $s|\overline{B}_{i+1}/\overline{\Delta}_{i+1}$ we get $(\overline{c}_{i+2}, s) = 1$. Also,

$$\left(z_{i+1}\overline{y}_{i+2}, \frac{u}{\overline{u}_{i+2}\overline{w}_{i+1}}\right) = 1 \quad \text{and} \quad (z_{i+1}\overline{y}_{i+2}, \overline{w}_{i+1}) = 1$$

imply that

$$(\overline{z}_{i+2}\overline{y}_{i+2}, u/\overline{u}_{i+2}) = 1.$$

In conclusion, the complete quotient, $\theta_{j_{i+2}}$ is

$$\theta_{j_{i+2}} = \frac{\omega_n + S_1/d - s\overline{r}_{i+2}\overline{l}_{i+2}\overline{u}_{i+2}\overline{u}'_{i+2}\overline{y}_{i+2}\overline{y}'_{i+2}\overline{c}_{i+2}x^{2n-\lambda_i}/d}{\overline{r}_{i+2}\overline{l}_{i+2}\overline{m}_{i+2}(s\overline{u}_{i+2}\overline{y}_{i+2}\overline{z}_{i+2})^2x^{2n-\lambda_i}} \quad (23.11)$$

where

$$\begin{aligned} \overline{r}_{i+2} &\in \{1, r\}, \quad \overline{l}_{i+2} \in \{1, |l|\}, \quad \overline{m}_{i+2} \in \{1, |m|\}, \quad \overline{u}_{i+2}|u, \quad \overline{u}'_{i+2}|u, \\ \overline{y}_{i+2}|y/d_y, \quad \overline{y}'_{i+2}|y, \quad \overline{z}_{i+2}|z/d_z, \quad (\overline{c}_{i+2}, s) &= 1, \quad (\overline{z}_{i+2}\overline{y}_{i+2}, u/\overline{u}_{i+2}) = 1. \end{aligned}$$

Furthermore,

$$\frac{\overline{A}_{i+1} - \overline{B}_{i+1}\overline{\theta}_{h_{i+1}}}{\overline{\Delta}_{i+1}} = \frac{\alpha}{d\overline{\Delta}_{i+1}},$$

hence,

$$\Psi_{j_{i+2}+1} = \left(\frac{\alpha}{d\overline{\Delta}_{i+1}}\right) \Psi_{h_{i+1}+1}.$$

23.4. Finishing the induction

Now, the complete quotient in (23.11) satisfies the same conditions as the complete quotient given in (23.5) on page 104. Hence we can find

$$\theta_{j_{i+3}} = \frac{\omega_n + S_2/d - s\overline{m}_{i+3}\overline{r}_{i+3}\overline{u}_{i+3}\overline{u}'_{i+3}\overline{z}_{i+3}\overline{z}'_{i+3}\overline{c}_{i+3}x^{\lambda_i+k-n}/d}{\overline{r}_{i+3}\overline{l}_{i+3}\overline{m}_{i+3}(s\overline{u}_{i+3}\overline{y}_{i+3}\overline{z}_{i+3})^2x^{\lambda_i+k-n}} \quad (23.12)$$

with

$$\begin{aligned} \overline{r}_{i+3} &\in \{1, r\}, \quad \overline{l}_{i+3} \in \{1, |l|\}, \quad \overline{m}_{i+3} \in \{1, |m|\}, \quad \overline{u}_{i+3}|u, \quad \overline{u}'_{i+3}|u, \\ \overline{y}_{i+3}|y/d_y, \quad \overline{z}_{i+3}|z/d_z, \quad \overline{z}'_{i+3}|z, \quad j_{i+3} &= j_{i+2} + \overline{p}_{i+2}, \\ (\overline{c}_{i+3}, s) &= 1, \quad (\overline{z}_{i+3}\overline{y}_{i+3}, u/\overline{u}_{i+3}) = 1, \end{aligned}$$

and

$$\Psi_{j_{i+3}+1} = \left(\frac{\beta}{d\overline{\Delta}_{i+2}}\right) \Psi_{j_{i+2}+1} = \left(\frac{\alpha\beta}{d^2\overline{\Delta}_{i+1}\overline{\Delta}_{i+2}}\right) \Psi_{h_{i+1}+1}.$$

Since $\varepsilon_i = 1$ we have $\lambda_{i+2} = \lambda_i + k - n$ and by renaming

$$\begin{aligned} m_{i+2} &:= \overline{m}_{i+3}, \quad r_{i+2} := \overline{r}_{i+3}, \quad l_{i+2} := \overline{l}_{i+3}, \quad u_{i+2} := \overline{u}_{i+3}, \quad u'_{i+2} := \overline{u}'_{i+3}, \\ y_{i+2} &:= \overline{y}_{i+3}, \quad z_{i+2} := \overline{z}_{i+3}, \quad z'_{i+2} := \overline{z}'_{i+3}, \quad h_{i+2} := j_{i+3}, \\ p_{i+2} &:= \overline{p}_{i+1} + \overline{p}_{i+2}, \quad \Delta_{i+1} := d\overline{\Delta}_{i+1}\overline{\Delta}_{i+2}, \end{aligned}$$

the complete quotient $\theta_{j_{i+3}}$ in (23.12) becomes

$$\theta_{h_{i+2}} = \frac{\omega_n + S_2/d - sr_{i+2}m_{i+2}u_{i+2}u'_{i+2}z_{i+2}z'_{i+2}c_{i+2}x^{\lambda_{i+2}}/d}{r_{i+2}l_{i+2}m_{i+2}(su_{i+2}y_{i+2}z_{i+2})^2x^{\lambda_{i+2}}}$$

with

$$\begin{aligned} r_{i+2} &\in \{1, r\}, \quad l_{i+2} \in \{1, |l|\}, \quad m_{i+2} \in \{1, |m|\}, \quad u_{i+2}|u, \quad u'_{i+2}|u, \\ y_{i+2}|y/d_y, \quad z_{i+2}|z/d_z, \quad z'_{i+2}|z, \quad (c_{i+2}, s) &= 1, \quad (z_{i+2}y_{i+2}, u/u_{i+2}) = 1, \end{aligned}$$

and

$$\Psi_{h_{i+2}+1} = \left(\frac{\alpha\beta}{d\Delta_{i+1}}\right) \Psi_{h_{i+1}+1}.$$

Combining both the $\varepsilon_i = 0$ and $\varepsilon_i = 1$ cases, we have

$$\begin{aligned}\Psi_{h_{i+2}+1} &= \left(\frac{\alpha^{\varepsilon_i} \beta}{d \Delta_{i+1}} \right) \Psi_{h_{i+1}+1} \\ &= \left(\frac{\alpha^{\varepsilon_i+1} \beta}{d^2 \Delta_i \Delta_{i+1}} \right) \Psi_{h_{i+1}},\end{aligned}$$

and

$$h_{i+2} = h_i + p_i + p_{i+1}. \quad (23.13)$$

This finally completes the induction process. Our next step will be to investigate the number of partial quotients in the period of the continued fraction expansion of ω_n .

24. DETERMINING THE NUMBER OF PARTIAL QUOTIENTS

By defining, $f_0 := h_0$ and $f_{i+1} := p_i$ for $i \geq 1$ we find that (23.13) becomes the more convenient,

$$h_i = \sum_{k=0}^i f_k. \quad (24.1)$$

Furthermore the value of f_i is dependent only upon the set

$$Z_i := \{l_i, m_i, r_i, s_i, s'_i, z_i, z'_i, u_i, u'_i, y_i, y'_i, c_i, L_i, x\}$$

where $L_i := \lambda_i \pmod{\mu}$. Moreover, if $Z_i = Z_j$ then $f_i = f_j$. Furthermore, once ε_i is determined then Z_{i+1} can be found from Z_i .

Letting $\theta := (n - \nu)/\mu$, then clearly all of the elements of Z_i are bounded independently of θ . Denote the total number of distinct sets Z_i by Z .

Since

$$\varepsilon_{2j-1} = \left\lfloor \frac{(j+1)k}{n} \right\rfloor - \left\lfloor \frac{jk}{n} \right\rfloor$$

we have that

$$\sum_{j=0}^{2nt-1} \varepsilon_j = \sum_{j=0}^{tn} \varepsilon_{2j-1} = \left\lfloor \frac{(tn+1)k}{n} \right\rfloor = tk.$$

So there are precisely tk values of $i \in \{1, \dots, 2tn-1\}$ where $\varepsilon_i = 1$. Let $i_1, \dots, i_{tk}$ represent these points. Then,

$$\begin{aligned}\sum_{j=0}^{i_h-1} \varepsilon_j = h-1 &\text{ implies } \left\lfloor \frac{\frac{1}{2}(i_h+2)k}{n} \right\rfloor = h-1, \\ \sum_{j=0}^{i_h} \varepsilon_j = h &\text{ implies } \left\lfloor \frac{\frac{1}{2}(i_h+3)k}{n} \right\rfloor = h.\end{aligned}$$

So,

$$\begin{aligned}n(h-1) &\leq (i_h+2)k/2 < hn, \\ nh &\leq (i_h+3)k/2 < (h+1)n.\end{aligned}$$

Combining these we have

$$\begin{aligned}nh &\leq (i_h+3)k/2 < hn + k/2, \\ \frac{2nh}{k} - 1 &\leq i_h + 2 < \frac{2nh}{k}.\end{aligned}$$

In other words,

$$i_h = \begin{cases} \frac{2nh}{k} - 3 & \text{if } k \mid 2nh, \\ \left\lfloor \frac{2nh}{k} \right\rfloor - 2 & \text{if } k \nmid 2nh. \end{cases}$$

From (24.1),

$$\begin{aligned} h_{2nt-1} &= \sum_{i=0}^{2nt-1} f_i \\ &= \sum_{i=0}^{i_1-1} f_i + \sum_{h=1}^{tk-1} \sum_{i=i_h}^{i_{h+1}-1} f_i \\ &= \sum_{i=0}^{i_1-1} f_i + \sum_{h=1}^{tk-1} f_{i_h} + \sum_{h=1}^{tk-1} \sum_{i=i_h+1}^{i_{h+1}-1} f_i. \end{aligned}$$

Now our interest falls upon the term $\sum_{i_h+1}^{i_{h+1}-1} f_i$. The number of summands is

$$\begin{aligned} i_{h+1} - 1 - (i_h + 1) + 1 &\geq \left\lfloor \frac{2n(h+1)}{k} \right\rfloor - 4 - \left(\left\lfloor \frac{2nh}{k} \right\rfloor - 1 \right) + 1 \\ &\geq \left\lfloor \frac{2n}{k} \right\rfloor - 2 \end{aligned}$$

which means that we can make the distance between i_h and i_{h+1} arbitrarily large. But Z is independent of n which by the box-principle means that for large enough n , there exists ρ_h and τ_h such that

$$Z_{i_h+\tau_h} = Z_{i_h+\tau_h+\rho_h} \quad (i_h + \tau_h + \rho_h \leq i_{h+1} - 1)$$

where $1 \leq \tau_h \leq Z$, $1 \leq \rho_h \leq Z$, (so then τ_h and ρ_h are both independent of θ since Z is).

Now examine $Z_{\kappa+i_h+\tau_h+j\rho_h}$ where $\kappa + i_h + \tau_h + j\rho_h \leq i_{h+1} - 1$ and $0 \leq \kappa < \rho_h$. Since $\varepsilon_{i_h+\tau_h} = 0$ and $\varepsilon_{i_h+\tau_h+\rho_h} = 0$ we have

$$Z_{i_h+\tau_h+1} = Z_{i_h+\tau_h+\rho_h+1} \quad \text{provided } i_h + \tau_h + \rho_h + 1 \leq i_{h+1} - 1.$$

Inductively,

$$Z_{i_h+\tau_h+u} = Z_{i_h+\tau_h+\rho_h+u} \quad \text{provided } i_h + \tau_h + \rho_h + u \leq i_{h+1} - 1.$$

Hence,

$$Z_{i_h+\tau_h+\kappa} = Z_{i_h+\tau_h+\kappa+j\rho_h} \quad \text{provided } i_h + \tau_h + \kappa + j\rho_h \leq i_{h+1} - 1$$

which implies

$$f_{i_h+\tau_h+\kappa} = f_{i_h+\tau_h+\kappa+j\rho_h}. \quad (24.2)$$

24.1. Counting the partial quotients

We now wish to establish a j which satisfies $i_h + \tau_h + \kappa + j\rho_h \leq i_{h+1} - 1$. Consequently, let

$$v := \left\lfloor \frac{\lfloor 2n/k \rfloor - \tau_h - \rho_h}{\rho_h} \right\rfloor - 1 = \left\lfloor \frac{2n - k\tau_h - k\rho_h}{k\rho_h} \right\rfloor - 1.$$

Then we have,

$$\begin{aligned}
i_h + \tau_h + \kappa + v\rho_h &\leq i_h + \tau_h + \kappa + \rho_h \left(\frac{\lfloor 2n/k \rfloor - \tau_h - \rho_h}{\rho_h} - 1 \right) \\
&= i_h + \lfloor 2n/k \rfloor + \kappa - 2\rho_h \\
&< i_h + \lfloor 2n/k \rfloor - 2 \\
&< i_{h+1}.
\end{aligned}$$

Then

$$\sum_{i=i_h+1}^{i_{h+1}-1} f_i = \sum_{j=1}^{\tau_h-1} f_{i_h+j} + \sum_{\kappa=0}^{\rho_h-1} \left(\sum_{j=0}^v f_{i_h+\tau_h+\kappa+j\rho_h} \right) + \sum_{j=i_h+\tau_h+\rho_h+v\rho_h}^{i_{h+1}-1} f_j$$

The number of terms in $\sum_{j=i_h+\tau_h+\rho_h+v\rho_h}^{i_{h+1}-1} f_j$ is

$$\begin{aligned}
&i_{h+1} - 1 - i_h - \tau_h - \rho_h - v\rho_h + 1 \\
&\leq i_{h+1} - i_h - \tau_h - \rho_h - \rho_h \left(\frac{\lfloor 2n/k \rfloor - \tau_h - \rho_h}{\rho_h} - 1 \right) \\
&= i_{h+1} - i_h - \lfloor 2n/k \rfloor + \rho_h \\
&\leq \rho_h + 2.
\end{aligned}$$

By (24.2),

$$\sum_{\kappa=0}^{\rho_h-1} \sum_{j=0}^v f_{i_h+\tau_h+\kappa+j\rho_h} = (v+1) \sum_{\kappa=0}^{\rho_h-1} f_{i_h+\tau_h+\kappa}.$$

Hence,

$$\sum_{i=i_h+1}^{i_{h+1}-1} f_i = \left\lfloor \frac{2n - k\tau_h - k\rho_h}{k\rho_h} \right\rfloor \zeta_h + \xi_h, \quad (24.3)$$

where ζ_h, ξ_h are independent of θ .

We now take

$$\begin{aligned}
\rho &= \prod_{i=1}^{tk} \rho_h \\
w &= \text{lcm}[k, \mu, \rho]
\end{aligned}$$

both of which are independent of θ . Write $n = w\gamma + \phi$, where $0 \leq \phi < w$.

From the original set I_ν , we now wish to consider the following subset,

$$I_{\nu, \phi} = \{n \in I_\nu : n \equiv \phi \pmod{w}\}$$

In the sequel, we shall suppose that $n \in I_{\nu, \phi}$ (again this loses no generality since every n lies in some $I_{\nu, \phi}$). Consequently, we obtain

$$\left\lfloor \frac{2n - k\tau_h - k\rho_h}{k\rho_h} \right\rfloor = \left\lfloor \frac{2\gamma w + 2\phi - k\tau_h - k\rho_h}{k\rho_h} \right\rfloor = 2\gamma \frac{w}{k\rho_h} + \left\lfloor \frac{2\phi - k\tau_h - k\rho_h}{k\rho_h} \right\rfloor.$$

Thus, the sum (24.3) becomes,

$$\sum_{i=i_h+1}^{i_{h+1}-1} f_i = 2\zeta_h \gamma \frac{w}{k\rho_h} + \zeta_h \left\lfloor \frac{2\phi - k\tau_h - k\rho_h}{k\rho_h} \right\rfloor + \xi_h,$$

which means we can now write h_{2nt-1} as

$$h_{2nt-1} = \sum_{i=0}^{i_1-1} f_i + \sum_{h=1}^{tk-1} f_{i_h} + 2\gamma \sum_{h=1}^{tk} \zeta_h \frac{w}{k\rho_h} + \sum_{h=1}^{tk} \left(\zeta_h \left\lfloor \frac{2\phi - k\tau_h - k\rho_h}{k\rho_h} \right\rfloor + \xi_h \right).$$

Now, let

$$x_t = 2 \sum_{h=1}^{tk} \zeta_h \frac{w}{k\rho_h}$$

$$y_t = \sum_{h=1}^{tk} \left(\zeta_h \left\lfloor \frac{2\phi - k\tau_h - k\rho_h}{k\rho_h} \right\rfloor + \xi_h \right) + \sum_{i=0}^{i_1-1} f_i + \sum_{h=1}^{tk-1} f_{i_h}$$

Both of which are integers and independent of γ . Then,

$$\begin{aligned} h_{2nt-1} &= \gamma x_t + y_t \\ &= ((n - \phi)/w) x_t + y_t \\ &= \frac{x_t}{w} n + y_t - \frac{\phi x_t}{w} \\ &= a_t n + b_t \end{aligned}$$

where a_t, b_t are both rational numbers which are independent of n for all $n \in I_{\nu, \phi}$.

This shows that the length of the expansion up to h_{2nt-1} is linear in n . It remains to show that there exists some h_{2nt-1} where $Q_{h_{2nt-1}} = 1$, and if $Q_j = 1$ then $j = h_{2nt-1}$ for some t independent of n . As noticed in the examples earlier, when square factors are present, it can be difficult determining whether or not the square factors are present in Q_{h_i} . This will be overcome by showing that an appropriate Q_{h_i} must exist, without our explicitly having to find it.

25. FINDING AN ELEMENT OF NORM 1

We now examine the product of the elements in the expansion. First, we recall that

$$\Psi_{h_{i+2}+1} = \frac{\alpha^{1+\varepsilon_i} \beta}{d^2 \Delta_i \Delta_{i+1}} \Psi_{h_i+1}$$

and

$$\Psi_{h_1+1} = \frac{\beta \alpha}{d \Delta_0}.$$

Then inductively,

$$\Psi_{h_{2i-1}+1} = \frac{(\alpha\beta)^i \alpha^{\sum_{j=1}^{i-1} \varepsilon_{2j-1}}}{d^{2i-1} \prod_{j=0}^{2i-2} \Delta_j}.$$

So,

$$\begin{aligned} \Psi_{h_{2nt-1}+1} &= \frac{(\alpha\beta)^{nt} \alpha^{\sum_{j=1}^{nt-1} \varepsilon_{2j-1}}}{d^{2nt-1} \prod_{j=0}^{2nt-2} \Delta_j} \\ &= \frac{(\alpha\beta)^{nt} \alpha^{kt}}{d^{2nt-1} \prod_{j=0}^{2nt-2} \Delta_j}; \end{aligned}$$

and

$$|N(\Psi_{h_{2nt-1}+1})| = |Q_{h_{2nt-1}}| = Q_{h_{2nt-1}}.$$

Hence,

$$Q_{h_{2nt-1}} \left(d^{2nt-1} \prod_{j=0}^{2nt-2} \Delta_j \right)^2 = |N((\alpha\beta)^{nt}) N(\alpha^{kt})| = |N((\alpha\beta)^n) N(\alpha^k)|^t$$

as well as,

$$Q_{h_{2n-1}} \left(d^{2n-1} \prod_{j=0}^{2n-2} \Delta_j \right)^2 = |N((\alpha\beta)^n) N(\alpha^k)|,$$

which implies that

$$(Q_{h_{2n-1}})^t \left(d^{2n-1} \prod_{j=0}^{2n-2} \Delta_j \right)^{2t} = |N((\alpha\beta)^n)N(\alpha^k)|^t.$$

So we get,

$$\frac{\left(d^{2n-1} \prod_{j=0}^{2n-2} \Delta_j \right)^t}{d^{2nt-1} \prod_{j=0}^{2nt-2} \Delta_j} = \sqrt{\frac{Q_{h_{2nt-1}}}{(Q_{h_{2n-1}})^t}} \in \mathbb{Q}.$$

Since $\lambda_{2j-1} \equiv jk \pmod{n}$ we have that $\lambda_{2ni-1} = 0$ for positive i . Hence, $\lambda_{2n-1} = \lambda_{2nt-1} = 0$ and so

$$\begin{aligned} Q_{h_{2n-1}} &= l_{2n-1} m_{2n-1} r_{2n-1} (su_{2n-1} z_{2n-1} y_{2n-1})^2, \\ Q_{h_{2nt-1}} &= l_{2nt-1} m_{2nt-1} r_{2nt-1} (su_{2nt-1} z_{2nt-1} y_{2nt-1})^2. \end{aligned}$$

Thus,

$$\sqrt{\frac{l_{2nt-1} m_{2nt-1} r_{2nt-1}}{(l_{2n-1} m_{2n-1} r_{2n-1})^t}} \in \mathbb{Q}.$$

Since $l_{2n-1}, m_{2n-1}, r_{2n-1}$ are each squarefree and relatively prime, if $2 \nmid t$ then

$$\sqrt{l_{2nt-1} m_{2nt-1} r_{2nt-1}} \in \mathbb{Q} \text{ which implies } l_{2nt-1} = m_{2nt-1} = r_{2nt-1} = 1.$$

Conversely, if $2 \nmid t$, then

$$\sqrt{\frac{l_{2nt-1} m_{2nt-1} r_{2nt-1}}{l_{2n-1} m_{2n-1} r_{2n-1}}} \in \mathbb{Q} \text{ so then } l_{2nt-1} m_{2nt-1} r_{2nt-1} = l_{2n-1} m_{2n-1} r_{2n-1}.$$

So if $l_{2n-1} m_{2n-1} r_{2n-1} \neq 1$ then $N(\Psi_{h_{2n-1}+1}) \neq 1$.

Explicitly, we have

$$|N(\Psi_{h_{2n-1}+1})| = l_{2n-1} m_{2n-1} r_{2n-1} (su_{2n-1} y_{2n-1} z_{2n-1})^2.$$

There a couple of ways of reaching our goal now. On the one hand we might note there are only finitely many possible values of $N(\Psi_{h_{2n-1}+1})$. Further, since each one is a bounded number, they can only occur finitely often in the primitive period. If the expansion defined by this procedure were a regular continued fraction expansion it would then hold that $\theta_{h_{2ni-1}} = \theta_{h_{2nj-1}}$ with i and j independent of n . However, our expansion is not necessarily regular and to make this idea work in this case would require some detailing.

25.1. Constructing an element of norm 1

Instead, we construct an element of norm 1. When $l_{2n-1} = m_{2n-1} = r_{2n-1} = 1$ we just take

$$\Gamma = \frac{\Psi_{h_{2n-1}+1}}{su_{2n-1} y_{2n-1} z_{2n-1}}.$$

Conversely, if $l_{2n-1} m_{2n-1} r_{2n-1} \neq 1$ then

$$N(\Psi_{h_{2n-1}+1}^2) = (l_{2n-1} m_{2n-1} r_{2n-1})^2 (su_{2n-1} y_{2n-1} z_{2n-1})^4.$$

Put,

$$\varepsilon = \begin{cases} 1 & \text{if } l_{2n-1} m_{2n-1} r_{2n-1} \neq 1 \\ 0 & \text{if } l_{2n-1} m_{2n-1} r_{2n-1} = 1 \end{cases}$$

and

$$\Gamma = \frac{\Psi_{h_{2n-1}+1}^{1+\varepsilon}}{(l_{2n-1}m_{2n-1}r_{2n-1})^\varepsilon (su_{2n-1}y_{2n-1}z_{2n-1})^{1+\varepsilon}}$$

Then $N(\Gamma) = 1$.

Lemma 44. *If $D_n = F_n^2 D'_n$ where D'_n is squarefree, then $su_i y_i z_i \mid F_n$.*

Proof. Since $d^2 D_n = c^2 E_n$, we see that if E'_n is the squarefree kernel of E_n then $E'_n = D'_n$. We now observe that

$$d^2 (F_n/c)^2 D'_n \equiv (s_2/c)^2 \pmod{4z^2} \quad \text{and} \quad z \mid s_2/c.$$

Recall that $2d_z = (s_2 - td, 2d)$. If $2 \nmid s_2$ then $t = 1$ and $2 \nmid z$. So

$$z^2 \mid d^2 (F_n/c)^2 \text{ implies } (z/d_z)^2 \mid (d/d_z)^2 (F_n/c)^2.$$

Now suppose that $p \mid (z/d_z, d/d_z)$ with $p > 1$, then $p \mid s_2/(cd_z)$ and

$$p \mid s_2/d_z - td/d_z \text{ which implies } p \mid (s_2/d_z - td/d_z)/2 \text{ because } p \text{ is odd.}$$

But $((s_2/d_z - td/d_z)/2, d/d_z) = 1$ which is a contradiction. Thus $(z/d_z, d/d_z) = 1$ and $z/d_z \mid F_n/c$.

Now suppose that $2 \mid s_2$ and $t = 0$, then $4 \mid D_n$ and $2 \mid F_n$. Also, $d_z = (s_2/2, d)$ and

$$(dF_n/2)^2 D'_n \equiv (s_2/2)^2 \pmod{c^2 z^2}.$$

If $2 \mid d$ then $2 \nmid c$ so $2c \mid F_n$ and

$$\begin{aligned} d^2 (F_n/2c)^2 D'_n &\equiv (s_2/2c)^2 \pmod{z^2} \\ (d/d_z)^2 (F_n/2c)^2 D'_n &\equiv (s_2/2cd_z)^2 \pmod{(z/d_z)^2}. \end{aligned}$$

If $p \mid (d/d_z, z/d_z)$ then $p \mid s_2/2cd_z$ so that $p \mid (s_2/d_z, d/d_z)$, which is impossible. Hence $(d/d_z, z/d_z) = 1$ and $z/d_z \mid F_n/2c$.

If $2 \nmid d$ then $2 \nmid d_z$ and

$$d^2 (F_n/c)^2 D'_n \equiv (s_2/c)^2 \pmod{4z^2}.$$

If $2 \mid s_2/c$ then

$$d^2 (F_n/2c)^2 D'_n \equiv (s_2/2c)^2 \pmod{z^2} \text{ which gives } z/d_z \mid F_n/2c.$$

If $2 \nmid s_2/c$ then

$$(d/d_z)^2 (F_n/c)^2 D'_n \equiv (s_2/cd_z)^2 \pmod{4(z/d_z)^2}.$$

Then, if $p \mid (z/d_z, d/d_z)$ with $p > 1$, we have $p \mid s_2/cd_z$ which implies that $p \mid s_2/2d_z$, which is impossible. Hence $(z/d_z, d/d_z) = 1$ and $z/d_z \mid F_n/c$.

Now, if $2 \mid s_2$ and $t = 1$ then $D_n \equiv 1 \pmod{4}$, thus $2 \mid d$ and $2 \nmid c$. Then since $d_z \mid s_2/2$ and $d_z \mid d/2$, we have

$$\begin{aligned} d^2 (F_n/c)^2 D'_n &\equiv (s_2/c)^2 \pmod{4z^2} \\ (d/2d_z)^2 (F_n/c)^2 D'_n &\equiv (s_2/2cd_z)^2 \pmod{(z/d_z)^2} \end{aligned}$$

And $(z/d_z, d/2d_z) = 1$, so then $z/d_z \mid F_n/c$.

By similar reasoning we can show that $y/d_y \mid F_n/c$. Hence $zy/(d_z d_y) \mid F_n/c$ and $su_i y_i z_i \mid F_n$. ■

25.2. Lucas functions

The lemma and above results imply that we have

$$\Gamma = V_1 + Y_1 \frac{F_n}{su_{2n-1}y_{2n-1}z_{2n-1}} \omega'_n$$

where

$$\omega'_n := \frac{\sigma'_n - 1 + \sqrt{D'_n}}{\sigma'_n} \quad \text{and} \quad \sigma'_n := \begin{cases} 2 & \text{if } D'_n \equiv 1 \pmod{4} \\ 1 & \text{otherwise} \end{cases}$$

and D'_n is the squarefree component of D_n , that is $D_n = F_n^2 D'_n$. If we now define V_j and Y_j by

$$\left(V_j + Y_j \frac{F_n}{su_{2n-1}y_{2n-1}z_{2n-1}} \omega'_n \right) = \Gamma^j.$$

Then Y_j/Y_1 is the Lucas function, $(\Gamma^j - \bar{\Gamma}^j)/(\Gamma - \bar{\Gamma})$ because $\Gamma\bar{\Gamma} = 1$. Furthermore, there must exist some minimal positive p such that

$$su_{2n-1}y_{2n-1}z_{2n-1} \mid Y_p.$$

Letting $g = (1 + \varepsilon)p$ we get

$$\Psi_{h_{2ng-1}+1} = \Gamma^p \sqrt{l_{2ng-1}m_{2ng-1}r_{2ng-1}} su_{2ng-1}y_{2ng-1}z_{2ng-1}.$$

If $\varepsilon = 1$ then $2 \mid g$ implies that $l_{2ng-1}m_{2ng-1}r_{2ng-1} = 1$. And if $\varepsilon = 0$ then $l_{2n-1}m_{2n-1}r_{2n-1} = 1$ so that $l_{2ng-1}m_{2ng-1}r_{2ng-1} = 1$. In either case, $(\Psi_{h_{2ng-1}+1})$ is primitive in $\mathcal{O}_n = \mathbb{Z} + \omega_n \mathbb{Z}$ and

$$\Gamma^p \in \mathbb{Z} + \omega_n \mathbb{Z}.$$

Thus we must also have $su_{2ng-1}y_{2ng-1}z_{2ng-1} = 1$. Hence, $N(\Psi_{h_{2ng-1}+1}) = 1$ which means that $Q_{h_{2ng-1}} = 1$. Furthermore, the values of p depend only on $su_{2n-1}y_{2n-1}z_{2n-1}$ which divides $su_{2n-1}y_{2n-1}z_{2n-1}$. Thus, there can only be a finite number of possible values for p .

Conversely, if $N(\Psi_{h_{2nc-1}+1}) = 1$ for some c then $1 + \varepsilon \mid c$ which means that

$$l_{2nc-1}m_{2nc-1}r_{2nc-1}su_{2nc-1}y_{2nc-1}z_{2nc-1} = 1.$$

And if we put $e = c/(1 + \varepsilon)$ then $\Psi_{h_{2nc-1}+1} = \Gamma^e$, and for some $z \in \mathbb{Z}$ we have

$$\frac{Y_e \omega_n}{su_{2n-1}y_{2n-1}z_{2n-1}} = \omega_n z.$$

This implies $su_{2n-1}y_{2n-1}z_{2n-1} \mid Y_e$, and by the theory of Lucas functions $p \mid e$. Hence $g \mid c$, which means that the first instance of $N(\Psi_{h_{2nc-1}+1}) = 1$ corresponds to the one found above. That is, our solution is fundamental, although this is superfluous in showing that D_n is a creeper.

26. RETURNING TO THE REGULAR CONTINUED FRACTION EXPANSION

There is just one final point which needs to be made in order to show creepiness. Up to now we have determined

$$\omega_n = [a_0, \dots, a_{h_0-1}, b_1, a_{h_0+1}, \dots, a_{h_1-1}, b_2, \dots, a_{h_{2n-1}-1}, 2a_0 - t],$$

but in this evaluation we never insisted that $b_i \geq 1$. In other words this expansion might not correspond to the regular continued fraction expansion of ω_n . This is equivalent to saying that the expansions of A_i/B_i might have an initial nonpositive partial quotient. Of course we know that such partial quotients can be easily removed but in doing so we might destroy the creepiness. It turns out that this doesn't happen.

It is important that the number of possible non-positive partial quotients are bounded independently of n .

Proposition 45. *In the expansion of ω_n given by the earlier procedure, the number of non-positive partial quotients is bounded independently of n .*

Proof. Let $b_i := \lfloor A_i/B_i \rfloor$ and suppose that $b_i < 1$. Then we have either,

$$\text{Case I:} \quad \theta_{h_i} = \frac{\omega_n + S_1/d - E_i x^{n-\lambda_i}/d}{E'_i x^{n-\lambda_i}} = \frac{A_i}{B_i} + e \quad (26.1a)$$

or

$$\text{Case II:} \quad \theta_{h_i} = \frac{\omega_n + S_2/d - E_i x^{\lambda_i}/d}{E'_i x^{\lambda_i}} = \frac{A_i}{B_i} + e \quad (26.1b)$$

Also recall that E_i, E'_i are independent of n . In Case I, (26.1a),

$$A_i = Ax^n - E_i x^{n-\lambda_i} \quad \text{and} \quad B_i = dE'_i x^{n-\lambda_i}.$$

If $b_i < 1$ then

$$Ax^n < (E_i + dE'_i)x^{n-\lambda_i} \quad \text{which implies} \quad x^{\lambda_i} < \frac{E_i + dE'_i}{A}.$$

Thus, $\lambda_i < K_1$ where K_1 is bounded independently of n . In Case II, (26.1b),

$$A_i = Ax^n - E_i x^{\lambda_i} \quad \text{and} \quad B_i = dE'_i x^{\lambda_i}.$$

If $b_i < 1$ then

$$Ax^n < (E_i + dE'_i)x^{\lambda_i} \quad \text{which implies} \quad x^{\lambda_i} > \frac{Ax^n}{(E_i + dE'_i)}.$$

Thus, $\lambda_i > n - K_2$ where K_2 is bounded independently of n . ■

This proposition means that there are only finitely many partial quotients which need to be altered in order to find the regular continued fraction expansion of ω_n .

26.1. Removing nonpositive partial quotients

The removal of nonpositive partial quotients is covered in Dirichlet [14]. Here we recap its details, it entails no more than running through the possible scenarios. Namely, suppose

$$\gamma = [\mu, \nu, p, q, r, \dots]$$

We are going to suppose that $\nu < 1$. The case $\nu = 0$ was treated in Chapter I. Suppose that $\nu < -1$ then we have

$$\begin{aligned} [\mu, \nu, p] &= [\mu - 1, 1, -\nu - 1, -p] \\ [\mu, \nu, p, q] &= [\mu - 1, 1, -\nu - 2, 1, p - 1, q] \end{aligned}$$

and not one of the partial quotients is negative. If $\nu = -1$ and $p > 1$ then we have

$$[\mu, -1, p, q] = [\mu - 2, 1, p - 2, q]$$

and if $\mu > 1$ then all the partial quotients are now non-negative. If $\mu = 1$ then we are in the same situation as when $\nu = -1$ and $p = 1$. There one finds,

$$[\mu, -1, 1, q, r, s] = [\mu - 2 - q, 1, r - 1, s]$$

and all the partial quotients are nonnegative except possibly $\mu - 2 - q$. The point is that the negative partial quotient has been moved to the left. This procedure can be repeatedly used to move any negative partial quotients to the left and keep the size of the partial quotients bounded until the complete quotient $\theta_{h_{i-1}}$ is reached. In other words, the only situation remaining is

$$[b_{i-1}, -1, 1, u, v, \dots] = [b_{i-1} - 2 - u, 1, v - 1, \dots]$$

where $u > 0$ and is bounded independently of n and $b_{i-1} = \lfloor A_{i-1}/B_{i-1} \rfloor$. In Case I, (26.1a), we have

$$\theta_{h_{i-1}} = \frac{\omega_n + S_2/d - E_{i-1}x^{\lambda_i}/d}{E'_{i-1}x^{\lambda_i}} = \frac{A_{i-1}}{B_{i-1}} + e$$

and from before, $\lambda_i < K_1$ which implies

$$\lfloor A_{i-1}/B_{i-1} \rfloor > x^{n-J_1}$$

where J_1 is bounded independently of n . Consequently,

$$b_{i-1} - 2 - u > 0$$

for all sufficiently large n .

Similarly, in Case II, (26.1b), we have

$$\theta_{h_{i-1}} = \frac{\omega_n + S_1/d - E_{i-1}x^{n-\lambda_{i-2}}/d}{E'_{i-1}x^{n-\lambda_{i-2}}} = \frac{A_{i-1}}{B_{i-1}} + e$$

and from before $\lambda_{i-2} > n - k - K_2$ which gives $n - \lambda_{i-2} < J_2$ with J_2 bounded independently of n . Hence,

$$\lfloor A_{i-1}/B_{i-1} \rfloor > x^{n-J_2} \text{ so then } b_{i-1} - 2 - u > 0 \text{ for all sufficiently large } n$$

and the same conclusion follows as before.

Finally, we note that a_{h_i+j} with $j > 0$ can not be at the end of a period because

Lemma 46. *If $B_i > 0$ and $\Delta_i = (A_i, B_i)$ and $A_i/B_i = [q_0, q_1, \dots, q_t]$ then $0 < q_j \leq B_i/\Delta_i$ for $j = 1, \dots, t$*

Proof. This is straightforward from observing $[q_1, \dots, q_t] = \frac{B_i/\Delta_i}{K}$ with $K \leq B_i/\Delta_i$. ■

Hence the lemma gives,

$$a_{h_i+j} < 2a_0 - t.$$

Furthermore, for sufficiently large n , each $b_j \neq b_{h_{2nt-1}}$ satisfies $b_j < x^n$ since $B_i \geq x$.

In conclusion, we have shown that,

$$lp(\omega_n) = h_{2ng-1} + c_n = a_g n + b_g + c_n = a_g n + b'_g.$$

where $c_n \in \mathbb{Z}$ can be bounded independently of n . Then a_g, b'_g are bounded independently of n .

Hence there must exist an infinitude of values of $n \in I$ such that

$$lp(\omega_n) = an + b$$

where $a, b \in \mathbb{Q}$ and are fixed independently of n . Hence the sequence of discriminants $\{D_n\}_{n \in I}$ is a creeper.

Chapter VIII. Higher order creepers

This chapter represents some musing on creepers and jeepers. It connects together the earlier work on exceptional function fields in Chapter III with numerical creepers.

27. MORE CREEPERS

Having completed our investigations of creepers, the next obvious question is what creepers are there which are not creepers? In other words what sequences of discriminants

$$D_n = a_{jn}x^{jn} + a_{(j-1)n}x^{(j-1)n} + \cdots + a_0 \quad a_i \in \mathbb{Q}$$

exist such that

$$lp(\omega_n) = an + b \quad \text{and} \quad R(D_n) = \Theta(n^2)$$

Kreepers are basically the $j = 2$ case with some additional conditions. It would seem a little over the top to just pick a j and attempt to emulate what we did with creepers. Rather we return to the origins of creepers. The basic premise of creepers is the ability to write them as

$$D_n = s_1^2 + 4Ax^n \quad \text{and} \quad D_n = s_2^2 + 4Bx^{n+k}$$

where $x^n | s_1 + s_2$. Returning to the origins of functions with a periodic continued fraction, recall that a particular example is

$$Y^2 = A_1(X)^2 + 4t_1X \quad \text{and} \quad Y^2 = A_2(X)^2 + 4t_2X^2$$

where $X | (A_1(X) + A_2(X))$. The connection between the two is obvious. Also remember that exceptional function fields were constructed by writing down a set of ideals which composed nicely to form a unit. This is exactly the same procedure as finding creepers except that we require a dependence on n , that is, an infinite family of cases.

27.1. An example

Consequently, we take the simplest non-trivial case of torsion in genus one; namely that of torsion five, which is given by

$$Y^2 = (X^2 - (t^2 - 6t + 1))^2 + 32t(X - (t - 1)).$$

We first translate X so that $Y^2 = (X^2 + 2(t - 1)X + 4t)^2 + 32tX$ and our idea then is to replace X by X^n . Some care is required in the selections of t and X so that we don't end up with a numerical sleeper. Anyway, $t = 2$, $X = 2$ will work and after removing a superfluous 2^6 we have

$$D_n = (2^{2n+1} + 2^n + 1)^2 + 4 \cdot 2^n \tag{27.1}$$

where

$$lp(\omega_n) = \begin{cases} 4n + 1 & n \equiv 0 \pmod{2} \\ 8n + 2 & n \equiv 1 \pmod{2} \end{cases}$$

ASIDE: We can also find

$$D_n = (2^{2n+2} + 2^n + 1)^2 + 4 \cdot 2^n$$

This is also a creeper. In fact,

$$lp(\omega) = \begin{cases} 7n + 1 & n \equiv 0 \pmod{2} \\ 14n + 2 & n \equiv 1 \pmod{2} \end{cases}$$

Here is the expansion of ω_n from (27.1) for $n = 14$,

h	a_h	P_h	Q_h	factors of Q_h
0	268443649	0	1	
1	32769	268443648	16384	2^{14}
2	16383	268443647	32770	
3	1	268427262	268451839	
4	1	24576	268435456	2^{28}
5	8191	268410879	65536	2^{16}
6	1	268394496	402628609	
7	3	134234112	134225920	$2^{13} J$
8	134221824	268443647	4	2^2
9	131076	268443648	4096	2^{12}
10	4095	268443647	131080	$2^3 J$
11	1	268328952	469680121	
12	7	201351168	67108864	2^{26}
13	2047	268410879	262144	2^{18}
14	1	268197888	503101441	
15	15	234903552	33556480	$2^{11} J$
16	33555456	268443647	16	2^4
17	524304	268443648	1024	2^{10}
18	1023	268443647	524320	$2^5 J$
19	1	267935712	519618529	
20	31	251682816	16777216	2^{24}
21	511	268410879	1048576	2^{20}
22	1	267411456	527482369	
23	63	260070912	8389120	$2^9 J$
24	8388864	268443647	64	2^6
25	2097216	268443648	256	2^8
26	255	268443647	2097280	$2^7 J$
27	1	266362752	530628481	
28	127	264265728	4194304	2^{22}
29	127	268410879	4194304	2^{22}

where $J = 5 \cdot 29 \cdot 113 = 2^{14} + 1$ and so $lp(\omega_n) = 57$ as expected. The presence of J which is a function of n shows that this is not a creeper, but it might be an extended-creeper.

27.2. Generalising this example

If this example were to be used as the basis for a more general family of creepers one might try to simultaneously satisfy the following relations

$$\begin{aligned}
D_n &= S_1^2 + 4t_1 X^n & X^n | S_1 + S_2 \\
&= S_2^2 + 4t_2 X^n J_n & X^n | S_1 + S_3 \\
&= S_3^2 + 4t_3 X^{3n} & J_n | S_2 + S_4 \\
&= S_4^2 + 4t_4 X^{2n} J_n & X^{2n} | S_3 + S_4
\end{aligned}$$

Writing $S_i = a_i X^{2n} + b_i X^n + c_i$, the condition $X^n | S_1 + S_2$ becomes $c_2 = -c_1$. Similarly,

$$\begin{aligned}
X^n | S_1 + S_3 &\text{ implies } c_3 = -c_1 \\
X^{2n} | S_3 + S_4 &\text{ implies } c_3 = -c_4 \text{ and } b_3 = -b_4
\end{aligned}$$

If we were to suppose that $J_n = X^n + r$ then

$$S_2 + S_4 = (a_2 + a_4)X^{2n} + (b_2 + b_4)X^n = X^n((a_2 + a_4)X^n + b_2 + b_4),$$

so

$$r = \frac{b_2 + b_4}{a_2 + a_4}.$$

Equating coefficients yields, $a_i = a$ for $i = 1, \dots, 4$. And the coefficients of X^{3n} give

$$ab_1 = ab_2 = ab_3 + 2t_3 = 2t_4 - ab_3.$$

Hence, $b_2 = b_1$ and $ab_3 = t_4 - t_3$. The coefficients of X^{2n} yield $ac_1 = t_2$ and $-ac_1 = t_4r$ so $t_2 = -t_4r$.

The coefficients of X^n give $b_1c_1 = t_2r - t_1$. Also,

$$r = \frac{b_2 + b_4}{a_2 + a_4} = \frac{b_2 - b_3}{2a} \text{ so then } b_3 = b_1 - 2ar.$$

From X^{3n} we have

$$ab_1 = ab_3 + 2t_3 = ab_1 - 2a^2r + 2t_3 \text{ thus } t_3 = a^2r.$$

Next,

$$ab_1 = -a(b_1 - 2ar) + 2t_4 \text{ implies } t_4 = ab_1 - 2a^2r + a^2r = a(b_1 - ar).$$

Thus, $t_2 = -ar(b_1 - ar)$. Hence,

$$\begin{aligned} -rb_1(b_1 - ar) &= -r^2a(b_1 - ar) - t_1 \\ t_1 &= rb_1^2 - 2a^2rb_1 + a^2r^3 = r(b_1 - ar)^2. \end{aligned}$$

And no further reductions can be made since all the conditions are now satisfied. Consequently, the four representations have become,

$$\begin{aligned} D_n &= (aX^{2n} + bX^n - r(b - ar))^2 + 4r(b - ar)^2X^n \\ &= (aX^{2n} + bX^n + r(b - ar))^2 - 4ar(b - ar)X^n(X^n + r) \\ &= (aX^{2n} + (b - 2ar)X^n + r(b - ar))^2 + 4a^2rX^{3n} \\ &= (aX^{2n} - (b - 2ar)X^n - r(b - ar))^2 + 4a(b - ar)X^{2n}(X^n + r) \end{aligned}$$

At this point we are not guaranteed to find a unit, in other words we still need to select a, b, r carefully.

If we take $r = -1, a = X - 1, b = -X$ then

$$Y^2 = ((X - 1)X^{2n} - X^{n+1} - 1)^2 - 4X^n \quad (27.2)$$

This function wonderfully displays a folly which is essential not to make. Over the integers, if one failed to remove square factors this would perhaps be silly but certainly not fatal, that is, the period length will not blow up. But in the function field case, square factors are to be avoided since not only can they cause an explosion in the period length they can completely destroy periodicity. In (27.2) there is a factor $(X - 1)^2$ regardless of n . Without removing it, only $n = 1, 2, 3$ will provide a unit with period lengths of 8, 14 and 24 respectively. However, removing $(X - 1)^2$ provides us with

$$W^2 = (X^{2n} - (X^n + X^{n-1} + \dots + 1))^2 - 4X^n(X^{n-1} + X^{n-2} + \dots + 1)$$

whose expansion is

h	$a_h(X)$	$P_h(X)$	$Q_h(X)$
0	$X^{2n} - (X^n + \dots + 1)$	0	1
1	$\frac{-1}{2}(X - 1)$	$X^{2n} - (X^n + \dots + 1)$	$-4X^n(X^{n-1} + \dots + 1)$
2	$2(X^n - 1)$	$X^{2n} - X^n + (X^{n-1} + \dots + 1)$	X^n
3	$\frac{-1}{2}(X^n(X - 1) - 1)$	$X^{2n} - (X^n + \dots + 1)$	$-4(X^{n-1} + \dots + 1)$
4	$2(X^n - 1)$	$X^{2n} - (X^n + \dots + 1)$	X^n
5	$\frac{-1}{2}(X - 1)$	$X^{2n} - X^n + (X^{n-1} + \dots + 1)$	$-4X^n(X^{n-1} + \dots + 1)$
6	$2(X^{2n} - (X^n + \dots + 1))$	$X^{2n} - (X^n + \dots + 1)$	1

Obviously, by instead considering $W^2/4$ we would find a unit in $\frac{1}{2}\mathbb{Z}[X]$ for every n and X . To my understanding, this is the first example of a non-trivial infinite family of function field sleepers.

27.3. Some further examples

Continuing on, we now choose $a = x^2$, $b = x^2 + 1$ and $r = 1$. (Other cases include $b = x^2 - 1$, $x^2 \pm 3$). In other words,

$$D_n = (x^{2n+2} + x^{n+2} + x^n - 1)^2 + 4x^n. \quad (27.3)$$

The other representations become

$$\begin{aligned} D_n &= (x^{2n+2} + (x^2 + 1)x^n + 1)^2 - 4x^{n+2}J_n \\ &= (x^{2n+2} + (1 - x^2)x^n + 1)^2 + 4x^{3n+4} \\ &= (x^{2n+2} - (1 - x^2)x^n - 1)^2 + 4x^{2n+2}J_n \end{aligned}$$

where $J_n = x^n + 1$. The period length is given by

$$lp(\omega_n) = \begin{cases} 2n + 1 & n \equiv 0 \pmod{4} \\ 8n + 10 & n \equiv 1, 3 \pmod{4} \\ 4n + 4 & n \equiv 2 \pmod{4} \end{cases}$$

The expansion for $x = 3$ and $n = 14$ can be found in the Tables at page 150.

Making the following abbreviations

$$\begin{aligned} D_n &= S_1^2 + 4x^n \\ &= S_2^2 - 4x^{n+2}J_n \\ &= S_3^2 + 4x^{3n+4} \\ &= S_4^2 + 4x^{2n+2}J_n \end{aligned}$$

and also $s_i = (S_i - t)/2$, we see that in detail, the expansion proceeds as

$$\begin{array}{lll} \omega & = & s_1 \quad - \quad (\bar{\omega} + s_1) \\ \frac{\omega + s_1}{x^n} & = & x^2(x^n + 1) \quad - \quad \frac{(\bar{\omega} + s_2 - x^n)}{x^n} \\ \frac{\omega + s_2 - x^n}{x^{2n+2} + (1 - x^2)} & = & 1 \quad - \quad \frac{(\bar{\omega} + s_3 - 2x^2)}{x^{2n+2} + (1 - x^2)} \\ \frac{\omega + s_3 - 2x^2}{x^2J_n} & = & x^n - 1 \quad - \quad \frac{(\bar{\omega} + s_4)}{x^2J_n} \\ \frac{\omega + s_4}{x^{2n}} & = & x^2 \quad - \quad \frac{(\bar{\omega} + s_3)}{x^{2n}} \\ \frac{\omega + s_3}{x^{n+4}} & = & x^{n-2} \quad - \quad \frac{(\bar{\omega} + s_4)}{x^{n+4}} \\ \frac{\omega + s_4}{x^{n-2}J_n} & = & x^4 - 1 \quad - \quad \frac{(\bar{\omega} + s_2 - x^{n-2}J_n)}{x^{n-2}J_n} \end{array}$$

$$\begin{aligned}
\frac{\omega + s_2 - x^{n-2}J_n}{S_2 - x^4(x^{n-6}J_n + 1)} &= 1 - \frac{(\bar{\omega} + s_2 - x^4)}{S_2 - x^4(x^{n-6}J_n + 1)} \\
\frac{\omega + s_2 - x^4}{x^4} &= x^{2n-2} + (x^2 + 1)x^{n-4} - 1 - \frac{(\bar{\omega} + s_1)}{x^4} \\
\frac{\omega + s_1}{x^{n-4}} &= x^{n+6} + (x^2 + 1)x^4 - 1 - \frac{(\bar{\omega} + s_2 - x^{n-4})}{x^{n-4}}
\end{aligned}$$

and at this point we don't need to do any more work since the pattern is evident. Indeed the term $(\bar{\omega} + s_2 - x^{n-4})/x^{n-4}$ corresponds to $(\bar{\omega} + s_2 - x^n)/x^n$ in the second line. (Also note that the sequence has 8 terms, hence the 8's that sort of occur in the period lengths). We can immediately conclude that the next section of the expansion will continue as

$$\begin{aligned}
\frac{\omega + s_2 - x^{n-4}}{Q} &= 1 - \frac{(\bar{\omega} + P)}{Q} \\
\frac{\omega + P}{x^6J_n} &= x^{n-4} - 1 - \frac{\bar{\omega} + s_4}{x^6J_n} \\
\frac{\omega + s_4}{x^{2n-4}} &= x^6 - \frac{(\bar{\omega} + s_3)}{x^{2n-4}} \\
\frac{\omega + s_3}{x^{n+8}} &= x^{n-6} - \frac{(\bar{\omega} + s_4)}{x^{n+8}} \\
\frac{\omega + s_4}{x^{n-6}J_n} &= x^8 - 1 - \frac{\bar{\omega} + s_2 - x^{n-6}J_n}{x^{n-6}J_n} \\
\frac{\omega + s_2 - x^{n-6}J_n}{Q'} &= 1 - \frac{(\bar{\omega} + s_2 - x^8)}{Q'} \\
\frac{\omega + s_2 - x^8}{x^8} &= x^{2n-6} + (x^2 + 1)x^{n-8} - 1 - \frac{(\bar{\omega} + s_1)}{x^8}
\end{aligned}$$

where $Q = x^{2n+2} - (x^{10} - x^6 - x^4 + 1)x^{n-4} + (1 - x^6)$ and P, Q' are equally terrible.

These awful P 's and Q 's are a bit annoying, but they only arise because of the minus sign in the representation

$$D = S_2^2 - 4x^{n+2}J_n.$$

Consequently, a simpler approach would be to ignore the minus sign, at the expense of some negative partial quotients. Taking this approach gives the expansion as

$$\begin{aligned}
\omega &= s_1 - (\bar{\omega} + s_1) \\
\frac{\omega + s_1}{x^n} &= x^{n+2} + x^2 + 1 - \frac{(\bar{\omega} + s_2)}{x^n} \\
\frac{\omega + s_2}{-x^2J_n} &= -x^n - \frac{(\bar{\omega} + s_4)}{-x^2J_n} \\
\frac{\omega + s_4}{-x^{2n}} &= -x^{n-2} - \frac{(\bar{\omega} + s_3)}{-x^{2n}} \\
\frac{\omega + s_3}{-x^{n+4}} &= -x^{n-2} - \frac{(\bar{\omega} + s_4)}{-x^{n+4}} \\
\frac{\omega + s_4}{-x^{n-2}J_n} &= -x^4 - \frac{(\bar{\omega} + s_2)}{-x^{n-2}J_n} \\
\frac{\omega + s_2}{x^4} &= x^{2n-2} + (x^2 + 1)x^{n-4} - \frac{(\bar{\omega} + s_1)}{x^4} \\
\frac{\omega + s_1}{x^{n-4}} &= x^4\alpha
\end{aligned}$$

where $\alpha = \omega + s_1/x^n$ represents the end of the sequence. Then we can see that

$$\begin{aligned}\omega_n &= [s_1, x^{n+2} + x^2 + 1, 0, -1, 1, -1, 0, x^n, x^2, x^4, \\ &\quad - (x^{2n-2} + (x^2 + 1)x^{n-4}), -x^4\alpha] \\ &= [s_1, x^{n+2} + x^2, 1, x^n - 1, x^2, x^{n-2}, x^4, \\ &\quad 0, -1, 1, -1, 0, x^{2n-2} + (x^2 + 1)x^{n-4}, x^4\alpha] \\ &= [s_1, x^{n+2} + x^2, 1, x^n - 1, x^2, x^{n-2}, x^4 - 1, \\ &\quad 1, x^{2n-2} + (x^2 + 1)x^{n-4} - 1, x^4\alpha].\end{aligned}$$

Furthermore, we can easily determine $x^4\alpha$ by multiplying the expansion of α by x^4 .

$$\begin{aligned}x^4\alpha &= [x^{n+6} + x^4(x^2 + 1), -x^{n-4}, -x^6, -x^{n-6}, -x^8, \\ &\quad x^{2n-6} + (x^2 + 1)x^{n-8}, x^8\alpha] \\ &= [x^{n+6} + x^4(x^2 + 1) - 1, 1, x^{n-4} - 1, 1, x^6, x^{n-6}, x^8 - 1, \\ &\quad 1, x^{2n-6} + (x^2 + 1)x^{n-8} - 1, x^8\alpha].\end{aligned}$$

When $n \equiv 0 \pmod{4}$ we have $Q_n = x^{n/2}$ and $Q_{n+1} = x^{n-n/2} = Q_n$ and hence the period length is $2n + 1$.

When $n \equiv 2 \pmod{4}$ then $Q_{2n+2} = J_n$ and $J_n|D$ implies this is halfway in the expansion, so $lp(\omega_n) = 4n + 4$.

When $n \equiv 1, 3 \pmod{4}$ we have $Q_{4n+5} = J_n$, so that $lp(\omega_n) = 8n + 10$.

27.4. Explanation of this creeper

Detailing the expansion makes it apparent why we are obtaining a unit. Indeed using the previous four representations for D we can write down the following four ideals

$$\begin{aligned}\mathfrak{a} &= \langle x^n, s_1 \rangle & x^n|s_1 + s_2 \\ \mathfrak{b} &= \langle -x^{n+2}J_n, s_2 \rangle & x^{n+2}J_n|s_2 + s_4 \\ \mathfrak{c} &= \langle x^{3n+4}, s_3 \rangle & x^{2n+2}|s_3 + s_4 \\ \mathfrak{d} &= \langle x^{2n+2}J_n, s_4 \rangle\end{aligned}$$

So, starting from the ideal $\mathfrak{a}$ we proceed to compose with the following sequence of ideals $\mathfrak{b}, \mathfrak{d}, \mathfrak{c}, \mathfrak{d}, \mathfrak{b}, \mathfrak{a}$ and then repeat the sequence until either a unit is obtained or the ideal has norm J_n ; the square of this ideal will then have norm 1.

We could also consider our form as a curve and look at the continued fraction expansion in the function field. Taking $n = 1$, we discover

h	$a_h(X)$	$P_h(X)$	$Q_h(X)$
0	$X^4 + X^3 + X - 1$	0	1
1	$\frac{1}{2}X^3 + \frac{1}{2}X^2 + \frac{1}{2}$	$X^4 + X^3 + X - 1$	$4X$
2	$-2X$	$X^4 + X^3 + X + 1$	$-X^2(X + 1)$
3	$\frac{-1}{2}X^2 - \frac{1}{2}X$	$X^4 + X^3 - X - 1$	$-4X^2$
4	$2X^2$	$X^4 + X^3 + X + 1$	$X(X + 1)$
5	$\frac{1}{2}X + \frac{1}{2}$	$X^4 + X^3 - X - 1$	$4X^3$
6	$-2X^3 - 2$	$X^4 + X^3 + X + 1$	$-(X + 1)$
7	$\frac{1}{2}X + \frac{1}{2}$	$X^4 + X^3 + X + 1$	$4X^3$
8	$2X^2$	$X^4 + X^3 - X - 1$	$X(X + 1)$
9	$\frac{-1}{2}X^2 - \frac{1}{2}X$	$X^4 + X^3 + X + 1$	$-4X^2$
10	$-2X$	$X^4 + X^3 - X - 1$	$-X^2(X + 1)$

$$\left[\begin{array}{c|c|c} 11 & \frac{1}{2}X^3 + \frac{1}{2}X^2 + \frac{1}{2} & X^4 + X^3 + X + 1 \\ 12 & 2X^4 + 2X^3 + 2X - 2 & X^4 + X^3 + X - 1 \end{array} \right] \begin{array}{c} 4X \\ 1 \end{array}$$

giving a genus 3 curve with regulator 25. Of course, the expansion does not match that with which we originally created the creeper but that is to be expected because we have changed rational numbers into variables, hence changing the curve.

Allowing n to run, we find:

$$\begin{aligned} n \equiv 1 \pmod{2}, & \quad lp(Y) = 6(n+1), & \quad R(Y) = 5n^2 + 12n + 8, \\ n \equiv 2 \pmod{4}, & \quad lp(Y) = 3n + 2, & \quad R(Y) = (5n^2 + 12n + 8)/2, \\ n \equiv 0 \pmod{4}, & \quad lp(Y) = (3n + 2)/2, & \quad R(Y) = (5n^2 + 12n + 8)/4. \end{aligned}$$

One might notice that if we take our standard creeper form

$$D_b = (qrx^b + mx^k - l)^2 + 4rlx^b$$

(here n has been replaced by b in order to avoid confusion). Then by taking $r = X^n + 1$, $l = -1$, $x^k = X^n$, $m = q = 1$, $x^b = X^{n+2}$ we obtain

$$D_n = ((X^n + 1)X^{n+2} + X^n + 1)^2 - 4X^{n+2}(X^n + 1)$$

or equivalently,

$$D_n = (X^{2n+2} + (X^2 + 1)X^n - 1)^2 + 4X^n$$

matching with (27.3), which the language introduced in §22 on page 98, says is an extended-kreeper. It would take us too far afield to show that every extended-kreeper does indeed creep. But it should at least be clear that if $l, m \in \mathbb{Q}[X]$ are independent of n , then the previous procedures apply and that in this case an extended-kreeper does creep. Hence, it would not have been necessary in this particular case to detail the full expansion in order to verify creepiness.

Why was this example an extended-kreeper? It happens that the original curve selected was not complicated enough. On the surface it did not seem to be a function field creeper, but if we take a closer look we see

$$\begin{aligned} Y^2 &= (X^2 + X + 2)^2 + 8X \\ &= X^4 + 2X^3 + 5X^2 + 12X + 4 \\ &= ((X + 2)X - X - 2)^2 + 4 \cdot 2 \cdot X(X + 2) \end{aligned}$$

which is a function field creeper with the selections of $q = 1$, $r = X + 2$, $n = k = 1$, $l = 2$, $m = -1$. So it comes as no surprise that the new creeper is an extended-kreeper. Thus, if we wish to come up with substantially new examples of creepers then perhaps we should start with even more complicated exceptional function fields.

28. FROM TORSION TO CREEPERS

The elliptic curve with torsion 7 can be represented as

$$Y^2 = (X^2 - (t^4 - 6t^3 + 3t^2 + 2t + 1))^2 + 32t^2(t - 1)(X - (t^2 - t - 1))$$

but we want the remainder to be a monomial so we translate and dilate to find

$$Y^2 = (X^2 + (t^2 - t - 1)X + t^2(t - 1))^2 + 4t^2(t - 1)X$$

As before, we now replace X by X^n , but somehow t needs to be selected appropriately. It helps to notice the following equations,

$$\begin{aligned} Y^2 &= (X^{2n} + (t^2 - t - 1)X^n - t^2(t - 1))^2 + 4t^2(t - 1)X^n(X^n + t(t - 1)) \\ &= (X^{2n} + (t^2 - 3t + 1)X^n - t^2(t - 1))^2 + 4(t - 1)X^n(X^n + t(t - 1))^2 \end{aligned}$$

These equations did not magically appear, §8 explains their existence. We could now examine the continued fraction expansion of Y over $\mathbb{Q}[X]$ and find suitable t 's, but it is easier to note that we are hoping to find a unit by composing the ideals coming from the above equations in some manner. If we call the ideals,

$$\begin{aligned}\mathfrak{a} &= \langle t^2(t-1)X^n, Y + X^{2n} + (t^2 - t - 1)X^n + t^2(t-1) \rangle \\ \mathfrak{b} &= \langle t^2(t-1)X^n(X^n + t(t-1)), Y + X^{2n} + (t^2 - t - 1)X^n - t^2(t-1) \rangle \\ \mathfrak{c} &= \langle (t-1)X^n(X^n + t(t-1))^2, Y + X^{2n} + (t^2 - 3t + 1)X^n - t^2(t-1) \rangle\end{aligned}$$

then in order for $\mathfrak{a}$ and $\mathfrak{b}$ to compose well we need,

$$t^2(t-1) \mid (X^{2n} + (t^2 - t - 1)X^n).$$

In other words,

$$X^n(X^n + t^2 - t - 1) \equiv 0 \pmod{t-1}$$

which is

$$X^n(X^n - 1) \equiv 0 \pmod{t-1}.$$

Either $X^n \equiv 0 \pmod{t-1}$ or $X^n \equiv 1 \pmod{t-1}$. On the other hand,

$$X^n(X^n + t^2 - t - 1) \equiv 0 \pmod{t}$$

gives

$$X^n(X^n - 1) \equiv 0 \pmod{t}.$$

Either $X^n \equiv 0 \pmod{t}$ or $X^n \equiv 1 \pmod{t}$. From the factorization of $X^n - 1$ we must have

$$X^n - 1 = (t-1)f(X) \quad \text{and} \quad X^n = tg(X),$$

so that $t = X^m$ and $m \mid n$. Taking $t = X$ we find

$$Y^2 = (X^{2n} + (X^2 - X - 1)X^n + X^2(X-1))^2 + 4X^2(X-1)X^n$$

which has a factor of $2^4 X^4$. It turns out to be better to leave the 2's in, so then,

$$Y^2 = (X^{2n+2} + (X^2 - X - 1)X^n + (X-1))^2 + 4(X-1)X^n \quad (28.1)$$

as well as,

$$\begin{aligned}&= (X^{2n+2} + (X^2 - X - 1)X^n - (X-1))^2 + 4(X-1)X^{n+1}(X^{n+1} + X - 1) \\ &= (X^{2n+2} + (X^2 - 3X + 1)X^n - (X-1))^2 + 4(X-1)X^n(X^{n+1} + X - 1)^2\end{aligned}$$

where the fact that $X \mid t^2(t-1)$ has been essential in order to obtain the exponent of $n+1$ in the second line; without it, there would be no way the expansion could ever creep. Over $\mathbb{Q}[X]$ the continued fraction expansion of Y satisfies:

$$lp(Y) = 12n, \quad R(Y) = 7n^2 + 9n + 3.$$

The following representations also exist,

$$\begin{aligned}Y^2 &= (X^{2n+2} + (X^2 + X - 1)X^n + (X-1))^2 - 4X^n J_n K_n \\ &= (X^{2n+2} - (X^2 - X + 1)X^n - (X-1))^2 + 4X^{2n+1}(X-1)K_n \\ &= (X^{2n+2} + (X^2 - X + 1)X^n + (X-1))^2 - 4X^{2n+1}J_n\end{aligned}$$

where $K_n = (X^{n+2} + X - 1) \mid Y^2$ and $J_n = X^{n+1} + X - 1$. But these are not necessary in order to demonstrate the existence of a unit. Here is the expansion for $n = 5$,

h	$a_h(X)$	$P_h(X)$	$Q_h(X)$
0	$X^{12} + X^7 - X^6 - X^5 + X - 1$	0	1
1	$\frac{1}{2}(X+1)(X^5 + X^3 + X + 1)$	$P_1(X)$	$4X^5(X-1)$
2	$2X^5$	$P_2(X)$	XJ_5
3	$\frac{-1}{2}X^2$	$P_3(X)$	$-4X^{10}$
4	$-2(X^3 + X^2 + X + 1)$	$P_4(X)$	$-X(X-1)K_5$
5	$\frac{-1}{2}(X-1)$	$P_5(X)$	$Q_1(X)$
6	$-2(X+1)$	$P_6(X)$	$-X^4(X-1)J_5$
7	$\frac{-1}{2}X^5$	$P_7(X)$	$-4XJ_5$
8	$2X$	$P_8(X)$	X^4K_5
9	$\frac{1}{2}(X^4 + X^3 + X^2 + X + 1)$	$P_4(X)$	$4X^7(X-1)$
10	$2(X-1)$	$P_9(X)$	$Q_2(X)$
11	$\frac{1}{2}(X+1)$	$P_{10}(X)$	$4X^4(X-1)J_5$
12	$2X^3(X-1)(X+1)(X^5 + X^3 + X + 1)$	$P_2(X)$	X^2
13	$\frac{1}{2}X^2(X+1)(X^5 + X^3 + X + 1)$	$P_1(X)$	$4X^3(X-1)$
14	$2X^3$	$P_2(X)$	X^3J_5
15	$\frac{-1}{2}X^4$	$P_3(X)$	$-4X^8$
16	$-2(X+1)$	$P_4(X)$	$-X^3(X-1)K_5$
17	$\frac{-1}{2}(X-1)$	$P_{11}(X)$	$Q_3(X)$
18	$-2(X^3 + X^2 + X + 1)$	$P_{12}(X)$	$-X^2(X-1)J_5$
19	$\frac{-1}{2}X^3$	$P_7(X)$	$-4X^3J_5$
20	$2X^3$	$P_8(X)$	X^2K_5
21	$\frac{1}{2}(X^2 + X + 1)$	$P_4(X)$	$4X^9(X-1)$
22	$2(X-1)$	$P_{13}(X)$	$Q_4(X)$
23	$\frac{1}{2}(X^3 + X^2 + X + 1)$	$P_{14}(X)$	$4X^2(X-1)J_5$
24	$2X(X-1)(X+1)(X^5 + X^3 + X + 1)$	$P_2(X)$	X^4
25	$\frac{1}{2}X^4(X+1)(X^5 + X^3 + X + 1)$	$P_1(X)$	$4X(X-1)$
26	$2X$	$P_2(X)$	X^5J_5
27	$\frac{-1}{2}J_5$	$P_3(X)$	$-4X^6$
28	$-2(X^5 + X^4 + X^3 + X^2 + X + 1)$	$P_2(X)$	$-(X-1)J_5$
29	$\frac{-1}{2}X$	$P_7(X)$	$-4X^5J_5$
30	$2(X^5 + 1)$	$P_8(X)$	K_5

where

$$\begin{aligned}
P_1(X) &= X^{12} + X^7 - X^6 - X^5 + X - 1 \\
P_2(X) &= X^{12} + X^7 - X^6 - X^5 - X + 1 \\
P_3(X) &= X^{12} + X^7 - X^6 + X^5 + X - 1 \\
P_4(X) &= X^{12} - X^7 + X^6 - X^5 - X + 1 \\
P_5(X) &= X^{12} - 2X^8 + X^7 + X^6 - X^5 - 2X^2 + 3X - 1 \\
P_6(X) &= X^{12} - 2X^{10} + X^7 + X^6 - 3X^5 + 2X^4 + X - 1 \\
P_7(X) &= X^{12} + X^7 - 3X^6 + X^5 - X + 1 \\
P_8(X) &= X^{12} + X^7 + X^6 - X^5 + X - 1 \\
P_9(X) &= X^{12} - X^7 - X^6 + X^5 + X - 1 \\
P_{10}(X) &= X^{12} - 2X^{10} + X^7 - X^6 - X^5 + 2X^4 + X - 1 \\
P_{11}(X) &= X^{12} - 2X^{10} + X^7 + X^6 - X^5 - 2X^4 + 2X^3 + X - 1 \\
P_{12}(X) &= X^{12} - 2X^8 + X^7 + X^6 - X^5 - 2X^3 + 2X^2 + X - 1 \\
P_{13}(X) &= X^{12} - 2X^9 + X^7 - X^6 + X^5 + X - 1
\end{aligned}$$

$$P_{14}(X) = X^{12} - 2X^8 + X^7 - X^6 + X^5 - 2X^3 + 2X^2 + X - 1$$

$$Q_1(X) = -4(X^{11} + X^{10} - X^7 + X^5 - X^4 - X + 1$$

$$Q_2(X) = X^{11} + X^{10} - X^5 - X^4 + 1$$

$$Q_3(X) = -4(X^{11} + X^{10} - X^7 + X^5 - X^3 - X^2 + 1)$$

$$Q_4(X) = X^{11} + X^{10} + X^9 - X^7 - X^5 - X^2 + 1$$

and $J_n = X^{n+1} + X - 1$ and $K_n = X^{n+2} + X - 1$ which divides Y^2 for all n . Furthermore, the fact that $(X^{n+1} + X - 1) \nmid Y^2$ implies that this creeper can not be a creeper since we find Q_h 's in the expansion which are not of the shape pX^v with $p|Y^2$. This fact also suggests that it is not an extended-kreeper, but of this we can not be sure.

The line containing $Q_{12} = X^2$, which is independent of n , represents the end of the first "cycle", hence the 12 in the period length. Upon numerating, we find that the period length of ω_n is $14n + 2$.

28.1. Inserting constant factors

The next idea is to attempt to place some nice constant factors in the remainders in a similar manner to the r , l and m in creepers. That is,

$$\begin{aligned} D_n &= S_1^2 + 4rk(x-1)x^n \\ &= S_2^2 + 4rl(x-1)x^{n+1}(x^{n+1} + x - 1) \\ &= S_3^2 + 4rm(x-1)x^n(x^{n+1} + x - 1)^2 \end{aligned} \quad (28.2)$$

In the earlier general case we have the factorization,

$$(x^{n+1} + x - 1)^2 - 1 = (x^n + 1)(x^{n+2} + x(x - 2))$$

and the two factors $x^n + 1$ and $x^{n+2} + x(x - 2)$ split off to $S_1 - S_3$ and $S_1 + S_3$ respectively. From (28.2),

$$\begin{aligned} S_1^2 - S_3^2 &= 4r(x-1)x^n (m(x^{n+1} + x - 1)^2 - k) \\ S_1 + S_3 &= 2qrx^n f(x) \\ S_1 - S_3 &= 2(x-1)g(x)/q \end{aligned}$$

where $f(x)g(x) = m(x^{n+1} + x - 1)^2 - k$. The difficulty now is that $f(x)g(x)$ must factorize in a non-trivial manner. In other words, the degree of $f(x)$ needs to be of order n so that $S_1 + S_3$ is of order $2n$ and the degree of $g(x)$ also needs to be of order n so that $S_1 - S_3$ is of order n . First observe that if $m = k$, so that we can use the earlier factorization, then we obtain nothing new. In this situation, the equations become

$$\begin{aligned} S_1 + S_3 &= 2qrmx^n (x^{n+2} + x(x - 2)) \\ S_1 - S_3 &= 2(x-1)(x^n + 1)/q \end{aligned}$$

which implies

$$\begin{aligned} S_1 &= qrmx^{2n+2} + (qrm(x^2 - 2x) + (x-1)/q)x^n + (x-1)/q, \\ S_3 &= qrmx^{2n+2} + (qrm(x^2 - 2x) - (x-1)/q)x^n - (x-1)/q. \end{aligned}$$

And,

$$\begin{aligned} S_1^2 - S_2^2 &= 4r(x-1)x^n (lx(x^{n+1} + x - 1) - m) \\ S_1 + S_2 &= 2q_2rx^n (lx(x^{n+1} + x - 1) - m) \\ S_1 - S_2 &= 2(x-1)/q_2 \end{aligned}$$

which imply that

$$\begin{aligned} S_1 &= q_2 r l x^{2n+2} + q_2 r (l x^2 - l x - m) x^n + (x - 1)/q_2 \\ S_2 &= q_2 r l x^{2n+2} + q_2 r (l x^2 - l x - m) x^n - (x - 1)/q_2 \end{aligned}$$

Equating coefficients on S_1 gives $q = q_2$, $l = m$, whereupon,

$$q r (l x^2 - l x - l) = q r l (x^2 - 2x) + (x - 1)/q$$

Hence $q^2 r l = 1$, and taking $r = -1$, $l = -1$ yields the same solution as $r = l = 1$. Moreover, it is clear that Y is invariant under $q \mapsto -q$.

28.2. Determining the constant factors

If $m(x^{n+1} + x - 1)^2 - k$ with $m, k \in \mathbb{Z}$ and $(m, k) = 1$ is to factorize in a non-trivial manner then it is necessary that $m, k \in \mathbb{Z}^2$. Denoting m by a^2 and k by b^2 we have

$$a^2(x^{n+1} + x - 1)^2 - b^2 = (ax^{n+1} + ax - (a + b))(ax^{n+1} + ax - (a - b)).$$

Then,

$$\begin{aligned} S_1 + S_3 &= 2q r x^n (ax^{n+1} + ax - (a + b)), \\ S_1 - S_3 &= 2(x - 1)/q (ax^{n+1} + ax - (a - b)). \end{aligned}$$

so,

$$S_1 = q r a x^{n+1} + (a x (x - 1)/q + q r a x - q r (a + b)) x^n + (a x - (a - b))(x - 1)/q.$$

Furthermore,

$$\begin{aligned} S_1 + S_2 &= 2q_2 r x^n (l x (x^{n+1} + x - 1) - b^2) \\ S_1 - S_2 &= 2(x - 1)/q_2 \end{aligned}$$

so,

$$S_1 = r q_2 l x^{2n+2} + q_2 r (l x (x - 1) - b^2) x^n + (x - 1)/q_2.$$

Equating coefficients, we find $a q_1 = l q_2 x$ and $a(b - a) = l x$, hence $l = a(a - b)/x$. Thus, we must have $x | a - b$. This means $x | S_1 - S_3$. This means we should instead be considering

$$\begin{aligned} S_1 + S_3 &= 2q_1 r x^{n+1} (ax^{n+1} + ax - (a + b)) \\ S_1 - S_3 &= 2(x - 1)/q_1 (ax^n + a - (a - b)/x) \end{aligned}$$

so

$$S_1 = q_1 r a x^{2n+2} + (q_1 r a x^2 - q_1 r x (a + b) + a(x - 1)/q_1) x^n + (a - (a - b)/x)(x - 1)/q_1.$$

Furthermore,

$$\begin{aligned} S_1 + S_2 &= 2q_2 r x^n (l x (x^{n+1} + x - 1) - b^2) \\ S_1 - S_2 &= 2(x - 1)/q_2 \end{aligned}$$

so

$$S_1 = r q_2 l x^{2n+2} + q_2 r (l x (x - 1) - b^2) x^n + (x - 1)/q_2.$$

Equating coefficients gives

$$r q_2 l = q_1 r a \quad \text{and} \quad q_1 = q_2 (a - (a - b)/x) \quad \text{implies} \quad l = a(a - (a - b)/x).$$

For simplicity, let us suppose that $q_1 = q_2 = 1$, then $l = a$ and $a - (a - b)/x = 1$. Equating the coefficient of x^n gives

$$r(1 - a)x^2 + (2ra - (1 + r))x + (1 - ar) = 0,$$

ignoring the solution $a = r = 1$ which corresponds to our earlier solution then we find $x = (1 - ar)/r(1 - a)$, which is never an integer. Hence it is impossible for both q_1 and q_2 to be equal to 1; in other words, $x \neq 2$. Relaxing the conditions on

q_i , we find a set of equations which are unsolvable for small values and most likely, are impossible.

On the other hand, there are also creepers which look like the above example but aren't quite. For example,

$$\begin{aligned} D_n &= (2 \cdot 3^{2n+2} + 5 \cdot 3^n + 1)^2 + 4 \cdot 3^n \\ &= (2 \cdot 3^{2n+2} + 5 \cdot 3^n - 1)^2 + 4 \cdot 3^{n+1} J_n \\ &= (2 \cdot 3^{2n+2} + 3^n - 1)^2 + 4 \cdot 3^n J_n^2 \end{aligned}$$

where $J_n = 2(3^{n+1} + 1)$.

29. UNUSUAL REPRESENTATIONS OF KREEPERS

As a slight detour, for the moment let us return to the original motivation of creepers, which was provided by Shanks. It is interesting to observe that the example highlighted by Shanks, 2089, which is a creeper via

$$D_n = (3 \cdot 2^n + (-2^3 - 1)/3)^2 + 4 \cdot 2^3 \quad (29.1)$$

can also be discovered in the following different manner. In the case 2089 we have

$$2089 = 43^2 + 4 \cdot 2^2 \cdot 15 = 45^2 + 4 \cdot 2^4 = 47^2 - 4 \cdot 2 \cdot 15$$

We generalize this to be

$$D_s = S_1^2 + 4 \cdot 2^a(2^s - 1) = S_2^2 + 4 \cdot 2^b = S_3^2 - 4 \cdot 2^c(2^s - 1)$$

where

$$S_2 - S_1 = S_3 - S_2 = 2.$$

Then one finds that we must have

$$\begin{aligned} D_s &= (2^{s+2} - 2^s - 2^2 - 1)^2 + 4 \cdot 2^2(2^s - 1) \\ &= (2^{s+2} - 2^s - 2^2 + 1)^2 + 4 \cdot 2^s \\ &= (2^{s+1} + 2^s - 2 + 1)^2 - 4 \cdot 2(2^s - 1) \end{aligned}$$

which as it is presented appears nothing like a creeper (also observe that $2^s - 1$ does not divide D_s). Yet from the above three equations a unit can be easily composed. Thankfully our results show that it *must* be a creeper and it is easy to rediscover (29.1). But this is more than just a cute example. It suggests that perhaps the results of §28 could be rephrased somehow to appear more like creepers. This suggestion is also reinforced from the fact that in a creeper there can only ever be one independent ideal of fixed norm. Returning to the curve given in (28.1), it turns out that the equation to study is not the reduced equation, but rather,

$$Y^2 = (X^{2b+2} + (X^2 + X - 1)X^b + (X - 1))^2 - 4X^b J_b K_b \quad (29.2)$$

this suggests considering $l = X^{b+1} + X - 1$ and $m = X^{b+2} + X - 1$. Whereupon one finds that by also taking $r = X - 1$, $q = 1$, $n = b + 1$ and $k = b$ then the standard creeper form,

$$Y^2 = (qrX^n + (mX^k + l)/q)^2 - 4mlX^k/q^2$$

and (29.2) coincide. But with this selection l does not divide Y^2 and hence this is not an extended-creeper. However, this realisation does suggest that further study of extended-creepers is warranted. Furthermore, the remark about there being only one independent ideal suggests the question as to whether every creeper can be represented in the usual form, that is

$$D_n = (qrX^n + (mX^k - l)/q)^2 + 4lrx^n,$$

but where q, r, l, m are functions of x and x^n , not necessarily dividing D_n .

30. LARGE TORSION CREEPERS

We can try to emulate the procedure of §28 with the other cases of torsion; remembering that, except for some isolated cases, these curves failed to satisfy the Schinzel condition. Hence, it can only be in these isolated cases that we could possibly hope to find a creeper since by taking $n = 1$ and varying X in a creeper we have a numerical sleeper. For torsion 10, by taking $t = X$ and dividing by X^6 , we can find the simple examples

$$E_n = (2^{2n+3} - 5 \cdot 2^n + 3)^2 - 12 \cdot 2^n$$

with period lengths:

$$lp(\omega_n) = \begin{cases} 20n - 8 & n \equiv 0 \pmod{6}, \\ 60n - 28 & n \equiv 2, 4 \pmod{6}. \end{cases}$$

And

$$D_n = (3^{2n+3} - 31 \cdot 3^n + 10)^2 + 40 \cdot 3^n,$$

this time with period lengths:

$$lp(\omega_n) = \begin{cases} 23n - 1 & n \equiv 9 \pmod{12}, \\ 69n - 35 & n \equiv 1, 5 \pmod{12}. \end{cases}$$

It is not surprising to find $t = 2, 3$ being the only obvious cases which work since $|t^2 - 3t + 1| = 1$ in these cases, and this was the term which prevented the Schinzel condition from being satisfied.

The expansion of

$$D_{21} = (3^{45} - 31 \cdot 3^{21} + 10)^2 + 40 \cdot 3^{21}$$

is included in the Tables on page 151 and the reader is recommended to have a look since it is quite a remarkable example, of a far more complicated manner than any creepers.

Among the Q_h 's of interest, the following terms arise:

$$I_n = 3^{n+1} - 2, \quad J_n = 3^{n+2} - 10, \quad K_n = 3^{n+3} - 10,$$

and none of these divide D_n . In other words

$$\left(\frac{D_n}{I_n}\right) = \left(\frac{D_n}{J_n}\right) = \left(\frac{D_n}{K_n}\right) = 1.$$

In the case $n = 21$ the only small factors of D_n are 2, 5 and 23. The presence of the norms $x^i I_n, x^j J_n, x^k K_n$, where each of I_n, J_n, K_n split, identifies this creeper as being distinct from any previous examples. The equations of interest are the following:

$$\begin{aligned} D_n &= (3^{2n+3} - 31 \cdot 3^n + 10)^2 + 4 \cdot 2 \cdot 5 \cdot 3^n \\ &= (3^{2n+3} - 31 \cdot 3^n - 10)^2 + 4 \cdot 2 \cdot 5 \cdot 3^{n+1} \cdot J_n \\ &= (3^{2n+3} - 29 \cdot 3^n + 10)^2 - 4 \cdot 3^{2n+1} \cdot J_n \\ &= (3^{2n+3} + 29 \cdot 3^n - 10)^2 - 4 \cdot 2 \cdot 5 \cdot 3^{2n+1} \cdot K_n \\ &= (3^{2n+3} - 49 \cdot 3^n + 10)^2 + 4 \cdot 3^n \cdot J_n \cdot K_n \\ &= (3^{2n+3} - 11 \cdot 3^n - 10)^2 - 4 \cdot 2 \cdot 5 \cdot 3^n \cdot I_n \cdot J_n \\ &= (3^{2n+3} - 25 \cdot 3^n + 10)^2 - 4 \cdot 3^n \cdot I_n \cdot K_n \\ &= (3^{2n+3} + 5 \cdot 3^n - 10)^2 - 4 \cdot 2 \cdot 3^{n+1} \cdot I_n \cdot K_n \\ &= (3^{2n+3} - 41 \cdot 3^n + 10)^2 + 4 \cdot 5 \cdot 3^{n+1} \cdot I_n^2 \end{aligned}$$

30.1. *The pattern of the expansion*

The pattern of the continued fraction expansion is formed by taking each of the ideals formed by the above equations and composing them precisely in the order above. (that is, if the ideals were labelled $\mathfrak{a}, \dots, \mathfrak{i}$ then the sequence $\mathfrak{a}, \mathfrak{b}, \dots, \mathfrak{h}, \mathfrak{i}, \mathfrak{h}, \dots, \mathfrak{a}$ will return to another element with a norm bounded independently of n). Further, since I_n, J_n, K_n each do not ramify, they can not be ignored as in earlier examples.

The set of ideals formed from the above equations will produce the unit; an interesting question is whether this set is minimal. Or, which other sets of ideals will suffice to construct the unit. Since I_n, J_n , and K_n don't ramify it would seem likely that at least four ideals are required.

30.2. *Generalizing this example*

Another question is whether this is an example of a more general family. The fact that the underlying function field sleeper fails the Schinzel condition would suggest that this is unlikely but does not prove it.

It is not at all difficult to satisfy the following set of equations

$$\begin{aligned} D_n &= S_1^2 + 4t_1x^n \\ &= S_2^2 + 4t_2x^{n+1}J_n \\ &= S_3^2 + 4t_3x^{2n+1}K_n \\ &= S_4^2 + 4t_4x^{2n+1}J_n \\ &= S_5^2 + 4t_5x^nJ_nK_n \end{aligned}$$

Indeed,

$$\begin{aligned} D_n &= (x^{2n+3} - (x^3 + x + 1)x^n + x^2 + 1)^2 + 4(x^2 + 1)x^n \\ &= (x^{2n+3} - (x^3 + x + 1)x^n - (x^2 + 1))^2 + 4(x^2 + 1)x^{n+1}J_n \\ &= (x^{2n+3} + (x^3 + x - 1)x^n - (x^2 + 1))^2 - 4(x^2 + 1)x^{2n+1}K_n \\ &= (x^{2n+3} - (x^3 + x - 1)x^n + (x^2 + 1))^2 - 4x^{2n+1}J_n \\ &= (x^{2n+3} - (x^3 + 2x^2 + x + 1)x^n + (x^2 + 1))^2 + 4x^nJ_nK_n \end{aligned}$$

where we have taken $J_n = x^{n+2} - (x^2 + 1)$ and $K_n = x^{n+3} - (x^2 + 1)$ (in fact these selections are suggested by the above equations). But the difficulty lies in finding an appropriate I_n . In other words, it is not obvious how to simultaneously satisfy the following equations

$$\begin{aligned} D_n &= S_1^2 + 4t_1x^{n+1}I_n^2 \\ &= S_2^2 + 4t_2x^nI_nJ_n \\ &= S_3^2 + 4t_3x^nI_nK_n \\ &= S_4^2 + 4t_4x^{n+1}I_nK_n \end{aligned}$$

In the situation of $x = 3$, we have

$$S_1^2 - S_4^2 = 4 \cdot 3^{n+1}I_n(t_4K_n - t_1I_n)$$

and $t_4K_n - t_1I_n = 23 \cdot 3^{n+1} - 30$. This extra factor of 3 is vital since

$$\begin{aligned} S_1 - S_4 &= -2(23 \cdot 3^n - 10) \\ S_1 + S_4 &= 2 \cdot 3^{n+2} I_n \end{aligned}$$

Seemingly, if we are to extend the $x = 3$ example further we will require

$$x | t_4K_n - t_1I_n$$

If we suppose that the t 's are factors of $x^2 + 1$ then after some fiddling we find

$$I_n = x^{n+1} - 2 \quad \text{and} \quad K_n = x^{n+3} - (3x^2 - 2x - 1)/2.$$

The above condition becomes $I_n \equiv 4 \pmod{x}$ whereupon $0 \equiv 6 \pmod{x}$ leaving only $x = 2$ and $x = 3$ as the possibilities.

31. JEEPERS?

Beyond creepers, very little is known. It is suspected that there are no other polynomially parametrized families of discriminants whose period length can be explicitly given. That is, we expect the regulators to be exponentially large. A jeeper would be an infinite sequence of discriminants $\{D_n\}$ which are polynomially parametrized and whose regulator $R(D)$ is of order $\Theta((\log D_n)^3)$. By the results in Chapter V, any sequence of discriminants possessing 2 independent ideals would have a regulator at least this large. In other words, we seek 3 ideals which compose to form a unit. The following example is a simple instance of the sort of thing which we desire.

Example 11. Consider $D = (2^{18} - 2)^2 - 3$. Then we also have

$$D = (2^{18} - 1)^2 - 4 \cdot 2^{17} \quad \text{and} \quad D = 260825^2 + 4 \cdot 2^{23} 3^{16}.$$

From these equations we can write the forms

$$\mathcal{B} = (2^2 3^{16}, 260825, -1) \quad \text{and} \quad \mathcal{C} = (2^{17}, 262143, 1)$$

We can't immediately write a form from $D = (2^{18} - 2)^2 - 3$, but it only takes a moment to find the principal form $\mathcal{A} = (3, 262141, -43690)$. Then,

$$\mathcal{A}^{16} \mathcal{B} = (4, 260825, -43046721) =: \mathcal{D}.$$

And,

$$\mathcal{D}^8 \mathcal{C} = (2, 262143, 2^{16})$$

so

$$(\mathcal{D}^8 \mathcal{C})^2 \mathcal{D} = (1, 260825, 2^{23} 3^{14}).$$

In other words, $\mathcal{A}^{48} \mathcal{B}^{17} \mathcal{C}^2 = (1, 260825, 2^{23} 3^{14})$.

The first set of compositions, between $\mathcal{A}$ and $\mathcal{B}$, is to find a form with a coefficient which is a small power of 2. In the expansion of ω we find that $Q_{87} = 4$. Then the form $\mathcal{C}$ is introduced in order to form the unit.

Only the first few lines of the expansion of ω are included in the Tables, at page 156, since it has a period length of 2111. Also, D has a class number of 163.

31.1. Further examples

Some other interesting examples similar to the one above are:

$$\begin{aligned} D &= (2^{20} + 42)^2 - 2 \cdot 3^4 2^{20} & \text{and} & & D &= (2^{18} + 15)^2 - 4 \cdot 3 \cdot 2^{18} \\ &= (2^{20} - 39)^2 + 3^5 & & & &= (2^{18} + 9)^2 + 2^4 3^2 \\ &= (2^{20} - 11930)^2 + 64 \cdot 3^{18} & & & &= (2^{18} - 137)^2 + 16 \cdot 3^{14} \end{aligned}$$

with period lengths (for ω) of 2521 and 874 respectively. They also have class numbers of 166 and 152. Also,

$$\begin{aligned} D &= (2^{16} + 14)^2 - 2 \cdot 2^{16} & \text{and} & & D &= (2^{23} + 62)^2 - 2^5 2^{23} \\ &= (2^{16} + 13)^2 + 3^3 & & & &= (2^{23} + 46)^2 + 2^6 3^3 \\ &= (2^{16} - 770)^2 + 64 \cdot 3^{13} & & & &= (2^{23} - 50609)^2 + 3^{25} \end{aligned}$$

have period lengths of 1148 and 2470 respectively; the respective class numbers are 32 and 1232.

Notice that the class numbers are smaller than what would be expected of a creeper for a similarly sized discriminant. Now suppose that we intend to use these examples for the basis of a jeeper. If we wish to compose three ideals/forms and produce a unit then we need to find 3 equations of the form $P^2 + 2^a 3^b$. It is easy to discover

$$D_n = (2^n + \frac{1}{4}(2^a 3^b + 2^{a'} 3^{b'}))^2 - 2^a 3^b 2^n \quad (31.1)$$

$$= (2^n - \frac{1}{4}(2^a 3^b + 2^{a'} 3^{b'}))^2 + 2^{a'} 3^{b'} 2^n \quad (31.2)$$

$$= (2^n + \frac{1}{4}(2^{a'} 3^{b'} - 2^a 3^b))^2 + 2^{a+a'-2} 3^{b+b'} \quad (31.3)$$

In fact, all of the above examples are of this shape. However, the ideals arising from these three equations are not independent, in the sense of Chapter V. Observe that the ideal from (31.3) is really only the composite of the ideals from (31.1) and (31.2). The above examples also have an extra equation of the form,

$$D = S^2 + 2^l 3^m \quad (31.4)$$

where m is slightly less than n . The equation, (31.4), is the mystery in these examples. Certainly, it seems necessary to have such an equation in order to remove powers of 3 from the norms involved. However, there appears little one can do to predict its existence; moreover as n increases the number of such examples decreases. It seems likely that these examples are just happenstance occurring for relatively small discriminants.

But there are also discriminants of the shape (31.1)–(31.3) with small period lengths which don't possess an equation of the type (31.4). For example:

$$(2^{27} + 3^8(3^5 + 2^9))^2 - 2^{11} 3^8 2^{27}, \quad (2^{23} + 3^{14})^2 - 2 \cdot 3^{14} 2^{23}, \quad (2^{30} + 3^{11}(1 + 3^6))^2 - 4 \cdot 3^{11} 2^{30}$$

with period lengths: 6842, 1631, and 7429 respectively. It would be interesting to discover why these examples have such small period lengths. Without another equation of the shape (31.4) the ideas used in the example can not be applied.

Ultimately, there does not appear any way in which the examples discussed here can be extended to include large discriminants, let alone an infinite sequence of such discriminants. It seems that there are no jeepers.

Conclusion

32.1. Results

The main result of this thesis is the determination of all creepers. Recall here that a ‘creeper’ is a sequence of discriminants $\{D_n\}$ of real quadratic orders which satisfy:

$$D_n = A^2 x^{2n} + Bx^n + C^2 \quad A, B, C \in \mathbb{Q}$$

with period length $lp(\omega_n) = an + b$ for some $a, b \in \mathbb{Q}$, and for which there exists an element in the principal cycle of ω_n which has norm x^g for some g , fixed independently of n .

We determine that a creeper must be given by

$$d^2 D_n = c^2 \left[(qrx^n + (mz^2 x^k - ly^2)/q)^2 + 4rly^2 x^n \right]$$

where: r, l, m are squarefree, q and r are positive, and

$$(rq, lymzx) = 1, \quad (ly, mz) = 1, \quad rly^2 mz^2 \mid d^2 D_n.$$

Conversely, such a sequence of discriminants is indeed a creeper.

We also discuss which quartics $f(X)$, with $\sqrt{f(X)}$ having a periodic continued fraction, possess a congruence class of values meeting the Schinzel condition. Doing that is equivalent to determining quartics $g(x)$ such that the period length of $\sqrt{g(x)}$, where the expansion is over the integers, is bounded. We also show that it is impossible, in general, for quartics with regulator 10 or 12 to satisfy the Schinzel condition.

A concluding result is the demonstration of the interplay between exceptional function fields and creepers. Using known cases of exceptional functional fields, we construct several instances of creepers which appear to be substantially different from creepers.

32.2. Future directions

For creepers, the remaining obstacles are the assumptions that A and C are squares, and that there exists an element with norm x^g in the principal cycle. We have resisted the *very* strong urge to give some wishy-washy justification for these assumptions. But we feel that these assumptions are true and hope that they will be proved sometime in the near future.

As for creepers beyond the known creepers, prior to this work there was nothing known. The information which we have given here is very brief, but it suggests that knowing exceptional function fields may help in determining more instances of creepers. Mind you, we are certainly not suggesting that this is the only way to produce creepers.

Since all quartics $f(X)$ with periodic continued fraction, $\sqrt{f(X)}$, are known, it seems feasible that one might be able to determine all creepers of the shape

$$D_n = ax^{4n} + bx^{3n} + cx^{2n} + dx^n + e.$$

Along these lines, it would certainly be of interest to know whether the incidental examples given in §30,

$$\begin{aligned} D_n &= (2^{2n+3} - 5 \cdot 2^n + 3)^2 - 12 \cdot 2^n \\ E_n &= (3^{2n+3} - 31 \cdot 3^n + 10)^2 + 40 \cdot 3^n \end{aligned}$$

are truly isolated as alleged, or whether they belong to some larger family.

A creeper would be an infinite sequence of polynomially parametrized discriminants $\{D_n\}$ which satisfy $lp(\omega_n) = \Theta(n^2)$ and $R(D_n)\Theta(n^3)$. Here we say “would

be” emphatically, since there as yet are no known instances. Rather than the numerical case, it may well be easier to deal with the question whether function field jeeppers exist. Doing that seems to require new information on possible torsion divisors on (the Jacobians of) hyperelliptic curves.

Our final remark is to point out that, given an isolated example of a discriminant with large class number, it remains difficult to determine whether that discriminant may be viewed as belonging to a family of creepers or not. If we consider Shanks’ problem as being the classification of discriminants with large class numbers, then we still have some work to do. However, we hope to have provided a little more insight into the issue of discriminants with large class numbers.

Tables

SLEEPERS

$$Y^2 = (2X^3 + X^2 - 4X + 2)^2 - 8X^3(X - 1)^2$$

$a_h(X)$	$P_h(X)$	$Q_h(X)$
$2X^3 - X^2 - 2$	0	1
$\frac{1}{4}X + \frac{1}{8}$	$2X^3 - X^2 - 2$	$16X^2 - 16X$
$8X + 8$	$2X^3 - X^2 - 2X + 2$	$\frac{1}{2}X^2 - \frac{3}{4}X + \frac{1}{2}$
$\frac{-1}{8}X^2 + \frac{1}{16}X$	$2X^3 - X^2 + 2$	$-32X$
$-8X^2 - 4X - 4$	$2X^3 - X^2 - 2$	$\frac{-1}{2}X + \frac{1}{2}$
$\frac{1}{4}X + \frac{1}{4}$	$2X^3 - X^2$	$16X^2 - 24X + 8$
$4X - 2$	$2X^3 - X^2 - 4X + 2$	X^2
$\frac{-1}{4}X$	$2X^3 - X^2 + 4X - 2$	$-16X^2 + 8X$
$-4X - 2$	$2X^3 - X^2 - 4X + 2$	$-X^2 + X$
$\frac{1}{2}X - \frac{1}{2}$	$2X^3 - X^2 + 2X - 2$	$8X^2 + 4X - 8$
$X + \frac{3}{2}$	$2X^3 - X^2 - 8X + 6$	$4X^2 - 8X + 4$
$-2X^2$	$2X^3 - X^2$	$-2X + 1$
$X + \frac{3}{2}$	$2X^3 - X^2$	$4X^2 - 8X + 4$
$\frac{1}{2}X - \frac{1}{2}$	$2X^3 - X^2 - 8X + 6$	$8X^2 + 4X - 8$
$-4X - 2$	$2X^3 - X^2 + 2X - 2$	$-X^2 + X$
$\frac{-1}{4}X$	$2X^3 - X^2 - 4X + 2$	$-16X^2 + 8X$
$4X - 2$	$2X^3 - X^2 + 4X - 2$	X^2
$\frac{1}{4}X + \frac{1}{4}$	$2X^3 - X^2 - 4X + 2$	$16X^2 - 24X + 8$
$-8X^2 - 4X - 4$	$2X^3 - X^2$	$\frac{-1}{2}X + \frac{1}{2}$
$\frac{-1}{8}X^2 + \frac{1}{16}X$	$2X^3 - X^2 - 2$	$-32X$
$8X + 8$	$2X^3 - X^2 + 2$	$\frac{1}{2}X^2 - \frac{3}{4}X + \frac{1}{2}$
$\frac{1}{4}X + \frac{1}{8}$	$2X^3 - X^2 - 2X + 2$	$16X^2 - 16X$
$4X^3 - 2X^2 - 4$	$2X^3 - X^2 - 2$	1

$$Y^2 = X^6 - 2X^5 - 4X^4 + 2X^3 + \frac{37}{4}X^2 - \frac{15}{2}X + \frac{9}{4}$$

$a_h(X)$
$X^3 - X^2 - \frac{5}{2}X - \frac{3}{2}$
$\frac{-2}{15}X^2 + \frac{2}{15}X + \frac{1}{3}$
$5X$
$\frac{-2}{25}X + \frac{2}{25}$
$\frac{25}{3}X^2 - \frac{125}{6}$
$\frac{-6}{125}X + \frac{6}{125}$
$\frac{125}{9}X$
$\frac{-9}{125}X + \frac{9}{50}$
$\frac{-100}{27}X - \frac{250}{27}$
$\frac{-27}{100}X + \frac{27}{50}$
$\frac{200}{81}X + \frac{200}{81}$
$\frac{-81}{500}X^2 + \frac{81}{250}X + \frac{81}{1000}$
$\frac{-1000}{243}X - \frac{1000}{243}$

$\frac{-243}{2500}X + \frac{243}{1250}$
$\frac{2500}{729}X + \frac{1250}{729}$
$\frac{-729}{625}X + \frac{6561}{1250}$
$\frac{-125}{2187}X - \frac{500}{2187}$
$\frac{-4374}{125}X + \frac{4374}{125}$
$\frac{125}{8748}X - \frac{125}{34992}$
$\frac{-46656}{625}X + \frac{81648}{625}$
$\frac{-625}{93312}X - \frac{625}{31104}$
$\frac{-20736}{625}X + \frac{55296}{625}$
$\frac{125}{2304}X + \frac{125}{13824}$
$\frac{-1024}{125}X + \frac{1536}{125}$
$\frac{-125}{1024}X - \frac{125}{512}$
$\frac{-2048}{625}X + \frac{6144}{625}$
$\frac{-625}{3072}X^2 - \frac{625}{1536}X - \frac{4375}{6144}$
$\frac{-2048}{3125}X + \frac{6144}{3125}$
$\frac{-3125}{1024}X - \frac{3125}{512}$
$\frac{-1024}{9375}X + \frac{512}{1875}$
$\frac{1875}{256}X + \frac{1875}{512}$
$\frac{-256}{1875}X + \frac{512}{1875}$
$\frac{-625}{128}X - \frac{1875}{128}$
$\frac{-128}{3125}X + \frac{64}{625}$
$\frac{3125}{96}X - \frac{3125}{192}$
$\frac{-48}{3125}X + \frac{24}{3125}$
$\frac{625}{18}X^2 - \frac{625}{36}X - \frac{6875}{72}$
$\frac{-18}{625}X + \frac{9}{625}$
$\frac{250}{27}X - \frac{125}{27}$
$\frac{27}{250}X + \frac{27}{250}$
$\frac{500}{81}X - \frac{2000}{81}$
$\frac{81}{2500}X + \frac{243}{5000}$
$\frac{10000}{243}X + \frac{5000}{243}$
$\frac{-81}{10000}X + \frac{27}{2500}$
$\frac{-4000}{9}X - \frac{52000}{27}$
$\frac{-1}{8000}X + \frac{19}{32000}$
$\frac{128000}{7}X - \frac{96000}{49}$
$\frac{-343}{11520000}X + \frac{833}{34560000}$
$\frac{207360000}{2401}X - \frac{1105920000}{2401}$
$\frac{2401}{5598720000}X + \frac{26411}{11197440000}$
$\frac{13436928000}{2401}X - \frac{20155392000}{2401}$
$\vdots$

EASY KREEPERS

$$D_n = \left(7 \cdot 2 \cdot 67^n + \frac{67-11}{7}\right)^2 + 4 \cdot 2 \cdot 11 \cdot 67^n$$

For $n = 6$,

h	a_h	P_h	Q_h	factors of Q_h
0	633208675188	0	1	
1	1	633208675188	723667057343	
2	1	90458382155	542750293034	
3	3	452291910879	361833528619	
4	1718341045	633208674978	737	(11)(67)
5	469	633208675187	2700250214	(2)(67) ⁵
6	282115694	633208675179	4489	(67) ²
7	2856	633208675187	443324662	(2)(11)(67) ⁴
8	1	632926559485	805722449401	
9	1	172795889916	460573994231	
10	1	287778104315	690740235002	
11	3	402962130687	345389967859	
12	382789	633207772890	3308393	(11)(67) ³
13	2105341	633208675187	601526	(2)(67) ³
14	62846	633208675179	20151121	(67) ⁴
15	12823440	633208675187	98758	(2)(11)(67) ²
16	1	633208612333	806024648881	
17	1	172816036548	460392674543	
18	1	287576637995	691264047434	
19	3	403687409439	344281903099	
20	85	629158299858	14851376177	(11)(67) ⁵
21	9450875749	633208675187	134	(2)(67)
22	14	633208675179	90458382169	(67) ⁶
23	57564425017	633208675187	22	
24	14	633208675187	90458382169	(67) ⁶
25	9450875749	633208675179	134	(2)(67)
26	85	633208675187	14851376177	(11)(67) ⁵
27	3	629158299858	344281903099	
28	1	403687409439	691264047434	
29	1	287576637995	460392674543	
30	1	172816036548	806024648881	
31	12823440	633208612333	98758	(2)(11)(67) ²
32	62846	633208675187	20151121	(67) ⁴
33	2105341	633208675179	601526	(2)(67) ³
34	382789	633208675187	3308393	(11)(67) ³
35	3	633207772890	345389967859	
36	1	402962130687	690740235002	
37	1	287778104315	460573994231	
38	1	172795889916	805722449401	
39	2856	632926559485	443324662	(2)(11)(67) ⁴
40	282115694	633208675187	4489	(67) ²
41	469	633208675179	2700250214	(2)(67) ⁵
42	1718341045	633208675187	737	(11)(67)
43	3	633208674978	361833528619	
44	1	452291910879	542750293034	

45	1	90458382155	723667057343	
46	1266417350376	633208675188	1	

$$D_n = \left(2 \cdot 5 \cdot 43^n + \frac{43^3 - 7}{2}\right)^2 + 4 \cdot 5 \cdot 7 \cdot 43^n$$

For $n = 11$,

h	a_h	P_h	Q_h
0	4646468697356133413	0	1
1	2	4646468697356133413	4646468697355994276
2	16697428968001	4646468697355855139	$7 \cdot 43^3$
3	159014	4646468697356133410	$5 \cdot 43^8$
4	1470084430	4646468697356093660	43^6
5	1806103728	4646468697356133410	$5 \cdot 7 \cdot 43^5$
6	3	4646468695886009230	2655196768442306089
7	2	3319121609440909037	3982041264480596084
8	2641	4644960919520283131	$7 \cdot 43^9$
9	1005185223873690	4646468697356133410	$5 \cdot 43^2$
10	3	4646468697356130640	2787881218413679201
11	2	3717174957884906963	2787881218413680276
12	1	1858587478942453589	6505056176298585949
13	6174709232366954	4646468697356132360	$5 \cdot 7 \cdot 43$
14	430	4646468697356133410	43^{10}
15	543637222214	4646468697356093660	$5 \cdot 43^4$
16	4884001	4646468697356133410	$7 \cdot 43^7$
17	2	4646467881900260339	3982687299551731076
18	3	3318906717203201813	2655124232124394801
19	976800	4646465979169982590	$5 \cdot 7 \cdot 43^7$
20	2718186111070	4646468697356133410	43^4
21	86	4646468697356093660	$5 \cdot 43^{10}$
22	30873546161834773	4646468697356133410	$7 \cdot 43$
23	2	4646468697356133263	4646468697356132876
24	5025926119368451	4646468697356132489	43^2
25	528	4646468697356133410	$5 \cdot 7 \cdot 43^9$
26	3	4641442771236725230	2653688991047520889
27	2	3319624201905837437	3983046449410452884
28	9030518641	4646468696915068331	$7 \cdot 43^5$
29	294016886	4646468697356133410	$5 \cdot 43^6$
30	795070	4646468697356093660	43^8
31	3339485793600	4646468697356133410	$5 \cdot 7 \cdot 43^3$
32	3	4646468697355298590	2787881218413429601
33	2	3717174957884990213	2787881218413707876
34	1	1858587478942425539	$7 \cdot 43^{11}$
35	1858587478942453364	4646468697356133410	5
36	1	4646468697356133410	$7 \cdot 43^{11}$
37	2	1858587478942425539	2787881218413707876
38	3	3717174957884990213	2787881218413429601
39	3339485793600	4646468697355298590	$5 \cdot 7 \cdot 43^3$
40	795070	4646468697356133410	43^8
41	294016886	4646468697356093660	$5 \cdot 43^6$
42	9030518641	4646468697356133410	$7 \cdot 43^5$
43	2	4646468696915068331	3983046449410452884

44	3	3319624201905837437	2653688991047520889
45	528	4641442771236725230	$5 \cdot 7 \cdot 43^9$
46	5025926119368451	4646468697356133410	43^2
47	2	4646468697356132489	4646468697356132876
48	30873546161834773	4646468697356133263	$7 \cdot 43$
49	86	4646468697356133410	$5 \cdot 43^{10}$
50	2718186111070	4646468697356093660	43^4
51	976800	4646468697356133410	$5 \cdot 7 \cdot 43^7$
52	3	4646465979169982590	2655124232124394801
53	2	3318906717203201813	3982687299551731076
54	4884001	4646467881900260339	$7 \cdot 43^7$
55	543637222214	4646468697356133410	$5 \cdot 43^4$
56	430	4646468697356093660	43^{10}
57	6174709232366954	4646468697356133410	$5 \cdot 7 \cdot 43$
58	1	4646468697356132360	6505056176298585949
59	2	1858587478942453589	2787881218413680276
60	3	3717174957884906963	2787881218413679201
61	1005185223873690	4646468697356130640	$5 \cdot 43^2$
62	2641	4646468697356133410	$7 \cdot 43^9$
63	2	4644960919520283131	3982041264480596084
64	3	3319121609440909037	2655196768442306089
65	1806103728	4646468695886009230	$5 \cdot 7 \cdot 43^5$
66	1470084430	4646468697356133410	43^6
67	159014	4646468697356093660	$5 \cdot 43^8$
68	16697428968001	4646468697356133410	$7 \cdot 43^3$
69	2	4646468697355855139	4646468697355994276
70	9292937394712266826	4646468697356133413	1

WHEN l IS NEGATIVE

$$D_n = \left(11 \cdot 7 \cdot 131^n + \frac{131 - (-23)}{11} \right)^2 + 4 \cdot 7 \cdot (-23) \cdot 131^n$$

For $n = 6$

h	a_h	P_h	Q_h	factors
0	194575656054823	0	1	
1	1	194575656054822	353773920099703	
2	10	159198264044880	35377392009943	
3	129157421874	194575656054549	3013	$(23)(131)$
4	1	194575656051812	388881255679081	
5	1440	194305599627268	270056427557	$(7)(131)^5$
6	22676493916	194575656054811	17161	$(131)^2$
7	1	194575656037664	389103897605209	
8	8206	194528241567544	47414487281	$(7)(23)(131)^4$
9	2	194555041060341	169187259713027	
10	3	143819478365712	101518539876539	
11	2	160736141263904	118437271018451	
12	1	76138400772997	270714020858353	
13	7526216	194575620085355	51706093	$(23)(131)^3$
14	1	194575604348732	389151244666921	
15	24729000	194575640318188	15736637	$(7)(131)^3$
16	1321396	194575656054811	294499921	$(131)^4$

17	1	194575361554904	389151014846809	(7)(23)(131) ²
18	140847787	194575653291904	2762921	
19	3	194575655213922	118439032972643	
20	3	160741443704006	101502637292669	
21	1	143766468174000	169363959562661	(23)(131) ⁵
22	1	25597491388660	219671614078009	
23	437	194074122689348	887328261973	
24	1	193688327792852	388263983846761	
25	424374386160	194575656053908	917	(7)(131)
26	76	194575656054811	5053913144281	(131) ⁶
27	1	189521742910544	384097398965209	(131) ⁶
28	2417088895089	194575656054664	161	
29	1	194575656054664	384097398965209	
30	76	189521742910544	5053913144281	
31	424374386160	194575656054811	917	(7)(131)
32	1	194575656053908	388263983846761	(23)(131) ⁵
33	437	193688327792852	887328261973	
34	1	194074122689348	219671614078009	
35	1	25597491388660	169363959562661	
36	3	143766468174000	101502637292669	(7)(23)(131) ²
37	3	160741443704006	118439032972643	
38	140847787	194575655213922	2762921	
39	1	194575653291904	389151014846809	
40	1321396	194575361554904	294499921	(131) ⁴
41	24729000	194575656054811	15736637	(7)(131) ³
42	1	194575640318188	389151244666921	(23)(131) ³
43	7526216	194575604348732	51706093	
44	1	194575620085355	270714020858353	
45	2	76138400772997	118437271018451	
46	3	160736141263904	101518539876539	(7)(23)(131) ⁴
47	2	143819478365712	169187259713027	
48	8206	194555041060341	47414487281	
49	1	194528241567544	389103897605209	
50	22676493916	194575656037664	17161	(131) ²
51	1440	194575656054811	270056427557	(7)(131) ⁵
52	1	194305599627268	388881255679081	(23)(131)
53	129157421874	194575656051812	3013	
54	10	194575656054549	35377392009943	
55	1	159198264044880	353773920099703	
56	389151312109645	194575656054822	1	

WHEN BOTH l AND m ARE NOT UNITS

$$D_n = (2^n - 5 \cdot 2^2 - 11)^2 + 4 \cdot 11 \cdot 2^n$$

For $n = 26$,

h	a_h	P_h	Q_h	factors
0	33554428	0	1	(2) ² (5)(11)
1	305040	33554427	220	
2	3	33554372	16777201	
3	1	16777230	50331646	
4	4194302	33554415	16	(2) ⁴

5	1	33554416	46137344	$(2)^{22}(11)$
6	2	12582927	20971505	
7	4	29360082	12583034	
8	1	20972053	54523621	
9	19064	33551567	3520	$(2)^6(5)(11)$
10	4	33553712	13631341	
11	1	20971651	50331764	
12	11	29360112	5242880	$(2)^{20}(5)$
13	1	28311567	61865759	
14	262143	33554191	256	$(2)^8$
15	23	33554416	2883584	$(2)^{18}(11)$
16	3	32768015	18087479	
17	1	21494421	36704366	
18	2	15209944	24371956	
19	1191	33533967	56320	$(2)^{10}(5)(11)$
20	4	33543152	13432621	
21	1	20187331	53479604	
22	203	33292272	327680	$(2)^{16}(5)$
23	1	33226767	66777119	
24	16383	33550351	4096	$(2)^{12}$
25	372	33554416	180224	$(2)^{14}(11)$
26	2	33488911	24371956	
27	1	15255000	36648046	
28	3	21393045	18233911	
29	74	33308687	901120	$(2)^{14}(5)(11)$
30	4	33374192	13386541	
31	1	20171971	53710004	
32	3275	33538032	20480	$(2)^{12}(5)$
33	1	33533967	67022879	
34	1023	33488911	65536	$(2)^{16}$
35	5957	33554416	11264	$(2)^{10}(11)$
36	1	33545231	54780581	
37	4	21235349	12321146	
38	2	28049234	27525041	
39	4	27000847	14417920	$(2)^{18}(5)(11)$
40	4	30670832	12845101	
41	1	20709571	54262964	
42	52427	33553392	1280	$(2)^8(5)$
43	1	33553167	66059039	
44	63	32505871	1048576	$(2)^{20}$
45	95325	33554416	704	$(2)^6(11)$
46	15	33554383	4194301	
47	1	29360131	62914484	
48	838859	33554352	80	$(2)^4(5)$
49	1	33554367	50331599	
50	3	16777231	16777216	$(2)^{24}$
51	1525201	33554416	44	$(2)^2(11)$
52	13421771	33554427	5	

$$D_n = (4 \cdot 11^n + 7 \cdot 11^2 - 3)^2 + 4 \cdot 4 \cdot 3 \cdot 11^n$$

In order to be a creeper this needs $n \equiv 3 \pmod{12}$. Here, $n = 15$,

h	a_h	P_h	Q_h
0	8354496338831726	0	1
1	1	8354496338831726	16708992677660912
2	6575754694081	8354496338829186	$(3)(7)(11)^2$
3	2	8354496338830635	7180724125937543
4	3	6006951913044451	4695088851574909
5	17	8078314641680276	$(2)^2(7)(11)^{13}$
6	1141246682444	8354496338830880	$(11)^4$
7	4880	8354496338831724	$(2)^2(3)(11)^{11}$
8	2	8353355092148436	5569283814460081
9	1	2785212536771726	11139708850800752
10	449132892	8354496314029026	$(3)(7)(11)^6$
11	1	8354496306943626	14321995043777552
12	6	5967498736833926	2386997607312481
13	253080	8354486907040960	$(2)^2(7)(11)^9$
14	77948684	8354496338830880	$(11)^8$
15	71452960	8354496338831724	$(2)^2(3)(11)^7$
16	2	8354496260882196	5569724720561761
17	1	2784953180241326	11139086395127792
18	30675	8354133214886466	$(3)(7)(11)^{10}$
19	1	8354107277462709	11934716868873767
20	2	3580609591411058	4774042371968272
21	1	5967475152525486	7160985746644911
22	1	1193510594119425	9548006930374333
23	3705346371	8354496336254908	$(2)^2(7)(11)^5$
24	5324	8354496338830880	$(11)^{12}$
25	1046142792240	8354496338831724	$(2)^2(3)(11)^3$
26	2	8354496338825556	6455747170913041
27	1	4556998003000526	7594996671666032
28	1	3037998668665506	$(3)(7)(11)^{14}$
29	2	4936747836582555	5696247503748983
30	3	6455747170915411	4936747836582349
31	54249976226179	8354496338831636	$(2)^2(7)(11)$
32	1	8354496338831496	12531744508247409
33	2	4177248169415913	4177248169415891
34	1	4177248169415869	12531744508247497
35	126583277861086	8354496338831628	$(2)^2(3)(11)$
36	44	8354496338831724	$(11)^{14}$
37	448346910960	8354496338830880	$(2)^2(7)(11)^3$
38	6	8354496338826400	2388343994684041
39	1	5975567629277846	14273572257328592
40	252	8298004628050746	$(3)(7)(11)^{12}$
41	1	8310558341556786	11110036453806512
42	2	2799478112249726	5576987225218201
43	8645808200	8354496338186676	$(2)^2(3)(11)^5$
44	644204	8354496338831724	$(11)^{10}$
45	30622697	8354496338830880	$(2)^2(7)(11)^7$
46	3	8354496182934356	4773997955229829
47	2	5967497682755131	7160996046178463
48	3711841	8354494409601795	$(3)(7)(11)^8$
49	1	8354493337806546	11139329595021872

50	2	2784836257215326	5569661582127721
51	590520	8354486907040116	$(2)^2(3)(11)^9$
52	9431790764	8354496338831724	$(11)^6$
53	2091	8354496338830880	$(2)^2(7)(11)^{11}$
54	1	8349931352101948	9545387251968373
55	1	1195455899866425	7162464179012631
56	2	5967008279146206	4773834872688592
57	1	3580661466230978	11935157804843087
58	54345080115	8354496338612109	$(3)(7)(11)^4$
59	1	8354496338625906	11185358734496432
60	2	2830862395870526	5523633943062841
61	40	8216405490255156	$(2)^2(3)(11)^{13}$
62	138090848575730	8354496338831724	$(11)^2$
63	1	8354496338831606	16708992677663312
64	795666317983972	8354496338831706	21
65	1	8354496338831706	16708992677663312
66	138090848575730	8354496338831606	$(11)^2$
67	40	8354496338831724	$(2)^2(3)(11)^{13}$
68	2	8216405490255156	5523633943062841
69	1	2830862395870526	11185358734496432
70	54345080115	8354496338625906	$(3)(7)(11)^4$
$\vdots$	$\vdots$	$\vdots$	$\vdots$

Square factors of D_n

$$D_n = (2 \cdot 509 \cdot 1319011^n + (3^2 11 \cdot 1319011 - 5^2 7)/2)^2 + 4 \cdot 509 \cdot 5^2 7 \cdot 1319011^n$$

For $n = 8$,

h	a_h	Q_h
0	4663426341517151789800515754636230026215015084843995	1
1	2	.
2	408144037624565082345625145279739816272309	$(3^2 11)(5^2 7)x$
3	1	.
4	2	.
5	3	.
6	1	.
7	1	.
8	2	.
9	7	.
10	1	.
11	1064	$(3^2 11)(5^2)509x^7$
12	1	.
13	74	.
14	23826253834950209922899154128767664449	$3^2 5^2 x^2$
15	3	.
16	4	.
17	1	.
18	2	.
19	2209257165	$3^2(5^2 7)509x^6$
20	1	.
21	1	.
22	19	.

23	1	.
24	2	.
25	4	.
26	234593849472344326647051426958	$(3^2 11)(5^2 7)x^3$
27	2	.
28	49	.
29	46359639830137340	$(3^2 11)509x^5$
30	3081352954682231959521312538	x^4
31	34592792081753937238521	$(5^2 7)509x^4$
32	1	.
33	1	.
34	1	.
35	1	.
36	2	.
37	1	.
38	8	.
39	1	.
40	134840323848799462	$(3^2 11)(5^2 7)x^5$
41	2	.
42	1	.
43	3	.
44	3	.
45	1	.
46	4	.
47	1	.
48	1	.
49	3226241544806307046226640448	$(3^2 11)(5^2)509x^3$
50	1	.
51	3	.
52	1	.
53	2	.
54	4	.
55	1	.
56	7871583281986	$3^2 5^2 x^6$
57	1	.
58	1	.
59	1	.
60	4	.
61	2	.
62	1	.
63	6687132706974518642407845671840488	$3^2(5^2 7)509x^2$
64	1	.
65	10	.
66	2	.
67	2	.
68	2	.
69	1	.
70	1	.
71	1	.
72	77503	$(3^2 11)(5^2 7)x^7$
73	6	.
74	5	.
75	2	.
76	1	.
77	140324570892532199234743419300499936832326	$(3^2 11)509x$
78	1018	x^8
79	104707860601002566147639983264355431405332923600	$(5^2 7)509$

80	4	.
81	1	.
82	39	$5^2 x^8$
83	1	.
84	11	.
85	1	.
86	1	.
87	5612982835701287969389736772019997473292	$(3^2 11) 5^2 \cdot 509x$
88	1	.
89	1	.
90	4	.
91	1	.
92	3	.
93	3	.
94	1	.
95	2	.
96	77503	$(3^2 11)(5^2 7)x^7$
97	2	.
98	3	.
99	10	.
100	2	.
101	2	.
102	1	.
103	6687132706974518642407845671840488	$3^2(5^2 7)509x^2$
104	1	.
105	8	.
106	196789582049686	$3^2 x^6$
107	2	.
108	1	.
109	80656038620157676155666011232	$(3^2 11)509x^3$
110	1	.
111	2	.
112	5	.
113	6	.
114	134840323848799463	$(3^2 11)(5^2 7)x^5$
115	10	.
116	3	.
117	2	.
118	2	.
119	34592792081753937238521	$(5^2 7)509x^4$
120	4	.
121	1	.
122	123254118187289278380852500	$5^2 x^4$
123	1	.
124	2	.
125	1	.
126	1	.
127	2	.
128	1	.
129	1854385593205492	$(3^2 11)5^2 \cdot 509x^5$
130	1	.
131	7	.
132	2	.
133	1	.
134	1	.
135	3	.
136	2	.

137	1	.
138	234593849472344326647051426958	$(3^2 11)(5^2 7)x^3$
139	32	.
140	1	.
141	4	.
142	3	.
143	2209257165	$3^2(5^2 7)509x^6$
144	1	.
145	1	.
146	3	.
147	1	.
148	595656345873755248072478853219191611232	$3^2 x^2$
149	2	.
150	1	.
151	26646	$(3^2 11)509x^7$
152	49	.
153	2	.
154	408144037624565082345625145279739816272309	$(3^2 11)(5^2 7)x$
155	1	.
156	2	.
157	3	.
158	373074107321372143184041260370898402097201206787519	5^2
159	3	.
160	2	.
:	:	:

Considering $D/(3 \cdot 5)^2$, we get

h	a_h	Q_h
0	310895089434476785986701050309082001747667672322933	1
1	2	.
2	6122160564368476235184377179196097244084652	$(7)(11)x$
3	2	.
4	2	.
5	1	.
6	4	.
7	15987	$(11)509x$
8	1	.
9	4	.
10	357393807524253148843487311931514966739	x^2
11	1	.
12	2	.
13	33138857487	$(7)509x^6$
14	1	.
15	2	.
16	5	.
17	2	.
18	3518907742085164899705771404380	$(7)(11)x^3$
19	2	.
20	2	.
21	1	.
22	4	.

23	27815783898082403	(11)509x ⁵
24	1	.
25	4	.
26	205423530312148797301420835	x ⁴
27	1	.
28	2	.
29	57654653469589895397535	(7)509x ⁴
30	1	.
31	2	.
32	5	.
33	2	.
34	2022604857731991948	(7)(11)x ⁵
35	2	.
36	2	.
37	1	.
38	4	.
39	48393623172094605693399606739	(11)509x ³
40	1	.
41	4	.
42	118073749229811	x ⁶
43	1	.
44	2	.
45	100306990604617779636117685077607343	(7)509x ²
46	1	.
47	2	.
48	5	.
49	2	.
50	1162556	(7)(11)x ⁷
51	2	.
52	2	.
53	1	.
54	4	.
55	84194742535519319540846051580299962099395	(11)509x
56	1	.
57	4	.
58	67	x ⁸
59	1	.
60	2	.
61	174513101001670943579399972107259052342221539333	(7)509
62	2	.
63	1	.
64	67	x ⁸
65	4	.
⋮	⋮	⋮

FUNCTION FIELD KREEPERS

$$Y^2 = X^{18} + 2X^{12} + 2X^{11} + 2X^9 + X^6 + 2X^5 + 5X^4 + 6X^3 + 6X^2 + 4X + 1$$

Here we have taken

$$m = -1, \quad l = X^3 + X^2 + X + 1, \quad q = r = k = 1, \quad n = 9.$$

h	$a_h(X)$
0	$X^9 + X^3 + X^2 + 1$
1	$\frac{1}{2}X^5 - \frac{1}{2}X^4 + \frac{1}{2}X - \frac{1}{2}$
2	$2X$
3	$\frac{-1}{2}X^7 - \frac{1}{2}X - \frac{1}{2}$
4	$-2X + 2$
5	$\frac{-1}{2}X - \frac{1}{2}$
6	$-2X^3 + 2X^2$
7	$\frac{-1}{2}X^3$
8	$2X^5$
9	$\frac{1}{2}X - \frac{1}{2}$
10	$2X^3 + 2X^2 + 2X + 2$
11	$\frac{1}{2}X - \frac{1}{2}$
12	$2X^5$
13	$\frac{-1}{2}X^3$
14	$-2X^3 + 2X^2$
15	$\frac{-1}{2}X - \frac{1}{2}$
16	$-2X + 2$
17	$\frac{-1}{2}X^7 - \frac{1}{2}X - \frac{1}{2}$
18	$2X$
19	$\frac{1}{2}X^5 - \frac{1}{2}X^4 + \frac{1}{2}X - \frac{1}{2}$
20	$2X^9 + 2X^3 + 2X^2 + 2$

h	$P_h(X)$	$Q_h(X)$
0	0	1
1	$X^9 + X^3 + X^2 + 1$	$4X^4 + 4X^3 + 4X^2 + 4X$
2	$X^9 - X^3 - X^2 - 2X - 1$	X^8
3	$X^9 + X^3 + X^2 + 2X + 1$	$-4X^2$
4	$X^9 + X^3 + X^2 - 2X - 1$	$-X^8 - X^7 - 2X^2 - 3X - 1$
5	$X^9 - 2X^7 + 3X^3 + X^2 - 2X - 1$	$-4X^8 + 4X^7 + 4X^4 - 4X^3 - 4X^2$
6	$X^9 - 2X^5 + X^3 + X^2 + 2X + 1$	$-X^6 - X^5 - X^4 - X^3$
7	$X^9 - X^3 - X^2 - 2X - 1$	$-4X^6$
8	$X^9 + X^3 + X^2 + 2X + 1$	X^4
9	$X^9 - X^3 - X^2 - 2X - 1$	$4X^8 + 4X^7 + 4X^6 + 4X^5$
10	$X^9 - 2X^5 + X^3 + X^2 + 2X + 1$	$X^6 - X^5 - X^2 + X + 1$
11	$X^9 - 2X^5 + X^3 + X^2 + 2X + 1$	$4X^8 + 4X^7 + 4X^6 + 4X^5$
12	$X^9 - X^3 - X^2 - 2X - 1$	X^4
13	$X^9 + X^3 + X^2 + 2X + 1$	$-4X^6$
14	$X^9 - X^3 - X^2 - 2X - 1$	$-X^6 - X^5 - X^4 - X^3$
15	$X^9 - 2X^5 + X^3 + X^2 + 2X + 1$	$-4X^8 + 4X^7 + 4X^4 - 4X^3 - 4X^2$
16	$X^9 - 2X^7 + 3X^3 + X^2 - 2X - 1$	$-X^8 - X^7 - 2X^2 - 3X - 1$
17	$X^9 + X^3 + X^2 - 2X - 1$	$-4X^2$
18	$X^9 + X^3 + X^2 + 2X + 1$	X^8
19	$X^9 - X^3 - X^2 - 2X - 1$	$4X^4 + 4X^3 + 4X^2 + 4X$
20	$X^9 + X^3 + X^2 + 1$	1

HIGHER CREEPERS

$$D_n = (3^{2n+2} + 3^{n+2} + 3^n - 1)^2 + 4 \cdot 3^n$$

For $n = 14$,

h	a_h	P_h	Q_h	factors of Q_h
0	102945589962169	0	1	
1	43046730	102945589962169	4782969	3^{14}
2	1	102945585179201	205891132094641	
3	4782968	102945546915440	43046730	$3^2 J$
4	9	102945585179200	22876792454961	3^{28}
5	531441	102945546915449	387420489	3^{18}
6	80	102945585179200	2541866359770	$3^{12} J$
7	1	100403723602400	203349313564489	
8	2541866418818	102945589962089	81	3^4
9	3486785210	102945589962169	59049	3^{10}
10	1	102945589903121	205887693080161	
11	59048	102942103177040	3486785130	$3^6 J$
12	729	102945585179200	282429536481	3^{24}
13	6561	102945546915449	31381059609	3^{22}
14	6560	102945585179200	31381066170	$3^8 J$
15	1	102914208896000	205859798851609	
16	31381066898	102945589955609	6561	3^8
17	282429602090	102945589962169	729	3^6
18	1	102945589961441	205608750328081	
19	728	102663160366640	282429595530	$3^{10} J$
20	59049	102945585179200	3486784401	3^{20}
21	81	102945546915449	2541865828329	3^{26}
22	531440	102945585179200	387420570	$3^4 J$
23	1	102945202541600	205890791972329	
24	387420578	102945589430729	531441	3^{12}
25	22876797769370	102945589962169	9	3^2
26	1	102945589962161	183014382686401	
27	8	80068792724240	22876797237930	$3^{14} J$
28	4782969	102945585179200	43046721	3^{16}
29	1	102945546915449	205891132094649	3^{30}
30	43046720	102945585179200	4782970	
31	1	102945585179200	205891132094649	3^{30}

where $J = 2 \cdot 5 \cdot 29 \cdot 16493 = 3^{14} + 1$.

$$D_n = (3^{2n+3} - 31 \cdot 3^n + 10)^2 + 40 \cdot 3^n$$

For $n = 21$,

h	a_h	Q_h	factors of Q_h
0	1477156353113281374680	1	
1	28242953645	104603532030	$2 \cdot 3^{21} \cdot 5$
2	10460353202	282429536451	$3 \cdot J_{21}$
3	1	2844893716833541560106	
4	25	109418989131512359209	3^{42}
5	1	2844893709249785488181	
6	348678439	8472886094130	$2 \cdot 3 \cdot 5 \cdot K_{21}$
7	8	328256967255065701591	
8	1	2626055738898274575346	
9	31381059606	94143178821	$3 \cdot I_{21}$
10	1	1969541804046438300886	

11	1	984770902148743388871	$3^{20} \cdot K_{21}$
12	1	1969541803889533002851	
13	5230176600	564859072926	$2 \cdot 3^2 \cdot I_{21}$
14	1	1750703825731111816451	
15	1	1203608880390847400879	
16	2	1094189891245387904070	$2 \cdot 3^{20} \cdot 5 \cdot I_{21}$
17	5	547094945162438411119	
18	232452293	12709329141195	$3^2 \cdot 5 \cdot K_{21}$
19	5	583567941474522555374	
20	15	182364981874231317345	$3^{19} \cdot 5 \cdot I_{21}$
21	1	2771947722866961277349	
22	1743392199	1694577218706	$2 \cdot 3^2 \cdot J_{21}$
23	1	1641284836286951122655	
24	1	1313027869751325269051	
25	3	656513934765828925914	$2 \cdot 3^{19} \cdot K_{21}$
26	1	2297798770676207333201	
27	2092070639	1412147682405	$3^{24} \cdot 5$
28	1	1750703825416138958886	
29	1	1203608880548914960399	
30	2	1094189891146595679379	
31	5	547094945599448722695	$3^{19} \cdot 5 \cdot J_{21}$
32	54709494559751162025	54	$2 \cdot 3^3$
33	1525119496830	1937102445	$3^{18} \cdot 5$
34	193710244	15251194968354	$2 \cdot 3^4 \cdot J_{21}$
35	2	1475130071734433497661	
36	363	8105110306037952534	$2 \cdot 3^{39}$
37	1	2946207482164183718231	
38	25828031	114383962270755	$3^4 \cdot 5 \cdot K_{21}$
39	1	1775019115536292522986	
40	1	1179293567625191414569	
41	1	1191451210206164487589	
42	1	1750703830374346376946	
43	47	60788327288827635855	$3^{17} \cdot 5 \cdot J_{21}$
44	1	2893524374063210521211	
45	581130732	5083731656334	$2 \cdot 3^4 \cdot I_{21}$
46	2	1458919853685789827819	
47	39	72945992751758769546	$2 \cdot 3^{17} \cdot K_{21}$
48	1	2881366706225779210601	
49	387420488	7625597484501	$3^5 \cdot I_{21}$
50	145	20262775763803479705	$3^{17} \cdot 5 \cdot I_{21}$
51	1	2350481961341523199726	
52	3	591673079803330177099	
53	1	2362639215017541112831	
54	4304671	686303773624530	$2 \cdot 3^5 \cdot 5 \cdot K_{21}$
55	3	886158664947699144501	
56	2	887509585295663405601	
57	1	2061399714123313868389	
58	217	13508517175868986470	$2 \cdot 3^{16} \cdot 5 \cdot I_{21}$
59	1	2940804166383108374399	
60	129140162	22876792452531	$3^5 \cdot J_{21}$
61	242	12157665458626461591	$3^{16} \cdot K_{21}$
62	1	2942154965139582630059	
63	38742047	76255974849870	$2 \cdot 3^{27} \cdot 5$

64	1	2658476118340136630889	
65	8	295836526902566944981	
66	1	2626055738124208438329	
67	72	40525551525885090570	$2 \cdot 3^{16} \cdot 5 \cdot J_{21}$
68	4052555152574160150	729	3^6
69	20589113207205	143489070	$2 \cdot 3^{15} \cdot 5$
70	14348906	205891132072779	$3^7 \cdot J_{21}$
71	1	2954162405721054383866	
72	19681	150094635296999121	3^{36}
73	1	2954156435484924321629	
74	478295	6176733962620770	$2 \cdot 3^7 \cdot 5 \cdot K_{21}$
75	1	2658921460670523346479	
76	9	295386303980582948611	
77	655	4502839058431676730	$2 \cdot 3^{14} \cdot 5 \cdot J_{21}$
78	1	2949809798746960776161	
79	43046719	68630377360509	$3^7 \cdot I_{21}$
80	1	2952961785941322345670	
81	2185	1350851717625162399	$3^{14} \cdot K_{21}$
82	1	2952961443103246139195	
83	7174452	411782264163054	$2 \cdot 3^8 \cdot I_{21}$
84	1	1477531486703632150859	
85	1	1476781219418327066471	
86	1967	1500946352874331830	$2 \cdot 3^{14} \cdot 5 \cdot I_{21}$
87	1	2362489189012631042101	
88	3	590922949778779823449	
89	1	2363384197763789282506	
90	318863	9265100943931155	$3^8 \cdot 5 \cdot K_{21}$
91	1	2363434229057836616711	
92	3	590872918170921893174	
93	1	2363289693483115542221	
94	11808	250157725479055305	$3^{13} \cdot 5 \cdot I_{21}$
95	1	2954061313363498321421	
96	2391483	1235346792436674	$2 \cdot 3^8 \cdot J_{21}$
97	1	1477381186135377613751	
98	1	1476931519902898777955	
99	3279	900567811750108266	$2 \cdot 3^{13} \cdot K_{21}$
100	1	2953411109586773360009	
101	2869780	1029455660473245	$3^{30} \cdot 5$
102	2	1181694898859031495409	
103	1	1181845199385444645949	
104	1	1772317412226757753926	
105	3936	750473176405279455	$3^{13} \cdot 5 \cdot J_{21}$
106	75047317640262225	39366	$2 \cdot 3^9$
107	1111812113189070	2657205	$3^{12} \cdot 5$
108	265720	11118121131930066	$2 \cdot 3^{10} \cdot J_{21}$
109	2	1477150794063175467605	
110	265719	11118121133111046	$2 \cdot 3^{33}$
111	1	2954218202824555450079	
112	35428	83385908495380395	$3^{10} \cdot 5 \cdot K_{21}$
113	2	1181715076106290874839	
114	2	1181715076106292055819	
115	35428	83385908489475495	$3^{11} \cdot 5 \cdot J_{21}$
116	1	2954225614486902870419	

117	797160	3706040377467486	$2 \cdot 3^{10} \cdot I_{21}$
118	2	1477130410862019453299	
119	29523	100063090194456474	$2 \cdot 3^{11} \cdot K_{21}$
120	1	2954207084452374806945	
121	531440	5559060566201229	$3^{11} \cdot I_{21}$
122	106288	27795302831006145	$3^{11} \cdot 5 \cdot I_{21}$
123	5	590841416890474961389	
124	5904	500315450972282370	$2 \cdot 3^{11} \cdot 5 \cdot K_{21}$
125	1	2067773554529532116269	
126	2	886339025579403839361	
127	3	886287141014117877381	
128	159431	18530201887337430	$2 \cdot 3^{10} \cdot 5 \cdot I_{21}$
129	1	2954277499052184580871	
130	177146	16677181697895099	$3^{11} \cdot J_{21}$
131	177146	16677181699076079	$3^{10} \cdot K_{21}$
132	1	2954240439066819310211	
133	53143	55590605665555230	$2 \cdot 3^{33} \cdot 5$
134	10	295430158812635093521	
135	53144	55590605659650330	$2 \cdot 3^{10} \cdot 5 \cdot J_{21}$
136	5559060565945350	531441	3^{12}
137	15009463528052445	196830	$2 \cdot 3^9 \cdot 5$
138	19682	150094635281055891	$3^{13} \cdot J_{21}$
139	1	2954162405721070305226	
140	14348905	205891132094649	3^{30}
141	1	2949809661904301305541	
142	655	4502839058750541330	$2 \cdot 3^{13} \cdot 5 \cdot K_{21}$
143	9	295386303980579755591	
144	1	2658921460670526889419	
145	478295	6176733962183370	$2 \cdot 3^8 \cdot 5 \cdot J_{21}$
146	1	2954256498156711818969	
147	59047	50031545095811061	$3^{13} \cdot I_{21}$
148	1	2954260821724040271286	
149	1594321	1853020188786231	$3^8 \cdot K_{21}$
150	1	2954010664312371812051	
151	9840	300189270574866366	$2 \cdot 3^{14} \cdot I_{21}$
152	1	1477081820471166095891	
153	1	1477230885650793121439	
154	1434890	2058911320815270	$2 \cdot 3^8 \cdot 5 \cdot I_{21}$
155	5	590592288620649227839	
156	437	6754258588125811995	$3^{14} \cdot 5 \cdot K_{21}$
157	5	590592357188264469134	
158	8609343	343151886802545	$3^7 \cdot 5 \cdot I_{21}$
159	1	2953411795472196675509	
160	3279	900567811686335346	$2 \cdot 3^{14} \cdot J_{21}$
161	1	1476931519902914743055	
162	1	1477381186135361648651	
163	2391483	1235346792524154	$2 \cdot 3^7 \cdot K_{21}$
164	1	2953560998330906421761	
165	3935	750473176484995605	$3^{36} \cdot 5$
166	1	1772317412226729060486	
167	1	1181845199385457396159	
168	2	1181694898859028324259	
169	2869781	1029455660363895	$3^7 \cdot 5 \cdot J_{21}$

170	102945566036025	28697814	$2 \cdot 3^{15}$
171	810511030514832030	3645	$3^6 \cdot 5$
172	364	8105110305177018114	$2 \cdot 3^{16} \cdot J_{21}$
173	2	1475130071734648730861	
174	193710243	15251194969974	$2 \cdot 3^{27}$
175	1	2893524364309856663591	
176	47	60788327293132307955	$3^{16} \cdot 5 \cdot K_{21}$
177	1	1750703830372796694666	
178	1	1191451210206853235449	
179	1	1179293567625019226389	
180	1	1775019115536464712786	
181	25828031	114383962262655	$3^5 \cdot 5 \cdot J_{21}$
182	1	2951610888616633753451	
183	1092	2701703435173797294	$2 \cdot 3^{16} \cdot I_{21}$
184	2	1476480892970680303739	
185	21523359	137260754724906	$2 \cdot 3^5 \cdot K_{21}$
186	1	2950260014189620043801	
187	728	4052555152760695941	$3^{17} \cdot I_{21}$
188	77484097	38127987422505	$3^5 \cdot 5 \cdot I_{21}$
189	1	2348860942330244667406	
190	3	605451741396098343739	
191	1	2130022986951760812271	
192	7	364729963758793847730	$2 \cdot 3^{17} \cdot 5 \cdot K_{21}$
193	3	853468114894319431461	
194	2	889941113812064644401	
195	1	2064371582309796911509	
196	116226145	25418658281670	$2 \cdot 3^4 \cdot 5 \cdot I_{21}$
197	1	2942155015559346004559	
198	242	12157665457765527171	$3^{17} \cdot J_{21}$
199	129140162	22876792454151	$3^4 \cdot K_{21}$
200	1	2913787132447201724699	
201	71	40525551530189762670	$2 \cdot 3^{39} \cdot 5$
202	1	2626055738120721654009	
203	8	295836526902609991621	
204	1	2658476118340093590729	
205	38742048	76255974841770	$2 \cdot 3^4 \cdot 5 \cdot J_{21}$
206	7625597484150	387420489	3^{18}
207	10941898911950232405	270	$2 \cdot 3^3 \cdot 5$
208	26	109418989119889744539	$3^{19} \cdot J_{21}$
209	1	2844893716845164174746	
210	10460353200	282429536481	3^{24}
211	1	2626055738749505107546	
212	8	328256967257390224531	
213	348678439	8472886093530	$2 \cdot 3^2 \cdot 5 \cdot J_{21}$
214	1	2917839701588037456401	
216	1	2917839707372612776870	
217	1162261465	2541865828239	$3^2 \cdot K_{21}$
218	1	2735474725812192055595	
219	12	218837978249077580814	$2 \cdot 3^{20} \cdot I_{21}$
220	1	1422446859204784054619	
221	1	1531865846917175162711	
222	1046035319	2824295364630	$2 \cdot 3^2 \cdot 5 \cdot I_{21}$
223	1	2166495983045210660581	

224	2	787816721863347585289	
225	1	1772587623604137199269	
226	2	787816721706442287254	
227	1	2166495984300453044861	
228	6276211920	470715894105	
229	1	2297798771245715452061	
230	3	656513934719338467234	$2 \cdot 3^{20} \cdot J_{21}$
231	1	1313027869762947883751	
232	1	1641284836275328507955	
233	1743392199	1694577218826	$2 \cdot 3 \cdot K_{21}$
234	1	2407217759502044926649	
235	4	547094945657561796045	$3^{42} \cdot 5$
236	2	1094189891137297587649	
237	1	1203608880551239483309	
238	1	1750703825413814436006	
239	2092070640	1412147682255	
240	141214768225	20920706406	$2 \cdot 3^{21}$
241	590862541245312549872	5	
242	141214768225	20920706406	$2 \cdot 3^{21}$
:	:	:	:
:	:	:	:

JEEPERS

The starred lines indicate the lines where Q_h is a product of a power of 2 and a power of 3.

h	a_h	P_h	Q_h	
0	131071	0	1	
1	2	131070	131070	
2	87380	131069	3	*
3	6	131070	43690	
4	29126	131069	3^2	*
5	1	131064	189320	
6	2	58255	72818	
7	1	87380	131070	
8	1	43689	116509	
9	2	72819	101940	
10	9708	131060	3^3	*
11	1	131055	150480	
12	1	19424	111658	
13	2	92233	77671	
14	1	63108	169908	
15	7	106799	33980	
16	3236	131060	3^4	*
17	5	131055	50160	
18	4	119744	56636	
19	2	106799	101940	
20	2	97080	76074	
21	1	55067	185966	
22	1078	130898	3^5	*
23	15	131055	16720	

24	1	119744	169908	
25	2	50163	86301	
26	9	122438	25358	
27	1	105783	$2^2 3^{10}$	*
28	358	130412	3^6	*
29	1	130569	179990	
30	2	49420	81878	
31	4	114335	50160	
32	1	86304	194002	
33	8	107697	28767	
34	3	122438	76074	
35	3	105783	$2^2 3^9$	*
36	119	130412	3^7	*
37	1	129840	146800	
38	1	16959	115068	
39	3	98108	65651	
40	2	98844	112860	
41	26	126875	9589	
42	1	122438	228222	
43	9	105783	$2^2 3^8$	*
44	39	130412	3^8	*
45	1	125466	219138	
46	5	93671	38356	
47	1	98108	196953	
48	6	98844	37620	
49	8	126875	28767	
50	1	103260	226540	
51	29	123279	$2^2 3^7$	*
52	13	130412	3^9	*
53	3	125466	73046	
54	1	93671	115068	
55	1	21396	145321	
56	20	123924	12540	
57	2	126875	86301	
58	1	45726	174838	
59	89	129111	$2^2 3^6$	*
60	4	130412	3^{10}	*
61	2	105783	101432	
62	2	97080	76455	
63	1	55829	183934	
64	62	128104	4180	
65	269	131055	$2^2 3^5$	*
66	1	130412	3^{11}	*
67	2	46734	84650	
68	9	122565	25485	
69	1	106799	226544	
70	20	119744	12540	
71	809	131055	$2^2 3^4$	*
72	30	131060	8495	
73	1	123789	218454	
74	6	94664	37620	
75	2427	131055	$2^2 3^3$	*
76	10	131060	25485	

77	3	123789	72818	
78	2	94664	112860	
79	7281	131055	$2^2 3^2$	*
80	3	131060	76455	
81	2	98304	$2^{15} 3$	*
82	3	98303	76457	
83	21844	131067	$2^2 3$	*
84	1	131060	229365	
85	6	98304	2^{15}	*
86	1	98303	229371	
87	65534	131067	2^2	*
88	1	131068	163837	
89	1	32768	$2^{15} 3$	*
90	1	65535	131070	
91	1	65534	98305	
92	1	32770	163834	
93	21844	131063	$2^2 3$	*
94	1	131064	141990	
95	1	10925	120151	
96	5	109225	43690	
97	1	109224	120156	
98	1	10931	141983	
99	7281	131051	$2^2 3^2$	*
100	5	131064	47330	
101	1	105585	127431	
102	1	21845	131070	
103	5	109224	40052	
104	1	91035	222015	
105	2426	130979	$2^2 3^3$	*
106	2	131028	103141	
107	1	75253	111658	
108	1	36404	141990	
109	5	105585	42477	
110	1	106799	135920	
111	1	29120	120156	
112	3	91035	74005	
113	808	130979	$2^2 3^4$	*
114	1	130812	208941	
115	3	78128	53008	
116	1	80895	200640	
117	17	119744	14159	
118	1	120958	180002	
119	2	59043	76074	
120	2	93104	111880	
121	269	130655	$2^2 3^5$	*
122	3	130812	69647	
123	1	78128	159024	
124	3	80895	66880	
125	5	119744	42477	
126	1	92640	202400	
127	9	109759	25358	
128	2	118462	124073	
129	89	129683	$2^2 3^6$	*

130	2	129840	110100	
131	2	90359	81878	
132	1	73396	144026	
133	2	70629	84645	
134	2	98660	87964	
135	1	77267	127431	
136	1	50163	115068	
137	1	64904	112690	
138	1	47785	132187	
139	2	84401	76074	
140	1	67746	165497	
141	4	97750	46070	
142	1	86529	210381	
143	29	123851	$2^2 3^7$	*
144	7	129840	36700	
145	9	127059	28215	
146	6	126875	38356	
147	1	103260	169905	
148	2	66644	74972	
149	1	83299	136595	
150	1	53295	$2^4 3^8$	*
151	1	51680	138210	
152	3	86529	70127	
153	9	123851	$2^2 3^8$	*
154	1	112344	173690	
155	2	61345	77243	
156	2	93140	110100	
157	27	127059	9405	
158	2	126875	115068	
159	4	103260	56635	
160	7	123279	$2^4 3^7$	*
161	3	121664	67940	
162	1	82155	153519	
163	2	71363	$2^2 3^9$	*
164	1	86100	124045	
165	1	37944	126888	
166	3	88943	73046	
167	83	130194	3135	
168	2	130010	88318	
169	1	46625	169905	
170	21	123279	$2^4 3^6$	*
171	1	121664	203820	
172	4	82155	51173	
173	5	122536	42296	
174	1	88943	219138	
175	250	130194	1045	
176	67	131055	$2^4 3^5$	*
177	2	129440	109250	
178	2	89059	84650	
179	1	80240	126888	
180	1	46647	118243	
181	1	71595	101940	
182	1	30344	159494	

183	83	129149	3135	
184	202	131055	$2^4 3^4$	*
185	3	130736	67573	
186	1	71982	177558	
187	6	105575	33980	
188	1	98304	$2^{13} 3^3$	*
189	27	122879	9405	
190	606	131055	$2^4 3^3$	*
191	1	130736	202719	
192	3	71982	59186	
193	2	105575	101940	
194	3	98304	$2^{13} 3^2$	*
195	9	122879	28215	
196	1820	131055	$2^4 3^2$	*
197	3	131024	84635	
198	10	122880	$2^{13} 3$	*
199	3	122879	84645	
200	5460	131055	$2^4 3$	*
201	1	131024	253905	
202	30	122880	2^{13}	*
203	1	122879	253935	
204	16382	131055	2^4	*
205	1	131056	237553	
206	9	106496	$2^{13} 3$	*
207	1	114687	163834	
208	1	49146	90117	
209	1	40970	172010	
210	5460	131039	$2^4 3$	*
211	1	131040	166550	
212	1	35509	95579	
213	1	60069	141990	
214	2	81920	$2^{13} 3^2$	*
215	1	65535	174760	
216	7	109224	30039	
217	1	101048	231992	
218	1819	130943	$2^4 3^2$	*
219	1	130992	142861	
220	1	11868	119268	
221	5	107399	47330	
222	25	129250	10013	
223	1	121074	251730	
224	605	130655	$2^4 3^3$	*
225	1	130704	222085	
226	5	91380	39756	
227	1	107399	141990	
228	1	34590	112565	
229	2	77974	98606	
230	8	119237	30039	
231	3	121074	83910	
232	201	130655	$2^4 3^4$	*
233	1	129840	247725	
234	18	117884	13252	
235	1	120651	197919	

236	3	77267	56636	
237	1	92640	151800	
238	2	59159	90117	
239	9	121074	27970	
240	67	130655	$2^4 3^5$	*
241	3	129840	82575	
242	6	117884	39756	
243	3	120651	65973	
244	1	77267	169908	
245	4	92640	50600	
246	2	109759	101432	
247	2	93104	83910	
248	1	74715	138210	
249	2	63494	95131	
250	22	126767	$2^4 3^6$	*
251	9	129840	27525	
252	2	117884	119268	
253	11	120651	21991	
254	2	121249	112690	
255	4	104130	56229	
256	5	120785	46070	
257	2	109564	112334	
258	7	115103	$2^4 3^7$	*
259	28	129840	9175	
260	2	127059	112860	
261	3	98660	65973	
262	2	99258	111066	
263	13	122873	18743	
264	1	120785	138210	
265	1	17424	122104	
266	4	104679	50955	
267	1	99140	144260	
268	1	45119	$2^4 3^8$	*
269	1	59856	129523	
270	2	69666	95166	
271	9	120665	27525	
272	6	127059	37620	
273	1	98660	197919	
274	6	99258	37022	
275	4	122873	56229	
276	1	102042	120346	
277	1	18303	$2^6 3^7$	*
278	14	121664	16985	
279	1	116125	217514	
280	7	101388	31722	
281	3	120665	82575	
282	20	127059	12540	
283	1	123740	148955	
284	1	25214	111066	
285	2	85851	88318	
286	2	90784	101200	
287	5	111615	$2^6 3^6$	*
288	4	121664	50955	

289	1	82155	204692	
290	23	122536	10574	
291	1	120665	247725	
292	61	127059	4180	
293	1	127920	195204	
294	3	67283	64817	
295	16	127167	$2^6 3^5$	*
296	1	121664	152865	
297	1	31200	106016	
298	1	74815	109250	
299	1	34434	146397	
300	7	111962	31722	
301	1	110091	159494	
302	1	49402	92411	
303	1	43008	$2^{11} 3^4$	*
304	20	122879	12540	
305	3	127920	65068	
306	1	67283	194451	
307	49	127167	$2^6 3^4$	*
308	1	126848	210082	
309	4	83233	48799	
310	2	111962	95166	
311	1	78369	115985	
312	1	37615	135920	
313	4	98304	$2^{11} 3^3$	*
314	6	122879	37620	
315	1	102840	175530	
316	3	72689	67771	
317	151	130623	$2^6 3^3$	*
318	2	130304	115940	
319	3	101575	59186	
320	1	75982	192719	
321	13	116736	$2^{11} 3^2$	*
322	2	122879	112860	
323	3	102840	58510	
324	1	72689	203313	
325	454	130623	$2^6 3^2$	*
326	3	130880	86635	
327	42	129024	$2^{11} 3$	*
328	3	129023	86677	
329	1364	131007	$2^6 3$	*
330	1	130880	259905	
331	126	129024	2^{11}	*
332	1	129023	260031	
333	4094	131007	2^6	*
334	1	131008	255937	
335	41	124928	$2^{11} 3$	*
336	1	126975	172010	
337	1	45034	88085	
338	1	43050	173994	
339	1364	130943	$2^6 3$	*
⋮				

Bibliography

1. W.W. Adams and M.J. Razar, *Multiples of points on elliptic curves and continued fractions*, Proc. London Math. Soc. **41** (1980), no. 3, 481–498.
2. T. Azuhata, *On the fundamental units and the class numbers of real quadratic fields*, Nagoya Math J. **95** (1984), 125–135.
3. ———, *On the fundamental units and the class numbers of real quadratic fields II*, Tokyo J. Math **10** (1987), no. 2, 259–270.
4. L. Bernstein, *Fundamental unit and cycles I*, J. of Numb. The. **8** (1976), no. 4, 446–491.
5. ———, *Fundamental units and cycles in the period of real quadratic number fields I,II*, Pac. J. Math. **63** (1976), 37–61, 63–78.
6. T.G. Berry, *On periodicity of continued fractions in hyperelliptic function fields*, Arch. Math. (Basel) **55** (1990), no. 3, 259–266.
7. Z. Borevich and I. Shafarevich, *Number Theory*, Academic Press, New York, 1966.
8. N. Buck and K. Williams, *Comparision of the lengths of the contnued fractions of $\sqrt{D}$ and $\frac{1+\sqrt{D}}{2}$* , Proc. Amer. Math. Soc. **120** (1994), no. 4, 992–1002.
9. D. Cantor, *Computing in the Jacobian of a Hyperelliptic curve*, Math. Comp. **48** (1987), no. 177, 95–101.
10. J.W.S Cassels and E.V. Flynn, *Prolegomena to a Middlebrow Arithmetic of Curves of Genus 2*, London Mathematical Society Lecture Note Series, vol. 230, Cambridge University Press, Cambridge, 1996.
11. H. Cohen and H.W. Lenstra Jr, *Heuristics on class groups of number fields*, Number Theory (Noordwijkerhout 1983), Lecture Notes in Math., no. 1068, Springer-Verlag, Berlin, 1984, pp. 33–62.
12. H. Cohn, *Advanced number theory*, Dover Publications, Inc., New York, 1980, Reprint of “A second course in number theory”.
13. D.A. Cox, *Primes of the form $x^2 + ny^2$. Fermat, class field theory and complex multiplication*, John Wiley & Sons, Inc., New York, 1989.
14. P.G.L. Dirichlet, *Lectures on number theory*, History of Mathematics, vol. 16, Amer. Math. Soc., Providence RI, 1999, Supplements by R. Dedekind. Translated from the 1863 original and with an introduction by John Stillwell.
15. N.D. Elkies, *Simple genus 2 Jacobians with high order torsion points*, <http://www.math.harvard.edu/~elkies/g2-tors.html>.
16. ———, *Heegner point computations*, Algorithmic number theory (Ithaca, NY, 1994), Lecture Notes in Comput. Sci., vol. 877, Springer, Berlin, 1994, pp. 122–133.
17. D. Flath, *Introduction to number theory*, John Wiley & Sons, Inc., New York, 1989.
18. E.V. Flynn, *Large rational torsion on abelian varieties*, J. of Numb. The. **36** (1990), no. 3, 257–265.
19. ———, *Sequences of rational torsion on abelian varieties*, Invent. Math. **106** (1991), no. 2, 433–442.
20. C.F. Gauss, *Diquistiones Arithmeticae*, Yale Univ. Press, New Haven, Conn., 1966.
21. F. Halter-Koch, *Reell-quadratische Zahlkörper mit gröser Grundeinheit*, Abh. Math. Sem. Univ. Hamburg **59** (1989), 171–181.
22. M.D. Hendy, *Applications of a continued fraction algorithm to some class number problems*, Math. Comp. **28** (1974), 267–277.
23. E. Howe, F. Leprévost, and B. Poonen, *Large torsion sungroups of split Jacobians of curves of genus 2 or 3*, Forum Math. **12** (2000), no. 3, 315–364.
24. D. Husemöller, *Elliptic curves*, Graduate Texts in Mathematics, vol. 111, Springer-Verlag, New York, 1987.
25. N. Ishii, P. Kaplan, and K. Williams, *On Eisenstein’s problem*, Acta. Arth. **54** (1990), no. 4, 323–345.
26. M.J. Jacobson Jr., *Subexponential Class Group Computation in Quadratic Orders*, Ph.D. thesis, Darmstadt, 1999.
27. M.J. Jacobson Jr. and A.J. van der Poorten, *Computational aspects of NUCOMP*, Algorithmic Number Theory (Proc. Fifth International Symposium, ANTS-V, Sydney, NSW, Australia, July 2002) (C. Fieker and D.R. Kohel, eds.), Springer Lecture Notes in Computer Science, no. 2369, 2002, pp. 390–404.
28. I. Kaplansky, *Letter to 3 Canadian number theorists*, 1998.
29. A. Khinchin, *Continued fractions*, third ed., Univ. of Chicago Press, 1964.
30. H. Koch, *Number Theory, Algebraic numbers and functions*, Graduate Series in Mathematics, vol. 24, Amer. Math. Soc., Providence RI, 2000, Translated from the 1997 German original by David Kramer.

31. D. Kubert, *Universal bounds on the torsion of elliptic curves*, Proc. London Math. Soc. **33** (1976), no. 3, 193–237.
32. S. Lang, *Algebraic number theory*, Springer-Verlag, New York, 1986.
33. H.W. Lenstra Jr, *On the calculation of regulators and class numbers of quadratic fields*, Number theory days, 1980 (Exeter 1980) (J.V. Armitage, ed.), London Math. Soc. Lecture Note Ser., no. 56, Cambridge Univ. Press, Cambridge, 1982, pp. 123–150.
34. F. Leprévost, *Familles de courbes de genre 2 munies d’une classe de diviseurs rationnels d’ordre 15, 17, 19, ou 21*, C. R. Acad. Sci. Paris Sér. I Math. **313** (1991), no. 11, 771–774.
35. ———, *Torsion sur des familles de courbes de genre g* , Manuscripta Math. **75** (1992), no. 3, 303–326.
36. ———, *Points rationnels de torsion de Jacobiennes de certaines courbes de genre 2*, C. R. Acad. Sci. Paris Sér. I Math. **316** (1993), no. 8, 819–821.
37. ———, *Jacobiennes de certaines courbes de genre 2: torsion et simplicité*, J. Théor. Nombres Bordeaux **7** (1995), no. 1, 283–306.
38. ———, *Sur certains sous-groupes de torsion de Jacobiennes de courbes hyperelliptiques de genre $g \geq 1$* , Manuscripta Math. **92** (1997), no. 1, 47–63.
39. C. Levesque and G. Rhin, *A few classes of periodic continued fractions*, Util. Math. **30** (1986), 79–107.
40. D. Lorenzini, *An Invitation to Arithmetic Geometry*, Graduate Series in Mathematics, vol. 9, Amer. Math. Soc., Providence RI, 1996.
41. R.E. MacRae, *On unique factorization in certain rings of algebraic functions*, J. Algebra **17** (1971), 243–261.
42. D. Madden, *Constructing families of long continued fractions*, Pac. J. Math. **198** (2001), no. 1, 123–147.
43. B. Mazur, *Modular curves and the Eisenstein ideal*, Inst. Hautes Études Sci. Publ. Math. **47** (1977), 33–186.
44. M. Mendès France, *Sur les fractions continues limitées*, Acta. Arith. **23** (1973), 207–215.
45. A. Menezes, Y-H Wu, and R. Zuccherato, *Algebraic aspects of cryptography*, Algorithms and Computation in Mathematics, no. 3, ch. Appendix, Springer-Verlag, 1998.
46. R.A. Mollin and K. Cheng, *Period lengths of continued fractions involving Fibonacci numbers*, Fibonacci Quarterly, to appear.
47. ———, *Beepers, creepers and sleepers*, Int. Math. J. **2** (2002), no. 9, 951–956.
48. ———, *Continued fractions beepers and Fibonacci numbers*, C. R. Math. Acad. Sci. Soc. R. Can. **24** (2002), no. 3, 102–108.
49. R.A. Mollin, K. Cheng, and B. Goddard, *Pellian polynomials and period lengths of continued fractions*, JP J. Algebra Number Theory Appl. (2002), no. 1, 47–60.
50. R.A. Mollin and A.J. van der Poorten, *A note on symmetry and ambiguity*, Bull. Austral. Math. Soc. **51** (1995), 215–233.
51. R.A. Mollin and H.C. Williams, *Computation of the class number of a real quadratic field*, Util. Math. **41** (1992), 259–308.
52. ———, *Consecutive powers in continued fractions*, Acta. Arith. **51** (1992), no. 3, 233–264.
53. ———, *On the period length of some special continued fractions*, J. Théor. Nombres Bordeaux **4** (1992), no. 1, 19–42.
54. R.A. Mollin, H.C. Williams, and A.J. van der Poorten, *Halfway to a solution of $X^2 - DY^2 = -3$* , J. Théor. Nombres Bordeaux **6** (1994), 421–457.
55. T. Nagell, *Problems in the theory of exceptional points on plane cubics of genus one*, Den 11te Skandinaviske Matematikerkongress (1949), 71–76.
56. M. Nyberg, *Culminating and almost culminating continued fractions*, Norsk. Mat. Tidsskr. **31** (1949), 95–99, In Norwegian.
57. S. Paulus and H-G Rück, *Real and imaginary quadratic representations of hyperelliptic function fields*, Math. Comp. **68** (1999), no. 227, 1233–1241.
58. O. Perron, *Die Lehre von den Kettenbrüchen*, Second ed., Chelsea Publishing Co., New York, 1950.
59. G.N. Raney, *On continued fractions and finite automata*, Math. Ann. **206** (1973), 265–283.
60. A. Schinzel, *On some problems of the arithmetical theory of continued fractions*, Acta. Arith. **6** (1961), 393–413.
61. ———, *On some problems of the arithmetical theory of continued fractions II*, Acta. Arith. **7** (1962), 287–298.
62. D. Shanks, *On Gauss’s class number problems*, Math. Comp. **23** (1969), 151–163.
63. ———, *Class number, a theory of factorization and genera*, 1969 Number Theory Institute, Proc. Sympos. Pure Math, vol. 20, Amer. Math. Soc., Providence RI, 1971, pp. 415–440.
64. ———, *The infrastructure of a real quadratic number field and its applications*, Proc. Number Theory Conf. (Univ. of Colorado, Boulder, Colo.), 1972, pp. 217–224.

65. ———, *On Gauss and composition I, II*, Number theory and applications (Banff, AB, 1988), NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci., 265, Kluwer Acad. Publ., Dordrecht, 1989, pp. 163–178, 179–204.
66. A. Silverberg, *Torsion points on abelian varieties of CM-type*, *Compositio Math.* **68** (1988), 241–249.
67. J. Silverman, *The arithmetic of elliptic curves*, Graduate Texts in Mathematics, vol. 106, Springer-Verlag, 1986.
68. H.P.F. Swinnerton-Dyer, *A brief guide to algebraic number theory*, London Mathematical Society Student Texts, vol. 50, Cambridge, University Press, Cambridge, 2001.
69. X.C. Tran, *Periodic continued fractions in function fields*, Ph.D. thesis, Macquarie University, 2000.
70. A.J. van der Poorten, *Some problems of recurrent interest*, Topics in classical number theory, Vol I, II (Budapest, 1981), *Colloq. Math. Soc. János Bolyai*, no. 34, North-Holland, Amsterdam, 1984, pp. 1265–1294.
71. ———, *An introduction to continued fractions*, Diophantine Analysis (Kensington, 1985) (J. Loxton and A.J. van der Poorten, eds.), London Math. Soc. Lecture Note Ser., vol. 109, Cambridge Univ. Press, Cambridge, 1986, pp. 99–138.
72. ———, *Explicit formulas for units in certain quadratic number fields*, Algorithmic number theory (Ithaca, NY, 1994), *Lecture Notes in Comput. Sci.*, vol. 877, Springer, Berlin, 1994, pp. 194–208.
73. ———, *Beer and continued fractions with periodic periods*, Number Theory (Ottawa, ON, 1996), CRM Proc. Lecture Notes, vol. 19, Amer. Math. Soc., Providence RI, 1999, pp. 309–314.
74. ———, *Reduction of continued fractions of formal power series*, Continued Fractions: From Analytic Number Theory to Constructive Approximation, (Columbia, MO, 1998), *Contemp. Math.*, no. 236, Amer. Math. Soc., Providence RI, 1999, pp. 343–355.
75. ———, *Non-periodic continued fractions in hyperelliptic function fields*, *Bull. Austral. Math. Soc.* **64** (2001), no. 2, 331–343.
76. ———, *Symmetry and folding of continued fractions*, *J. Théor. Nombres Bordeaux* **14** (2002), no. 2, 603–611.
77. ———, *A note on NUCOMP*, *Math. Comp.* **72** (2003), no. 244, 1935–1946.
78. ———, *Periodic continued fractions and elliptic curves*, High Primes and Misdemeanours: Lectures in honour of the 60th birthday of Hugh Cowie Williams (A.J. van der Poorten and A. Stein, eds.), Fields Institute Communications Series, Amer. Math. Soc., Providence RI, 2004, p. 13pp.
79. A.J. van der Poorten and X.C. Tran, *Periodic continued fractions in elliptic function fields*, Algorithmic Number Theory (Proc. Fifth International Symposium, ANTS-V, Sydney, 2002) (C. Fieker and D.R. Kohel, eds.), *Lecture Notes in Computer Science*, no. 2369, Springer, 2002, pp. 390–404.
80. A.J. van der Poorten and H.C. Williams, *On certain continued fraction expansions of fixed period length*, *Acta. Arith.* **89** (1999), no. 1, 23–35.
81. A. Weil, *Number Theory, an approach through history. From Hammurapi to Legendre*, Birkhäuser Boston, Inc., Boston, MA, 1984.
82. H.C. Williams, *A note on the period length of the continued fraction expansion of certain $\sqrt{D}$* , *Util. Math.* **28** (1985), 201–209.
83. ———, *Some generalisations of the S_n sequence of Shanks*, *Acta. Arith.* **69** (1995), no. 3, 199–215.
84. ———, *Édouard Lucas and primality testing*, Canadian Mathematical Society Series of Monographs and Advanced Texts, no. 22, John Wiley & Sons, Inc., New York, 1998.
85. ———, *Solving the Pell Equation*, Number theory for the millennium, III (Urbana, IL, 2000), A K Peters, Natick, MA, 2002, pp. 397–435.
86. Y. Yamamoto, *Real quadratic number fields with large fundamental units*, *Osaka. J. Math.* **8** (1971), 261–270.
87. J. Yu, *Arithmetic of hyperelliptic curves*, (1999), see 4–6, manuscript marked *Aspects of Mathematics*.